

REFERENCES

- Akbuga, J. , and Ermantas, K. Role of disintegrants in temperature- and moisture-related physical changes of ketoconazole tablets. Pharmazie 47 (1992) : 64-65.
- Aker, M.J. Antioxidants in pharmaceutical products. J. Pharm. Sci.Tech. 36 (1982) : 222-228.
- Alton, K.B. , Determination of the antifungal , Ketoconazole , in human plasma by high performance liquid chromatography. J. Chromatogr. 221 (1980) : 337-344.
- Badcock, N.R. Micro-determination of ketoconazole in plasma or serum by high-performance liquid chromatography. J. Chromatogr. 306 (1984) : 436-446.
- Banker, G.S. , and Rhodes, C.T. Modern Pharmaceutics. pp. 209-237. New York : Marcel Dekker Inc. 1990.
- Beak, P. , and Messer, W. Photorearrangements of some N-methyldiazoles. Tetrahedron . 25 (1960) : 328-3295.
- Berson, J.A. , and Brown, E. Studies on dihydropyridines II. The photochemical disproportionation of 4-(2-nitrophenyl)-1,4-dihydrpyridines. J. Am. Chem. Soc. 77 (1955) : 447-450.
- Borgers, M. and Van der Bossche, H. The mode of action of antifungal drugs in Levine (eds.) , Ketoconazole in the management of fungal disease , Sydney : ADIS Press , 1982.
- Cole, E.R. , Crank, G. , and Lye E. Photochemistry of heterocyclic II . Photolysis of benzimidazole and its 2-alkyl derivatives. Aust. J. Chem. 31 (1978) : 2675-2684.

- Conners, K.A. , Amidon, G.L. , and Stella, V.J. , eds. Chemical stability of pharmaceuticals : A handbook for pharmacists. 2 nd ed. pp. 82-114. New York : A Wiley-Interscience , 1986.
- Cooper, G. and Irwin , W.J. Photocyclisation of 1-Stylimidazoles. A novel route to N-bridgehead compounds. J. Chem. Soc. Perkin I. 75 (1976) : 75-80.
- Daneshmend, T.K. , and Warnock, D.W. Clinical pharmacokinetics of ketoconazole. Clinical Pharmacokinetics. 14 (1988) : 13-34.
- Dawson, B.A. ¹H- and ¹³C-NMR chemical shift assignment for ketoconazole. Canad. J. Spectr. 35 (1990) : 27-30.
- Felmeister, A. , and Dischler, C.A. Photodegradation of chlorpromazine hydrochloride. J. Pharm. Sci. 53 (1964) : 756-762.
- Gainsford, G.J., and Woolhouse, A.D. Photolysis of Imidazole N (3)-oxides. Formation of unsymmetrical benzil di-imines ; X-Ray crystal structure of one such di-imine. J. Chem. Soc., Chem. Commun. 20 (1978) : 857-858.
- Graybill, J.R. , Craven, P.C. Antifungal agents used in systemic mycoses activity and therapeutic use. Drugs 25 (1983) : 41-62.
- Haddadin,M.J.,Hawi,A.A.,andNazer,M.Z. Photolysis of some 1-hydroxybenzimidazole-3-oxide. Tetrahedron Lett. 46 (1978) : 857-858.
- Hamlin, W.E. , Chulski, T. , Johnson, R.H. , and Wagner, J.G. A note on the photolytic degradation of anti-inflammatory steroid. J. Am. Pharm. Assoc. Sci. 49 (1960) : 253-255.
- Heel, R.C. , Brogden, R.N. , Carmine, A. , Morley, P.A. , Speight, T.M., and Avery, G.S. Ketoconazole : A review of its therapeutic efficacy in superficial and system fungal infection. Drugs 23 (1982) : 1-36.

- Heeres, J. , Backx, L . J. J., Mostmans,J.H., Van Cutsem , J. Antimicotic Imidazoles . 4 . Synthesis and antifungal activity of ketoconazole , a new potent orally active broad spectrum antifungal agent . J. Med. Chem. 22 1979) : 1003 -1005.
- Henderson, W.A. ,and Zweig , A. Photolytic rearrangement and halogen dependent photocyclization of halophenylnaphthalene. J. Am. Chem. Soc. 89 (1967) : 6778-6779.
- Hey, D.H. , Jones, G.H. , and Perkins, M.J. Interneuclear cyclisation. Part XXVI. Photolysis of 2-iodo-N-methyl-benzanilide in benzene . J. Chem. Soc. (C) 1971 : 116-122.
- Holmes, S.E. , and Aldous, S. Stability of miconazole in peritoneal dialysis fluid. Am. J. Hosp. Pharm. 48 (1991) : 286-290.
- Kharach, N. , and Sharma, R.K. The photolysis of iodoaromatic compounds. Angew. Chem. 7 (1968) : 36-44.
- Kumer, K.P. , Okonomah, A.D. , Bradshaw, W.G. , and Soliman, K.F.A. Stability of ketocnazole in ethanolic solutions. Drug Dev. Ind. Pharm. 17 (1991) : 577-580.
- Lachman, L. , Lieberman, H.A. , and Kanig, J.L., eds. The Theory and Practice of Industrial Pharmacy. 3 rd ed. pp. 760-803. Philadelphia. Lea & Febiger , 1986.
- _____. , Swartz , C. J. , and Cooper, J. A comprehensive pharmaceutical stability testing laboratory III , A light stability cabinet for evaluating the photosensitivity of pharmaceuticals. J. Am. Pharm. Assoc Sci. 49(1960) : 213-218.
- Lin , S.L., and Lachman , L. Photochemical considerations of parenteral products. Bull. Parenter. Drug Assoc. 23 (1969) : 149-165.

- McEvoy, G.K. , ed. 1989. AHFS Drug Information. Bethesda , MD : American Society of Hospital Pharmacists , Inc. : 82-86 .
- Millican, J.A. , Ahuja, S. , and Cohen, J. Stability of pharmaceuticals. J. Pharm. Sci. 67 (1978) : 443-465.
- Moffat, A. C. , Jackson , J. V. , Moss, M.S. , and Widdop , B. Clarke's Isolation And Identification of Drugs. 2 nd ed . London : The Pharmaceutical Press , 1986
- Nijhoff, D.F. , and Havinga, N. Photoreaction of aromatic compounds VIII. A photochemical substitution of halogen in 2-bromo-(chloro)-4-nitroanizole. Tetrahedron Lett. 47 (1965) : 4199-4201.
- Nishikawa, M. , and Fujii, K. Effect of autoxidation of hydrogenated castor oil containing 60 oxyethylene groups on degradation of miconazole. Chem. Pharm. Bull. 39 (1991) : 2408-2411.
- Ogata,M.,Matsumoto,H.,Takahashi,S.,andKano,H.Organicphotochemical reaction VII. Photolysis of 1-Benzyl-2-ethylbenzylimidazole-3-oxide. Chem. Pharm. Bull. 18 (1970) : 964-969.
- Oyler, A.R. , Naldi, R.E. , Facchine, K.L. , and Burinsky, D.J. Characterization of autoxidation products of the antifungal compounds econazole nitrate and miconazole nitrate. Tetrahedron 47 (1991) : 6549-6560.
- Pinhey, J.T. ,and Rigby, R.D.G. Photo-reduction of chloro- and bromo-aromatic compounds. Tetrahedron Lett. 16 (1969) : 1267-1270.
- Robinson, G.E. , and Vernon, J.M. Photochemical phenylation and oxidation of halogen-substituted di-phenylmethanes in benzene J. Chem. Soc. (C) 1970 : 2586-2591.

Sane, R.T. , Tendolkar, R.V. , Gangal, D.P. , Ladage, K.D. , and Kolthorkar, R.M. An extraction colorimetric method for the determination of ketoconazole from pharmaceutical preparations. Ind. J. Pharm. Sci. Nov-Dec (1988) : 347-348.

Stewart, P.J. , and Tucker, I.G. Prediction of drug stability Part 3 : Oxidation and photolytic degradation. Aust. J. Hosp. Pharm. 15 (1985) : 111-117.

Swezey, S.F. , Giacomini, K.M. Abang, A. , Brass, A. , Stevens, D.A. , and Blaschke, T.F. Measurement of ketoconazole , a new antifungal agent , by high-performance liquid chromatography. J. Chromatogr. 227 (1982) : 510-515.

The United States Pharmacopoeia XXII. National Formulary XVII.
Rockville , MD : The United States Pharmacopeial Convention , Inc. 1990 : 747-748.

Vargas, F. , Rivas, C. , and Canudas, N. Formation of a perbenzoic acid derivative in the photodegradation of fenofibrate : phototoxicity studies on erythrocytes. J. Pharm. Sci. 82 (1993) : 590-591.


Wad , N., Degradation of clonazepam in serum by light confirmed by means of a high performance liquid chromatographic method. Ther. Drug Monit. 8 (1986) : 358-360.

Wilson, S. , and Ruenitz, P.C. Structure characterization and biological effects of photocyclized products of tamoxifen irradiation. J. Pharm. Sci. 82 (1993) : 571-574.



APPENDICES

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย



Silica gel GF 254 / solvent system ,chloroform : methanol (95 : 5)

= fluorescent spot under UV light (254 nm) and
positive with iodine vapour

Figure 3: TLC chromatogram of ketoconazole in various solvents on photolysis at 15 hrs.

- a) in ethylacetate b) in chloroform
- c) in acetone d) in methanol
- e) in chloroform at 0 hr.




Figure 4 :The UV spectrum of ketoconazole (in CHCl_3)




Figure 5 : The IR spectrum of ketoconazole (KBr disc)




Figure 6 : the EIMS spectrum of Ketoconazole




Figure 7 : The 500 MHz ^1H - NMR spectrum of ketoconazole in chloroform - d (Full spectrum)




Figure 8 : The 500 MHz ^1H - NMR spectrum of ketoconazole in chloroform - *d* (Expansion between 2.1 - 3.9 ppm)




Figure 9 : The 500 MHz ^1H - NMR spectrum of ketoconazole in chloroform - *d* (Expansion between 5.9 - 7.6 ppm)




Figure 10 : The 125 MHz ^{13}C - NMR spectrum of ketoconazole in chloroform - d




Figure 11 : The UV spectrum of compound A (in CHCl_3)




Figure 12 : The IR spectrum of compound A (KBr disc)




Figure 13 : the EIMS spectrum of compound A.




Figure 14 : The 500 MHz ^1H - NMR spectrum of compound A in chloroform - d (Full spectrum)




Figure 15 : The 500 MHz ¹H - NMR spectrum of compound A in chloroform - *d* (Expansion between 2.1 - 4.4 ppm)




Figure 16 : The 500 MHz ^1H - NMR spectrum of compound A in chloroform - d (Expansion between 4.7 - 8.1 ppm)




Figure 17 : The 125 MHz ¹³C - NMR spectrum of compound A in chloroform - d




Figure 18 : The ¹³C -DEPT spectrum of compound A in chloroform - d

a) ¹³C DEPT - 135

b) Normal spectrum




Figure 19 : The 500 MHz ¹H - ¹H COSY spectrum of compound A in chloroform - d (Full spectrum)




Figure 20 : The 500 MHz ^1H - ^1H COSY spectrum of compound A in chloroform - d
 (Expansion between 1.0 - 4.0 ppm)




Figure 21 : The 500 MHz ^1H - ^1H COSY spectrum of compound A in chloroform - d
 (Expansion between 3.9 - 4.9 ppm)




Figure 22 : The 500 MHz ^1H - ^1H COSY spectrum of compound A in chloroform - d
 (Expansion between 6.8 - 8.2 ppm)




Figure 23 : The 125 MHz ^{13}C - ^1H COSY spectrum of compound A in chloroform - d (Full spectrum)




Figure 24 : The 125 MHz ^{13}C - ^1H COSY spectrum of compound A in chloroform - d
 (Expansion : X scale between 20-80 ppm and Y scale between 2.0 - 4.8 ppm)




Figure 25 : The 125 MHz ^{13}C - ^1H COSY spectrum of compound A in chloroform - d
 (Expansion : X scale between 100 - 140 ppm and Y scale between 6.6 - 8.4 ppm)




Figure 26 : The 500 MHz COLOC spectrum (at 4 Hz) of compound A in chloroform - *d* (Full spectrum)





Figure 27 : The 500 MHz COLOC spectrum of compound A (at 4 Hz) in chloroform - d
 (Expansion : X scale between 60 -74 ppm and Y scale between 4.0 - 4.7 ppm)






Figure29 :The 500 MHz COLOC spectrum (at 8 Hz) of compound A in chloroform - d (Full spectrum)




Figure 30 : The 500 MHz COLOC spectrum of compound A (at 8 Hz) in chloroform - *d*
 (Expansion : X scale between 35 - 60 ppm and Y scale between 2.9 -3.9 ppm)




Figure 31 : The 500 MHz COLOC spectrum of compound A (at 8 Hz) in chloroform - *d*
 (Expansion · X scale between 99 - 135 ppm and Y scale between 4.0 - 4.6 ppm)




Figure 32 : The 500 MHz COLOC spectrum of compound A (at 8 Hz) in chloroform - *d*

(Expansion : X scale between 110 - 160 ppm and Y scale between 6.7 - 8.4 ppm)




Figure 33 : The 500 MHz COLOC spectrum (at 12 Hz) of compound A in chloroform - d (Full spectrum)




Figure 34 : The 500 MHz COLOC spectrum of compound A (at 12 Hz) in chloroform-*d*

(Expansion : X scale between 39-110 ppm and Y scale between 3.7 - 4.8 ppm)




Figure 35: The 500 MHz COLOC spectrum of compound A (at 12 Hz) in chloroform - *d*
 (Expansion : X scale between 112 - 158 ppm and Y scale between 6.7 - 7.7 ppm)



VITA

Mr. Paiboon Nuntanakorn was born on February 1, 1969 , in Rachaburi ,Thailand. He graduated with a Bachelor degree of Pharmacy from Faculty of Pharmaceutical Science, Chulalongkorn University , Bangkok in 1992.

ศูนย์วิทยบริพยากร
จุฬาลงกรณ์มหาวิทยาลัย