

References

- Aloya VJ and Walker RF. Alpha adrenergic control of serotonin release from rat pineal glands. Neuroendocrinology 48 (1988): 1; 61-66.
- Aoyagi N, Kimura R and Murata T. P. *P. incarnata* dry extract I. Isolation of maltol and pharmacological action of maltol and ethyl maltol. Chemical and Pharmaceutical Bulletin. 22 (1974): 1008-1013.
- Battery EJ and Newman EV. Chronotropic effects of norepinephrine and tyramine: A system response analysis. Circulation Research 15 (1964): 112.
- Bergner P. Passiflora: Passionflower. Medical herbalism 7 (2001): (1&2); 13-14, 26.
- Borne RF. Serotonin: The neurotransmitter for '90s. Drug Topics (1994): 108.
- Caughey B and Kirshner N. Effects of reserpine and tetrabenazine on catecholamine and ATP storage in cultured bovine adrenal medullary chromaffin cells. Journal of Neurochemistry. 49 (1987); 2: 563-573.

Chopra RN, Nayar SL, Chopra IC. Glossary of Indian Medicinal plants CSIR, New delhi, Indian (1956): 186-187.

Cohen HW, Gibson G and Alderman MH. Excess risk of myocardial infarction in patients treated with antidepressant medications: association with use of Tricyclic agents. AM J Med 108 (2000): 87-88.

Copstead LC and Banasik JL. Pathophysiology 3rd ed. St. Louis: Elsevier Saunders (2005).

Davy M, Grandcourt O, Midol-Munnet M, and Cohen Y. 5-Hydroxytryptamine and β -adrenoceptors in rat isolated atria. Cell. Mol. Sci. 53(1997): 263-266.

Dedeogin A and Fisher LA. Central nervous actions of serotonin and a serotonin 1A receptor agonist: Cardiovascular excitation at low doses. American Society for Pharmacology and Experimental therapeutics 257(1991): 1; 425-432.

Dhawan K, Dhawan S, and Sharma A. *Passiflora*: a review update. J Ethnopharmacol 94(2004): 1-23.

Dhawan K, Kumar S, and Sharma A. Anxiolytic activity of aerial and underground parts of *Passiflora incarnata*. Fitoterapia 72(2001): 922-926.

Fawaz G and Simaan J. On the mechanism of the presence action of tyramine. Neunyn-Schmicdeberg's Archives of Pharmacology. 247 (1964): 5; 456-460.

Fernandaz A, Bang SE, Srivathsan K, Vieweg WVR. Cardiovascular side effects of newer antidepressants. Anadolu Kardiyol Derg. (2007); 7: 305-309.

Formica JV and Regelson W. Review of the biology of quercetin and related bioflavonoids. Food Chem Toxicol. 33 (1995): 1061-1080.

Friccione GL, Woznicki RM, Klesmer J and Vlay SC. Vasoconstrictive effects and SSRIs. J Clin Psychiatry 54 (1993): 71-72.

Glassman AH. Cardiovascular effects of antidepressant drugs: updated. J Clin Psychiatry. 59 (1998): 15; 13-18.

Itoigawa M, Kazumi T, Chihiro I and Hirochi F. Structure-activity relationship of cardiotonic flavonoids in guinea-pig papillary muscle. Journal of Ethnopharmacology. 65(1999): 267-272.

Jiang W, and Davidson JRT. Antidepressant therapy in patients with ischemic heart disease. Am Heart J. (2005); 150: 871-881.

Jirajariyaweg Waraporn. Study the positive chronotropic and inotropic effects of serotonin on the isolated right and left atria. Master's thesis, Department of Pharmaceutical Science, Graduate School, Chulalongkorn University, 1988.

Katzung B.G., Trevor A.J., and Masters S.B. Katzung and Trevor's Pharmacology. 7th ed. New York. McGraw-Hill Medical (2004)..

Kaumann AJ and Levy FO. 5-Hydroxytryptamine receptors in the human cardiovascular system. Pharmacology & Therapeutics 111 (2006); 3: 674-706.

Klabunde RE. Cardiovascular Pharmacology Concepts: Excitation-contraction coupling. Available from ; <http://www.cypharmacology.com>. [Mar 2007]

Kirtikar KR, Basu BD. Indian Medicinal Plants Periodical Experts. Dhradun (1975): 1103.

Kranias RE. Cardiac function: A simple view. University of Cincinnati. Available from: http://www.med.uc.edu/kranias/cardiac_function.thm. [Feb 2005]

Maluf E, Barros HMT, Frochtengarten ML, Benti R and Leite JR. Assessment of the hypnotic/sedative effects and toxicity of *Passiflora edulis* aqueous extract in rodents and humans. Phytotherapy Research 5 (1991): 262-266.

Medina JH, Paladini AC, Wolfman C, LevideStein M, Calvo D, Diaz LE and Pena C. Chrysin (5, 7-di-OH-flavone) a naturally occurring ligand for benzodiazepine receptors, with anticonvulsant properties. Biochemistry and Pharmacology. 40 (1990): 2227-2231.

Moongkarndi P, Kosem N, Luanratana O, Jongsomboonkusol S and Pongpan N. Antiproliferative activity of Thai medicinal plant extracts on human breast adenocarcinoma cell line. Fitoterapia 75(2004): 375-377.

Nishio H, Morimoto Y, Hisaoka K, Nakata Y and Watanabe T. 5-HT-induced, 5-HT₃ receptor-mediated, and ruthenium red- and capsaicin-sensitive positive chronotropic effects in the isolated guinea pig atrium. The Japanese Journal of Pharmacology. 89 (2002); 3: 242-248.

Nwosu MO. Herbs for mental disorders. Fitoterapia 70 (1999): 58-63.

Petry RD, Reginatto F, de-Paris F, Gosmann G, Salgueiro JB, Quevedo J, Kapezinski F, Ortego GG and Schenkel EP. Comparative pharmacological study on hydroethanol extracts of *Passiflora alata* and *Passiflora edulis* leaves. Phytotherapy Research 15 (2001): 162-164.

Ramussen SL, Overo KF, and Tanghoj P. Cardiac safety of citalopram: Prospective trials and retrospective analyses. J Clin Psychopharmacol 19(1999): 407-415.

Rang HP, Dale MM and Ritter JM. Other peripheral mediators: 5-hydroxytryptamine and purines. Pharmacology 4th ed. Edinburgh, UK. Harcourt Publisher Ltd. (2001):165-176.

Rodriquez de la Torre B, Dreher J, Malevany I, Bagli M, Kolbinger M, Omran H, Lüderitz B and Rao ML. Serum levels and cardiovascular effects of triacyclic antidepressants and selective serotonin reuptake inhibitor in depressed patients. Ther Drug Monit 23 (2001): 4; 435-440.

Roose SP, Glassman AH, Attia E, Woodring S, Giardina EV and Bigger JT. Cardiovascular effects of fluoxetine in depressed patients with heart disease. AM J Psychiatry 155(1998): 660-665.

Roose SP and Spatz E. Treating depression in patients with ischemic heart disease.

Which agents are best to use and to avoid? Drug Saf 20(1999): 459-465.

Sharma A, Dhawan K, and Kumar S. Anxiolytic activity of aerial and underground parts

of Passiflora incarnata. Fitoterapia 72(2001): 922-926.

Speroni E and Minghetti A. Neuropharmacological activity of extract from *P. incarnata*.

Planta Medica. 54 (1988): 488-491.

Strik JJMH, Honig A, Lousberg R, Lousberg AHP, Cheriex EC, Tuynman-Qua HG,

Kuijpers PMJC, Wellens HJJ and Praag HMV. Efficacy and safety of fluoxetine in

the treatment of patients with major depression after first myocardial infarction:

finding from a double-blinded placebo-controlled trial. Psychosom Med

62(2000): 783-789.

Tellioghr T and Roberson D. Genetic or aquired deficit in the norepinephrine

transposters: current understanding of clinical implications. Expert Reviews in

Molecular Medicine. Available from: <http://www.expertreviews.org/> [Nov 2001].

Thanacoody HK and Thomas SH. Tricyclic antidepressant poisoning: cardiovascular

toxicity. Toxicological reviews 24 (2005): 3; 205-214.

Trendelburg U. The action of histamine and 5-hydroxytryptamine on isolated mammalian atria. J Pharmaco Exp Ther. 130 (1996): 450-460.

Van den Buuse and Wegener N. Involvement of serotonin 1A receptors in cardiovascular responses to stress: a radio-telemetry study in four rat strains. European Journal of Pharmacology 507(2005): 187-198.

Wijagkanalan Panu. Antidepressant activity of extracts of *Passiflora foetida* in mice and the involvement of D₁ and 5-HT_{1A} receptors. Master's thesis, Department of Pharmaceutical Science, Graduate School, Chulalongkorn University, 2005.

Wouters W, Tulp M T.M., and Beran P. Flesinoxan lowers blood pressure and heart rate in cat via 5-HT_{1A} receptors. Eur J Pharmacol 149(1988): 3; 213-223.

Yamoto T, Aomine M, Noto H, Ikeda M and Ohta C. Capsaicin does not inhibit the intracellular calcium handing process in rat ventricular papillary muscle. Gen Pharmaco. 27 (1996); 1: 105-108 (4).

Yildiz A, Gönül AS, and Tamam L. Mechanism of action of antidepressant: Beyond the receptors. Bull Clin Psychopharmacol 12 (2002): 194-200.

Zanoli P, Avallone R and Baraldi M. Behavioral characterization of the flavonoids apigenin and chrysin. Fitoterapia. 71 (2000): S117-S123.

Zibadi S and Watson RR. Passion fruit (*Passiflora edulis*) composition, efficacy, and safety. Evid Based Integrative Med 1(2004): 3; 183-187.

APPENDICES

Table 2 Chemical compound for Krebs-Henseleit Solution (mM/L)

Chemicals	mM
Sodium chloride (NaCl)	118.0
Potassium chloride (KCl)	4.70
Magnesium sulfate ($MgSO_4 \cdot 7H_2O$)	1.64
Calcium chloride ($CaCl \cdot 2H_2O$)	2.52
Sodium bicarbonate ($NaHCO_3$)	24.88
Potassium dibasic phosphate (KH_2PO_4)	1.18
Glucose	11.10

Table 3 Effects of DMSO (0.25 % v/v) on rate and force of contraction on isolated left and right atria (n=6) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	94.92 \pm 2.79	97.16 \pm 1.59	99.83 \pm 0.60
3 min	91.95 \pm 3.66	96.86 \pm 1.78	99.22 \pm 0.55
5 min	91.80 \pm 5.04	96.77 \pm 1.94	100.21 \pm 1.28
10 min	86.71 \pm 4.19	94.98 \pm 2.78	98.36 \pm 1.05
15 min	86.47 \pm 4.78	95.78 \pm 2.92	98.79 \pm 1.10

Table 4 Effects of PF002-(1-4) (100 μ g/ml) on rate and force of contraction on isolated left and right atria (n=6) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	93.59 \pm 2.68	89.80 \pm 5.31	98.32 \pm 0.82
3 min	92.38 \pm 4.10	89.93 \pm 6.06	100.24 \pm 1.79
5 min	92.32 \pm 4.92	93.58 \pm 3.43	101.70 \pm 2.74
10 min	95.02 \pm 6.40	88.88 \pm 4.37	103.42 \pm 3.70
15 min	92.17 \pm 6.21	87.56 \pm 6.00	104.13 \pm 3.94

Table 5 Effects of PF002-(5-6) (100 µg/ml) on rate and force of contraction on isolated left and right atria (n=6) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min	91.23±3.63*	88.20±4.32	98.30±0.70
3 min	92.02±4.00	90.07±5.57	99.23±1.30
5 min	91.92±3.93	88.05±6.28	100.44±1.82
10 min	94.60±2.50	86.42±7.19	100.05±2.13
15 min	94.09±2.17	84.38±7.54	100.12±1.90

*p<0.05, significantly different from DMSO 0.25% group (unpaired *t*-test)

Table 6 Effects of PF002-7 (100 µg/ml) on rate and force of contraction on isolated left and right atria (n=6) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min	81.10±3.16*	85.00±4.04	96.46±2.21
3 min	90.35±4.91	92.66±5.47	95.81±2.37
5 min	95.24±5.10	94.62±5.96	95.78±2.64
10 min	100.87±5.22	97.32±5.59	93.66±2.63*
15 min	100.72±5.35	96.17±7.26	90.05±2.79*

*p<0.05, significantly different from DMSO 0.25% group (unpaired *t*-test)

Table 7 Effects of PF003-1 (100 µg/ml) on rate and force of contraction on isolated left and right atria (n=6) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min	96.55±1.98	92.67±1.50	96.44±2.15
3 min	95.78±3.31	92.66±2.54	97.91±1.18
5 min	94.65±2.80	88.23±3.76	98.98±0.91
10 min	95.06±3.24	84.29±3.25*	99.69±1.00
15 min	90.76±3.67	81.49±4.01*	99.52±1.35

*p<0.05, significantly different from DMSO 0.25% group (unpaired *t*-test)

Table 8 Effects of PF003-2 (100 µg/ml) on rate and force of contraction on isolated left and right atria (n=6) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min	84.48±1.22**	101.17±3.93	99.42±1.01
3 min	108.76±4.01*	103.69±8.44	98.49±0.59
5 min	113.20±3.18**	108.20±6.32	98.41±0.20
10 min	119.48±2.90**	111.99±4.66*	99.18±0.40
15 min	119.22±4.95**	115.41±3.47*	99.62±0.62

*p < 0.05, **p<0.001, significantly different from DMSO group (unpaired *t*-test)

Table 9 Effects of PF003-(3-5) (100 µg/ml) on rate and force of contraction isolated on left and right atria (n=6) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min	88.05±4.53	95.76±4.75	92.84±1.47
3 min	91.53±3.27	113.58±8.66	90.39±1.53*
5 min	93.51±1.96	115.83±7.59	87.97±1.35*
10 min	92.80±4.13	116.32±10.82	86.05±1.00*
15 min	95.24±4.77	110.15±8.31	85.13±1.17*

*p<0.05, significantly different from DMSO 0.25% group (unpaired *t*-test)

Table 10 Effects of NE (0.1 µM) on rate and force of contraction on isolated left and right atria (n=4) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min	165.91±11.40*	148.11±15.14*	132.22±6.30*
3 min	182.41±13.65*	125.93±13.85	133.69±6.06*
5 min	179.60±18.13*	125.33±12.81	134.26±6.33*
10 min	166.62±19.59*	113.48±8.97	132.49±6.53*
15 min	151.88±20.34*	112.02±12.05	130.47±6.48*

*p<0.05, significantly different from response at 0 min (paired *t*-test)

Table 11 Effects of propranolol (10 μ M) on rate and force of contraction isolated on left and right atria (n=6) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	95.76 \pm 0.86**	100.36 \pm 5.80	87.85 \pm 1.24*
3 min	89.99 \pm 1.67**	107.77 \pm 3.62	81.94 \pm 1.69*
5 min	85.10 \pm 2.80**	110.36 \pm 3.20*	79.81 \pm 1.98*
10 min	82.72 \pm 5.61**	114.03 \pm 3.45*	76.57 \pm 2.85*
15 min	84.86 \pm 3.35**	109.70 \pm 3.74*	74.99 \pm 3.53*

*p<0.05, ** p<0.001, significantly different from response at 0 min (paired *t*-test)

Table 12 Effects of NE (0.1 μ M) in the presence of propranolol (10 μ M) on rate and force of contraction on isolated left and right atria (n=4) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min.	100.22 \pm 2.68*	75.63 \pm 12.97*	97.20 \pm 2.28*
3 min.	96.84 \pm 5.34*	72.33 \pm 16.44*	95.98 \pm 2.88*
5 min.	96.94 \pm 4.90*	71.87 \pm 17.74*	95.34 \pm 4.31*
10 min.	85.37 \pm 2.50*	67.26 \pm 13.88*	92.94 \pm 4.66*
15 min.	83.64 \pm 3.33*	60.73 \pm 14.54*	89.34 \pm 2.18*

*p < 0.05, significantly different from NE group in the absence of propranolol (unpaired *t*-test)

Table 13 Effects of PF003-2 (100 µg/ml) in the presence of propranolol (10 µM) on rate and force of contraction on isolated left and right atria (n=6) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min.	93.58±0.84*	95.33±0.80	96.77±0.63*
±3 min.	91.71±2.90*	99.81±2.94	94.87±0.96*
5 min.	92.76±4.04*	93.89±4.55	94.15±1.33*
10 min.	93.79±4.35*	101.06±10.31	93.72±2.05*
15 min.	92.38±5.42*	102.11±12.64	93.20±1.88*

* $p < 0.05$, significantly different from PF003-2 group in the absence of propranolol (unpaired *t*-test)

Table 14 Effects of serotonin (1 µM) on rate and force of contraction on isolated left and right atria (n=4) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min	116.66±1.54*	113.84±2.27*	106.21±2.50
3 min	123.26±2.59*	108.60±4.17	108.77±3.14
5 min	123.32±1.74*	104.55±4.45	109.43±2.80
10 min	117.96±2.93*	102.41±5.71	107.40±2.60
15 min	112.78±2.46*	97.99±6.82	106.62±2.70

* $p<0.05$, significantly different from response at 0 min (paired *t*-test)

Table 15 Effects of ketanserin (10 μ M) on rate and force of contraction on isolated left and right atria (n=6) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	95.62 \pm 1.43*	93.10 \pm 6.90	96.98 \pm 0.57*
3 min	92.70 \pm 0.87*	93.26 \pm 4.20	93.99 \pm 0.72*
5 min	91.46 \pm 0.72*	90.21 \pm 5.87	91.24 \pm 1.11*
10 min	90.34 \pm 1.95*	95.96 \pm 2.94	85.53 \pm 1.92*
15 min	88.46 \pm 2.60*	93.08 \pm 2.40	87.30 \pm 2.29*

*p<0.05, significantly different from response at 0 min (paired *t*-test)

Table 16 Effects of serotonin (10 μ M) in the presence of ketanserin (10 μ M) on isolated rate and force of contraction of left and right atria (n=4) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min.	97.59 \pm 0.81**	96.94 \pm 1.29*	97.18 \pm 0.62*
3 min.	98.87 \pm 0.73*	99.92 \pm 3.80	100.97 \pm 1.94
5 min.	96.29 \pm 0.58**	95.65 \pm 1.08	103.26 \pm 3.11
10 min.	94.45 \pm 1.02**	95.10 \pm 3.24	108.71 \pm 5.93
15 min.	92.99 \pm 1.70*	99.43 \pm 5.79	111.98 \pm 6.91

*p < 0.05, ** p<0.001, significantly different from serotonin group (unpaired *t*-test)

Table 17 Effects of PF003-2 (100 µg/ml) in the presence of ketanserin (10 µM) on isolated rate and force of contraction of left and right atria (n=6) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min.	85.16±3.07	95.62±0.85	97.11±0.40
3 min.	92.14±2.09*	109.34±2.22	97.60±1.30
5 min.	94.18±1.84**	111.16±2.52	98.12±1.84
10 min.	97.44±1.92**	113.33±2.23	100.21±2.30
15 min.	97.83±1.52*	114.42±3.23	98.33±1.55

*p < 0.05, **p < 0.001 significantly different from PF003-2 group (unpaired t-test)

Table 18 Effects of PF003-2 (100 µg/ml) in the presence of propranolol (10 µM) and ketanserin (10 µM) on rate and force of contraction on isolated left and right atria (n=5) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min.	80.56±1.42	96.23±2.68	93.21±1.08*
3 min.	91.38±2.31*	98.96±2.55	93.08±2.22*
5 min.	97.32±2.38*	105.39±3.44	93.16±2.74
10 min.	99.61±3.03*	101.63±2.28	93.51±3.80
15 min.	93.74±1.64*	101.96±2.18	94.70±4.77

*p < 0.05, significantly different from PF003-2 group in the absence of ketanserin and propranolol (unpaired t-test)

Table 19 Effect of caffeine (10 mM) and PF003-2 (100 µg/ml) on the relationships between the Ti / Tss and the rest interval of range 10 to 300 seconds (n=4) (mean ± SEM).

Rest interval second	Post-rest contraction (PRC)			
	control	caffeine	control	PF003-2
10	2.64 ± 0.16	1.15 ± 0.07*	2.64 ± 0.16	3.15 ± 0.16
20	2.70 ± 0.11	1.33 ± 0.87*	2.70 ± 0.11	3.18 ± 0.26
30	2.66 ± 0.11	1.39 ± 0.07*	2.66 ± 0.11	3.00 ± 0.28
40	2.50 ± 0.13	1.60 ± 0.12*	2.50 ± 0.13	3.07 ± 0.39
50	2.34 ± 0.24	1.67 ± 0.21	2.34 ± 0.24	2.94 ± 0.48
60	2.30 ± 0.21	1.82 ± 0.18	2.30 ± 0.21	2.92 ± 0.43
120	2.22 ± 0.08	1.84 ± 0.27	2.22 ± 0.08	2.95 ± 0.41
300	2.06 ± 0.03	1.95 ± 0.26	2.06 ± 0.03	2.71 ± 0.40

*p< 0.05, significantly different from control group (paired *t*-test)

Table 20 Effects of tyramine (10 μ M) on rate and force of contraction on isolated left and right atria (n=4) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	287.47 \pm 29.30*	195.22 \pm 32.13	143.32 \pm 4.29*
3 min	304.22 \pm 25.93*	165.35 \pm 28.18	150.88 \pm 13.68*
5 min	287.28 \pm 23.69*	135.98 \pm 25.50	163.10 \pm 3.54*
10 min	278.50 \pm 25.28*	116.01 \pm 18.02	166.00 \pm 5.83*
15 min	262.84 \pm 19.95*	108.39 \pm 16.55	160.27 \pm 8.72*

*p<0.05, significantly different from response at 0 min (paired *t*-test)

Table 21 Effects of tyramine (10 μ M) on rate and force of contraction on isolated left and right atria in rat pretreated with reserpine (5 mg/kg. i.p. for 5 days)(n=4) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	99.90 \pm 1.15*	105.01 \pm 4.58*	106.12 \pm 3.23*
3 min	101.29 \pm 2.23*	100.60 \pm 3.22*	108.91 \pm 3.86*
5 min	99.54 \pm 2.04*	97.52 \pm 2.66*	109.22 \pm 3.99*
10 min	95.31 \pm 1.78*	94.56 \pm 2.25	108.26 \pm 3.53*
15 min	93.06 \pm 1.62*	91.87 \pm 1.87	107.10 \pm 3.19*

*p < 0.05, significantly different from tyramine group in normal rats (unpaired *t*-test)

Table 22 Effects of PF003-2 (100 µg/ml) on rate and force of contraction on isolated left and right atria in rat pretreated with reserpine (5 mg/kg. i.p. for 5 days)(n=4) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min	92.92±1.30*	98.53±2.12	95.92±0.62*
3 min	104.17±2.09	109.65±2.53	97.36±0.79
5 min	108.30±2.78	108.48±1.72	99.75±0.99
10 min	112.12±4.10	109.60±3.13	103.88±1.46*
15 min	108.46±2.79	107.49±4.29	106.81±1.55*

*p < 0.05, significantly different from PF003-2 group in normal rat (unpaired *t*-test)

Table 23 Effects of NE (0.1 µM) on rate and force of contraction on isolated left and right atria in 5 min (n=6) (mean ± SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100±0.00	100±0.00	100±0.00
1 min	152.82±5.18**	131.24±8.14*	139.50±5.00*
2 min	156.80±3.99**	107.15±5.45	146.02±5.42**
3 min	156.39±3.94**	99.78±4.70	148.25±5.53**
4 min	156.08±4.70**	94.78±5.38	146.16±5.61**
5 min	155.07±5.23**	90.66±5.14	149.06±5.75**

*p < 0.05, **p < 0.001, significantly different from response at 0 min (student's *t*-test)

Table 24 Effects of NE (0.1 μ M) on rate and force of contraction on isolated left and right atria (5 min) in the presence of amitriptyline(10 μ M) ((n=6) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	163.92 \pm 17.97	140.05 \pm 11.53	133.30 \pm 8.38
2 min	155.74 \pm 16.73	107.62 \pm 8.93	141.65 \pm 9.27
3 min	148.26 \pm 16.15	102.36 \pm 9.62	143.56 \pm 9.31
4 min	143.81 \pm 16.07	96.98 \pm 8.47	142.12 \pm 9.47
5 min	139.54 \pm 14.86	99.16 \pm 9.07	139.82 \pm 9.60

Table 25 Effects of NE (0.1 μ M) on rate and force of contraction on isolated left and right atria (5 min) in the presence of PF003-2 (100 μ g/ml) ((n=6) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	107.34 \pm 2.04*	113.50 \pm 3.87	106.95 \pm 2.60*
2 min	117.46 \pm 4.29*	118.43 \pm 6.10	117.97 \pm 4.96*
3 min	123.44 \pm 4.04	118.03 \pm 7.58*	124.48 \pm 5.50
4 min	129.78 \pm 4.42	118.29 \pm 6.20*	127.97 \pm 5.78
5 min	134.65 \pm 5.48	115.06 \pm 8.07*	130.33 \pm 6.29

* $p < 0.05$, significantly different from NE group (unpaired *t*-test)

Table 26 Effects of serotonin (1 μ M) on rate and force of contraction on isolated left and right atria (5 min) (n=4) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	120.05 \pm 1.78**	109.16 \pm 1.78*	102.35 \pm 0.30*
2 min	130.08 \pm 3.00*	107.90 \pm 1.82*	102.68 \pm 0.22*
3 min	132.76 \pm 3.65**	109.82 \pm 3.27*	102.57 \pm 0.24*
4 min	136.61 \pm 4.01*	109.02 \pm 1.96*	102.15 \pm 0.33*
5 min	133.00 \pm 4.88*	107.79 \pm 1.40*	101.94 \pm 0.42*

*p < 0.05, **p < 0.001, significantly different from response at 0 min (paired t-test)

Table 27 Effects of fluoxetine (1 μ M) on rate and force of contraction on isolated left and right atria (5 min) (n=4) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	94.91 \pm 1.72*	95.72 \pm 1.63	97.57 \pm 0.64*
2 min	92.64 \pm 2.70*	98.46 \pm 3.97	96.53 \pm 0.66*
3 min	90.64 \pm 2.84*	95.74 \pm 2.50	95.27 \pm 0.98*
4 min	90.02 \pm 2.46*	97.96 \pm 2.77	94.62 \pm 0.97*
5 min	89.00 \pm 2.01*	96.25 \pm 2.43	94.14 \pm 1.14*

*p < 0.05, significantly different from response at 0 min (paired t-test)

Table 28 Effects of serotonin (1 μM) on rate and force of contraction on isolated left and right atria (5 min) in the presence of fluoxetine (1 μM) ((n=4) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	111.60 \pm 2.93*	103.45 \pm 2.08	100.81 \pm 0.34*
2 min	120.59 \pm 1.43*	98.34 \pm 4.37	100.64 \pm 0.44*
3 min	124.64 \pm 0.82	104.73 \pm 2.94	100.42 \pm 0.45*
4 min	126.90 \pm 1.56	105.70 \pm 3.59	99.80 \pm 0.56*
5 min	128.01 \pm 0.50	108.59 \pm 6.53	99.60 \pm 0.59*

*p < 0.05, significantly different from serotonin group (unpaired *t*-test)

Table 29 Effects of serotonin (1 μM) on rate and force of contraction on isolated left and right atria (5 min) in the presence of PF003-2 (100 $\mu\text{g}/\text{ml}$) ((n=4) (mean \pm SEM).

Time	Response (%)		
	Left force	Right force	Right rate
Control	100 \pm 0.00	100 \pm 0.00	100 \pm 0.00
1 min	109.86 \pm 0.99	100.08 \pm 2.02	101.23 \pm 0.69
2 min	115.46 \pm 3.13	103.27 \pm 3.53	102.25 \pm 0.98
3 min	116.90 \pm 3.68	106.42 \pm 6.66	103.68 \pm 1.40
4 min	120.15 \pm 2.56	103.10 \pm 4.21	104.38 \pm 1.51
5 min	124.07 \pm 2.90	104.08 \pm 4.96	104.86 \pm 1.74

CURRICULUM VITAE

Miss Salinee Sirodom was born in October 23, 1977 in Bangkok, Thailand. She graduated with a Bachelor of Science (Pharmacy) in 2002 from the Faculty of Pharmaceutical Science, Mahidol University, Thailand. After graduation, she worked as a pharmacist in Bangplee Hospital, Samutprakarn, Thailand, for 3 years.