

REFERENCES

- Aida, M., Shinomiya, K., Hano, Y. and Nomura, T. 1993. Artonins J, K and L, three new isoprenylated flavones from the root bark of *Artocarpus heterophyllus* Lamk. Heterocycles 36: 575-583.
- Aida, M., Shinomiya, K., Matsuzawa, K., Hano, Y. and Nomura, T. 1994. Artonins Q, R, S, T, and U, five new isoprenylated phenols from the bark of *Artocarpus heterophyllus* Lamk. Heterocycles 39: 847-858.
- Altman, L. J. and Zito, S. W. 1976. Sterols and triterpenes from the fruit of *Artocarpus altilis*. Phytochemistry 15: 829-830.
- An, B. J., Choi, J. Y., Kwon, I. B., Nishioka, I. and Choi, C. 1992. Structure and isolation of glucosyl transferase inhibitor from jackfruit. Han'guk Saenghwa Hakhoechi 25: 347-353. (CA. 118: 87474j).
- Arora, J. S., Sanhu, R. S., Kamboj, S. S. and Chopra, S. K. 1987. Occurrence and characterization of lympho-agglutinins in Indian plants. Vox Sang 52: 134-137. (CA. 106: 211018g).
- Ayensu, E. S. 1981. Medicinal Plants of the West Indies. U. S. A.: Reference Publication.
- Barik, B. R., Bhaumik, T., Dey, A. K. and Kundu, A. B. 1994. Triterpenoids from *Artocarpus heterophyllus*. Phytochemistry 35: 1001-1004.
- Barton, D. H. R. 1951. Cycloartenone, a triterpenoid ketone. Ibid.: 1444-1451.
- Britton, G. 1983. The biochemistry of natural pigments. London: Cambridge University Press. pp. 222-239.
- Chakraborty, D. P. and Mandal, A. K. 1981. Aurantinamide acetate from *Artocarpus integrifolia* Linn. J. Indian Chem. Soc. 58: 103. (CA. 94:153452q).
- Chatterjee, B. P., Sarkar, N. and Rao, A. S. 1982. Serological and chemical investigations of the anomeric configuration of the sugar units in the D-galacto-D-mannan of fenugreek (*Trigonella foenum-graecum*) seed. Carbohydr. Res. 104: 348-353. (CA. 97: 69258k).
- Chauhan, J. S. and Kumari, G. 1979. A new glycoflavonol from the root bark of *Artocarpus lakoocha*. Planta Med. 37: 86-88.

- Chauhan, J. S., Kumari, G., Kumar, S. and Chaturvedi, R. 1982. Chemical examination of the root bark of *Artocarpus lakoocha*. Proc. Natl. Acad. Sci, India, Sect. A 52: 217-218. (CA. 99: 102301k).
- Chauhan, J. S., Kumari, G. and Saraswat, M. 1979. A new flavonol glycoside from the root bark of *Artocarpus lakoocha*. Indian J. Chem. 18B: 473-475.
- Chen, C. C., Huang, Y. L., Ou, J. C., Lin, C. F. and Pan, T. M. 1993. Three new prenylflavones from *Artocarpus altilis*. J. Nat. Prod. 56: 1594-1597.
- Chung, M. I., Lu, C. M., Huang, P. L. and Lin, C. N. 1995. Prenylflavonoids of *Artocarpus heterophyllus*. Phytochemistry 40: 1279-1281.
- Daulatabad, C. D. and Mirajkar, A. M. 1989. Ricinoleic acid in *Artocarpus integrifolia* seed oil. J. Am. Oil. Chem. Soc. 66: 1631.
- Dayal, R. and Seshadri, T. R. 1974. Courless components of the roots of *Artocarpus heterophyllus*: Isolation of a new compound, artoflavanone. Indian J. Chem. 12: 895-896.
- Durand, E., Ellington, E. V., Feng, P. C., Haynes, L. J., Magnus, K. E. and Philip, N. 1962. Simple hypotensive and hypertensive principle from some West Indian medicinal plants. J. Pharm. Pharmacol. 14: 562-566.
- Farnsworth, N. R. and Bunyapraphatsara, N. 1992. Thai Medicinal Plants Recommended for Primary Health Care System. pp. 67-69. Bangkok: Faculty of Pharmacy, Mahidol University.
- Ferrari, F., Messana, I. and do Mesquita de Araujo, M. 1989. Structure of three new flavone derivatives, brosimones G, H and I, from *Brosimopsis oblongifolia*. Planta Med. 55: 70-72.
- Ferreira De Miranda-Santos, I. K., Delgado, M., Bonini, P. V., Bunn, M. M. M. and Campos, N. A. 1992. A crude extact of *Artocarpus integrifolia* contains two lectins with distinc biological activities. Immunol. Lett. 31: 65-71. (CA. 116: 78788w).
- Fujimoto, Y., Agusutein, S. and Made, S. 1987. Isolation of a chalcone derivative and antitumor compositions containing it. Jpn. Kokai Tokkyo Koho JP 62270544 (CA. 110: 13561y).
- Fujimoto, Y., Uzava, J., Suhanda, S., Soemartono, A., Sumatra, M. and Koshihara, Y. 1987. Isolation and structural elucidation of new lipoxygenase inhibitors from Indonesian *Artocarpus communis*. Tennen Yuki Kagobutsu Toronkai Koen Yoshishu 29: 721-728. (CA. 108: 164664u).

- Fujimoto, Y., Zhang, X. X., Kirisawa, M., Uzawa, J. and Sumatra, M. 1990. New flavones from *Artocarpus communis* Forst. Chem. Pharm. Bull. 38: 1787-1789.
- Fukai, T. and Nomura, T. 1993. ¹H NMR spectra of prenylated flavonoids and pyranoflavonoids. Heterocycles 36: 329-343.
- Gelder, C. W. G., Flurkey, W. H. and Wicher, H. J. 1997. Sequence and structural features of plant and fungal tyrosinases. Phytochemistry 45: 1309-1323.
- Gerber, N. N. 1986. Phenolics from osage wood cleavage of oxyresveratrol. Phytochemistry 25: 1697-1699.
- Hagiwara, K., Collet, C. D., Kobayashi, K. and Vaerman, J. P. 1988. Jacalin: Isolation, characterization and influence of various factors on its interaction with human IgAI, as assessed by precipitation and latex agglutination. Molec. Immunol. 25: 69-83.
- Hano, Y., Aida, M. and Nomura, T. 1990. Two new natural Diels-Alder-type adducts from the root bark of *Artocarpus heterophyllus*. J. Nat. Prod. 53: 391-395.
- Hano, Y., Aida, M., Shiina, M., Nomura, T., Kawai, T., Ohe, H. and Kagei, K. 1989. Artonins A and B, two new prenylflavones from the root bark of *Artocarpus heterophyllus* Lamk. Heterocycles 29: 1447-1453.
- Hano, Y., Inami, R. and Nomura, T. 1990. Components of the bark of *Artocarpus rigida* Bl. 1. Structures of two new isoprenylated flavones, Artonins G and H. Heterocycles 31:2137-2179.
- Hano, Y., Inami, R. and Nomura, T. 1993. Components of the bark of *Artocarpus rigida* Bl. 2: Structures of four new isoprenylated flavone derivatives artonins M, N, O, and P. Heterocycles 35: 1341-1350.
- Hano, Y., Inami, R. and Nomura, T. 1994. Constituents of Moraceae plants. 20. A novel flavone, artonins V, from the root bark of *Artocarpus altilis*. J. Chem. Res. Synop. 348-349. (CA. 251263z).
- Hano, Y., Matsumoto, Y., Shinohara, K., Sun, J. Y. and Nomura, T. 1990. Cudraflavones C and D, two new prenylflavones from the root bark of *Cudrania tricuspidata* (Carr.) Bur. Heterocycles 31: 1339-1344.
- Hano, Y., Yamagami, Y., Kobayashi, M., Isohata, R. and Nomura, T. 1990. Artonins E and F, two new prenylflavones from the bark of *Artocarpus communis* Forst. Heterocycles 31:877-882.

- Hashim, O. H., Gendeh, G. S. and Jaafar, M. I. N. 1992. Lectin extracts of Champedek seeds. demonstrate selective stimulation of T lymphocyte proliferation. Biochem. Int. 27:139-143.
- Heupel, R. C., Sauvaire, Y., Le, P. H., Parish, E. J. and Nes, W. D. 1986. Sterol composition and biosynthesis in sorghum: importance to developmental regulation. Lipids 21: 69-75.
- Iida, K., Hase, K., Shimomura, K., Sudo, S., Kadota, S. and Namba, T. 1995. Potent inhibitors of tyrosinase activity and melanin biosynthesis from *Rhem officinale*. Planta Med. 61:425-428.
- Iribarren, A. M. and Pomilio, A. B. 1985. Sitosterol 3-O- α -D-riburonofuranoside from *Bauhinia candicans*. Phytochemistry 24: 360-361.
- Kapil, R. S. and Joshi, S. S. 1960. Chemical constituents of *Artocarpus lakoocha*. J. Sci. Ind. Research (India) 19B: 498. (CA. 55:15536a).
- Khalil, M. W. and Idler, D. R. 1980. Sterols of Scollop. III. Characterization of some C-24 epimeric sterols by high resolution (220 MHz) nuclear magnetic resonance spectroscopy. Lipids 15: 69-73.
- Kielland, I. S. and Malterud, K. E. 1994. Triterpenoids from *Artocarpus intregrifolia* fruits. Planta Med. 60:196.
- Kijjoa, A., Cidade, H. M., Gonzalez, M. T. G., Afonzo, C. M., Silva, A. M. S. and Herz, W. 1998. Further prenylflavonoids from *Artocarpus elasticus*. Phytochemistry 47: 875-878.
- Kijjoa, A., Cidade, H. M., Pinto, M. M. M., Gonzalez, M. J. T. G., Anantachoke, C., Gedris, T. E. and Herz, W. 1996. Prenylflavonoids from *Artocarpus elasticus*. Phytochemistry 43: 691-694.
- Kingrungpet, B. 1994. Phytochemical study of Artocarpus gomezianus Wall. ex Trec. Leaves. (Master's thesis, Department of Pharmacognosy, Graduate School, Chulalongkorn University).
- Kirtikar, K. R. and Basu, B. D. 1980. Indian medicinal plants vol. 3. India: Lalit Mohan Basu. pp. 119-133.
- Kochummen, K. M. 1978. Tree flora of Malaya vol. 3. Malaysia: Forest Department, Ministry of Primary Industries. pp. 119-133.
- Koshihara, Y., Fujimoto, Y. and Inoue, H. 1988. A new 5-lipoxygenase selective inhibitor derived from *Artocarpus communis* strongly inhibits arachidonic acid-induced ear edema. Biochem. Pharmacol. 37: 2161-2165.

- Kubo, I. 1997. Tyrosinase inhibitor from plants. Phytochemical for Pest Control. pp. 311-326. Washington: American Chemical Society.
- Kubo, I., Yokokawa, Y. and Kinst-Hori, I. 1995. Tyrosinase inhibitors from bolivian medicinal plants. J. Nat. Prod. 58: 739-734.
- Kumar, N. S., pavanarasivam, G., Sultanbawa, M. U. S. and Mageswaran, R. 1977. Chemical investigation of Ceylonese plants. Part 24. New chromenoflavonoids from the bark of *Artocarpus nobilis* Thw. (Moraceae). J. C. S. Perkin I: 1243-1251.
- Lin, C. N. and Lu, C. M. 1993. Heterophylol, a phenolic compound with novel skeleton from *Artocarpus heterophyllus*. Tetrahedron Lett. 34: 8249-8250.
- Lin, C. N., Lu, C. M. and Huang, P. L. 1995. Flavonoids from *Artocarpus heterophyllus*. Phytochemistry 39: 1447-1451.
- Lin, C. N. and Shieh, W. L. 1991. Prenylflavonoids and a pyranodihydrobenzoxanthrone from *Artocarpus communis*. Phytochemistry 30: 1669-1671.
- Lin, C. N. and Shieh, W. L. 1992. Pyranoflavonoids from *Artocarpus communis*. Phytochemistry 31:2922-2924.
- Lin, C. N., Shieh, W. L. and Jong, T. T. 1992. A pyranodihydrobenzoxanthone epoxide from *Artocarpus communis*. Phytochemistry 31: 2563-2564.
- Lu, C. M. and Lin, C. N. 1993. Two 2',4',6'-trioxygenated flavones from *Artocarpus heterophyllus*. Phytochemistry 33: 909-911.
- Lu, C. M. and Lin, C. N. 1994. Flavonoids and 9-hydroxytridecyl docosanoate from *Artocarpus heterophyllus*. Phytochemistry 35: 781-783.
- Mahato, S. B., Banerjee, S. K. and Chakravarti, R. N. 1971. Triterpenoids of the stem bark of *Artocarpus chaplasha*. Phytochemistry 10: 1351-1354.
- Markham, K. R. 1982. Techniques of flavonoids identification. London: Academic Press.
- Masamoto, Y., Iida, S. and Kubo, M. 1980. Inhibitory effect of chinese crude drugs on tyrosinase. Planta Med. 40: 361-365.
- Mongolsuk, S., Robertson, A. and Towers, R. 1957. 2,4,3',5'-Tetrahydroxystilbene from *Artocarpus lakoocha*. J. Chem. Soc.: 2231-2233.
- Morita, H., Kayashita, T., Kobata, H., Gonda, A., Takeya, K. and Itokawa, H. 1994. Pseudostellalin A-C, new tyrosinase inhibitory cyclic peptides from *Pseudostellaria heterophylla*. Tetrahedron 50: 6797-6804.

- Mu, Q. Z. and Li, Q. X. 1982. The isolation and identification of morin-calcium chelated compound from *Artocarpus pithecogallus* C. Y. Wu and *Artocarpus heterophyllus* Lamk. Chih Wu Hsueh Pao 24: 147-153. (CA. 97: 107043w).
- Nakajima, K., Taguchi, H., Endo, T. and Yosioka, I. 1978. The constituents of *Scirpus fluviatilis* (Torr.) A Gray. I. The structures of two new hydroxystilbene dimers, Scirpusin A and B. Chem. Pharm. Bull. 26: 3050-3057.
- Namjuntra, P. and Chulavatnatol, M. 1984. Purification and characterization of a lectin from seeds of jackfruit (*Artocarpus heterophyllus*). Abst. 10 th Conference of Science and Technology, Thailand. Changmai Univer., Changmai, Thailand. pp. 356-357.
- Nath, M. C. and Mukherjee. 1939. Crystallographic investigation of artostenone, the stenone isolated from the Indian summer fruit, *Artocarpus integrifolia* by means of Goniometer and X-rays. J. Indian Chem. Soc. 16: 229-234.
- Nogueira, P. L. and Correia, A. A. 1958. Chemical study of a tropical wood. Garcia orta 6: 679-688. (CA. 53: 20792a).
- Nomura, T., Hano, Y. and Aida, M. 1998. Isoprenoid-substituted flavonoids from *Artocarpus* plants (Moraceae). Heterocycles 47: 1179-1205.
- Pant, R. and Chaturvedi, K. 1989. 4-Hydroxyundecyl docosanoate and cycloartenone in *Artocarpus integrifolia* latex. Phytochemistry 28: 2197-2199.
- Paolo, P., Alessandro, P., Giorgio, H., Holisotan, H. E., Lukman, M., Aritin, A. S. and Babara, G. 1998. A new prenylated flavone from *Artocarpus champeden* inhibits the 'K⁺-dependent amino acid transport in *Bombyx mori* midgut. Biochem. Biophys. Res. Commun. 244: 445-448. (CA. 128: 306539w).
- Parthasarathy, P. C., Radhakrishnan, P. V., Rathi, S. S. and Venkataraman, K. 1969. Coloring matters of the wood of *Artocarpus heterophyllus*. V. Cycloartocarpesin and oxydihydroartocarpesin, two new flavones. Indian J. Chem. 7: 101-102. (CA. 71: 70451n).
- Pavanarasivum, G. and Sultanbawa, M. U. S. 1973. Cycloartenyl acetate, cycloartenol and cycloartenone in the bark of *Artocarpus* species. Phytochemistry 12: 2725-2726.
- Pavanarasivum, G., Sultanbawa, M. U. S. and Mageswaran, R. 1974. Chemical investigation of Ceylonese plants. X. New chromenoflavonoids from the bark of *Artocarpus nobilis* (Moraceae). Chem. Ind. (London): 875-876. (CA. 82: 121639s).

- Pavaro, J. and Reutrakul, V. 1976. New flavones in *Artocarpus lakoocha*. Mahidol Univ. J. Pharm. Sci. 3: 161-164.
- Pendse, A. D., Pendse, R., Rao, A. V. R. and Venkataraman, K. 1976. Integrin, cyclointegrin and oxyisocyclointegrin, three new flavones from the heartwood of *Artocarpus integer*. Indian J. Chem. 14B: 69-72.
- Pereira, J. R., Medina, H. and Bustos, R. E. 1962. On the presence of acetylcholine in seeds. of *Artocarpus integrifolia* (Moraceae) and of *Annona squamosa* (Annonaceae). An. Fac. Med. Univ. Parana 5: 45-47. (Biological Abstracts 46: 12891).
- Perry, L. M. 1980. Medicinal plants of East and Southeast Asia. Cambridge: The MIT Press.
- Poopyruchpong, N., Rungruangsak, K., Nimmanpisut, S., Panijpan, B. and Ratanabanangkoon, K. 1978. Some physico-chemical properties of 2,4,3',5'-Tetrahydroxystilbene. J. Sci. Soc. Thailand 4: 163-167.
- Radhakrishnan, P. V., Rao, A. V. R. and Venkataraman, K. 1965. Two new flavones from *Artocarpus heterophyllus*. Tetrahedron Lett. 663-667.
- Rao, A. V. R., Rathi, S. S and Venkataraman, K. 1972. Chaplashin, a flavone containing an oxepine ring from the heartwood of *Artocarpus chaplasha* Roxb. Indian J. Chem. 10:905-907.
- Rao, A. V. R., Varadan, M. and Venkataraman, K. 1971. Coloring matters of the wood of *Artocarpus heterophyllus*: part VI. Cycloheterophyllin, a flavone linked to three isoprenoid groups. Indian J. Chem. 9: 7-13.
- Rao, A. V. R., Varadan, M. and Venkataraman, K. 1973. Coloring matters of the wood of *Artocarpus heterophyllus*: part VII. Isocycloheterophyllin, a new flavone. Indian J. Chem. 11: 298-299. (CA. 79: 31957c).
- Reddy, G. R., et al. 1991. A prenylflavone, artonin E, as arachidonate 5-lipoxygenase inhibitor. Biochem. Pharmacol. 41: 115-118.
- Shieh, W. L. and Lin, C. N. 1992. A quinonoid pyranobenzoxanthone and pyranodihydrobenzoxanthone from *Artocarpus communis*. Phytochemistry 31: 364-367.
- Shimizu, K., Kondo, R., Sakai, K., Lee, S. H. and Sato, H. 1998. The Inhibitory Components from *Artocarpus incisus* on Melanin Biosynthesis. Planta Med. 64: 408-412.

- Smitinand, T. 1980. Thai plant names (Botanical names-vernacular names). pp. 34-35. Bangkok: Funny Publishing.
- Sultanbawa, M. U. S. and Surendrakumar, S. 1989. Two pyranodihydrobenzo-xanthones from *Artocarpus nobilis*. Phytochemistry 28: 599-605.
- Sultankhodzhaew, M. N. and Tadzhibaev, M. M. 1976. Phenyl- β -naphthylamine from three species of plants. Khim. Prir. Soedin 3: 406-407. (CA. 85: 139721j).
- Suresh, G. K., Appukuttan, P. S. and Basu, D. K. 1982. Alpha-D-galactose specific lectin from jackfruit (*Artocarpus integrifolia*) seed. J. Biol. Sci. 4: 257-261. (Medicinal Aromatic Plant Abstracts 83: 83010447).
- Takasuki, M., Munoz, L. and Mazamune, T. 1978. Stilbene phytoalexins from diseased mulberry. Chemistry Lett. 1241-1242.
- Ultee, A. J. 1949. Components of samples of latex (from a number of tropical trees). Pharm. Weekblad 84: 65-70. (CA. 43: 4041i).
- Venkataraman, K. 1972. Wood phenolics in the chemotaxonomy of the Moraceae. Phytochemistry 11: 1571-1586.
- Wright, J. L. C., et al. 1978. Identification of C-24 alkyl epimers of marine sterols by ^{13}C nuclear magnetic resonance spectroscopy. Can. J. Chem. 56: 1898-1903.
- Yamazaki, M., Okuyama, E., Matsudo, T., Takamaru, T. and Kaneko, T. 1987. Principle of Indonesian herbal drugs having antiulcerogenic activity I., isolation and identification of (\pm)-catechin from *Artocarpus integrifolia* Merr. Yakugaku Zasshi 107: 914-916. (CA. 108: 68693g).

VITA

Mr. Boonchoo Sritularak was born on January 5, 1973 in Phuket, Thailand. He received his Bachelor's degree of Science in Pharmacy in 1996 from the Faculty of Pharmaceutical Sciences, Prince of Songkla University, Thailand.

Publications

1. Sritularak, B., De-Eknamkul, W. and Likhitwitayawuid, K. 1998. Tyrosinase Inhibitors from *Artocarpus lakoocha* Roxb. pp. 462-463. 24th Congress on Science and Technology of Thailand, 19-21 October 1998, Queen Sirikit National Convention Center, Bangkok.
2. Sritularak, B., De-Eknamkul, W. and Likhitwitayawuid, K. 1998. Tyrosinase Inhibitors from *Artocarpus lakoocha*. Thai J. Pharm. Sci. 22: 149-155.

