

REFERENCES

1. Polunin, M.; and Robbins, C. The Natural Pharmacy. 4nd ed. London: Dorling Kindersley Press, 1992.
2. Silveira, E. R.; and McChesney, D. J. 6,7-Oxygenate neo-clerodane furan diterpenes from *croton sonderianus*. Phytochemistry 36 (1994): 1457-1463.
3. Krebs, H. C.; and Ramiarantsoa, H. Clerodane diterpenes and other constituents of *croton hovarum*. Phytochemistry 41 (1996): 561-563.
4. Sato, A.; Kurabayashi, M.; Nagahori, H.; Ogiso, A.; and Mishima, H. Chettaphanin I, A novel furanoditerpenoid. Tetrahedron Lett. (1970): 1095-1098.
5. Sato, A.; Kurabayashi, M.; Nagahori, H.; and Ogiso, A. Chettaphanin II, A novel furanoditerpenoids. Tetrahedron Lett. (1971): 839-842.
6. Pongboonrod, S. Mai Tet Muang Thai. Bangkok: Kasembunnakich Press, 1976.
7. Faculty of Pharmaceutical Science, Mahidol University. Saim Paisutchayaprug. Bangkok: Mahidol University, 1995.
8. Taingbooranatam, W. Dictionary of Animals and Plants in Thailand. 7th ed. Bangkok: Bamroongsarn Press, 1993.
9. Shaw, H. K. The Euphorbiaceae of saim. Kew Bulletin. 26(1972): 191-233.
10. Whitmore, T. C.; and Longman, F. L. S. Forest Department Ministry of Prymary Industry Malasia. Tree Frora of malaya, 2 (1973):78.
11. Ayres, D. C.; ed al. Dictionary of Natural Product. London: Chapman & Hall chemical data base (1994) 7.
12. Boonyarathanakornkit, L.; Chun-tao, C.; Fong, H.S.; and Farnsworth. N. R. Constituents of *croton crassifolius* Roots. Planta Med. 54(1988): 61-63.
13. Siems, K.; Jakupovic, J.; Castro, V.; and Poveda, L. Constituents of two Acalypha species. Phytochemistry 41(1996): 851-858.

14. Aiyar, V., N.; and Seshadri, T. R. 11-Dehydro(-)hardwickiic acid from *Croton oblongifolius* Roxb. Phytochemistry 11(1972): 1473.
15. Wen-Chiung, S.; Jim-Min, F.; and Yu-Shia, C. Labdanes from *cryptomeria japonica*. Phytochemistry 37(1994): 1109.
16. Bohlmann, F.; Dutta, L.; Robinson, H.; and King, R. M. A series of furanoid 7 β -acetoxylabdanes from *Austro eupatorium* chaparense. Phytochemistry 18(1979): 1839.
17. Rolando, A. S.; and Alejandro, J. V. 7- α -Acetoxyhardwiic acid: A furanoid clerodane Phytochemistry 35 (1994): 537.
18. Rahman, A.; ed al. New furanoid diterpenoidal constituteds of *Tinospora malabarica*. Tetrahedron 50(1994): 12109.
19. Isao, K.; Partomuan, S.; Tomokazu, W.; Hirotaka, S.; Satoshi, F.; and Yuriko, Y.; Motomasa, K., Indonesian medicinal plants. XI chemical structures of Caesaldekariins a and b, two new cassane-type furanoditerpenes. Chem. Pharm. Bull. 42(1994): 1798.
20. Anderson, M. M.; Neill, M. J.; Phillipson, J. D.; and Warhurst, D. C. In vitro Cytotoxicity of a series of Quassinoids from *Brucea Javanica* Fruits Against KB cell. Planta Med., 57(1991): 62-64.
21. Solis, P. N.; Wright, C. W.; Anderson, M. M.; Gupta, M. P.; and Phillipson, J. D., A microwell cytotoxicity Assay using *Atemia salina* (Brine shrimp). Planta Med., 59(1993): 250-252.
22. Fressenden, R. J.; and Fressenden, J. S. Techniques and Experiments for Organic Chemistry. London: Willard Grant Press, (1983): 227.
23. Randerath, K. Thin-Layer Chromatography. 2 nd rev. U.S.A.: Academic Press, (1996): 441.
24. Raksilp, T. Chemical Constituents of the Tuberous Roots of *Butea superba* Roxb. Master's Thesis, Department of Chemistry, Graduate school, Chulalongkorn University, (1995).

25. Calegate, S.M.; and Molyneux, R.J. Bioactive Natural Products, London: CRC Press, (1993): 441.
26. The Official Methods of Analysis of the Association of Official Analytical Chemistry, 11th ed, Official First Action, (1970): 843.
27. Hostellmann, K.; Marston, A.; and Maillard, M. The role of TLC in the investigations of medicinal plants of Africa, South America and other topical regions. GIT laboratory Journal (1997): 36-39.
28. Hostellmann, K.; Terreaux, C.; Marston, A.; and Potterate, O. The role of Planar Chromatography in the Rapid Screening and Isolation of Bioactive compounds from medicinal plants. J. Planar Chromatography 10(1997): 251-257.
29. Hostellmann, K.; Potelrat, O.; and Wolfender, J. Strategy in the search for new bioactive plant constituents. Pharm. Ind. 59(1997): 339.
30. Pinjinda, S. Chemical constituents and biological activities of the fruits of *Xylocarpus granatum* Koen. Master's Thesis, Department of Chemistry, Graduate school, Chulalongkorn University, (1986).
31. Weckert, E.; Hummer, K.; Addae-Mensah, I.; and Achenbach, H. The absolute configuration of Chirimonide. Phytochemistry 31(1992): 2170.
32. Bruno, M.; Fazio, C.; Piozzi, F.; Rodriguez, B.; and DELA Torre, M. C. Neo-clerodane diterpenoids from *Teucrium corymbosum*. Phytochemistry 40(1995): 1481-1483.
33. Englert, G.; Akemann, T.; Schiedt, K.; and Liaaen, S. Structure elucidation of the algal carotenoid (3S,5R,6R,3'S,5'R,6'S-cis-7',8'-Dihydroneoxanthin-20'-Al3'-(-)-Lactoside(P457). Part 2 NMR studies. J. Nat. Prod. 58(1995): 1675.
34. Martin, T. S.; Ohtani, K.; Kasai, R.; and Yamasaki, K. Furano diterpene Glycoside from *Tinospora Rumphii*. Phytochemistry 42(1966): 153-158.
35. Maldonado, E.; Cardenas, J.; Bojorquez, H.; and Escamilla, E.M.; Ortega, A. Amarisolide, a neo-clerodane diterpene glycoside from *salvia amarissima*. Phytochemistry 42 (1996): 1105-1108.

36. Hanson, J.R.; Nasir, H.; and Parvez, A. The Hydroxylation of testosterone and some relatives by *Cephalosporium aphidicola*. Phytochemistry 42(1996): 411-415.
37. Mopondo, E. M.; and Chulia, A. J. 6'-O-(-D-Glycosyl Gentiopicroside: A new secoiridoid from *Gentiana asclepiadea*. Planta Med. (1988): 185.
38. Takano, S.; and Kawaminami, S. Cyperanyl acetate and cyperanal from *Cirsium dipsacolepis*. Phytochemistry. 27(1988): 1197-1199.
39. Jacobs, H.; Lachmaning, S. S.; Ramdayal, F.; McLean, S.; Puzzuoli, F.; and Reynolds, W.F. Applications of 2D-NMR spectroscopy to phytochemical studies; cyperenol and cyperenopic acid. J. Nat. Prod. 50(1987): 835-842.
40. Boonyaratavej, S.; and Roengsumran, S. X-Rays structure of cyperenoic acid from *Croton crassifolius*. J. Nat Prod, 51(1988): 769-770.
41. Inagaki, F.; and Abe, A. Analysis of ¹H- and ¹³C-Nuclear Magnetic Resonance Spectra of Spathulenol by Two-dimensional Methods. J. Chem. Soc. Perkin Trans II (1985): 1173-1178.
42. Tringali, C.; Piattelli, M.; and Spatafora, C. Sesquiterpenes and geranylgeranylglycerol from the brown algae *Taonia lacheana* and *Taonia atomaria* F. ciliata: their chemotaxonomic significance. Phytochemistry 40(1995): 827-831.
43. Beechan, C. M.; and Djerassi, C. Terpenoids-LXXIV: The sesquiterpenes from the soft coral *simularia mayi*. Tetrahedron 34(1978): 2503-2508.
44. Bohlmann, F.; Grenz, M.; Jakupovic, J.; King, R. M.; and Robinson, H. Four heliangolides and other sesquiterpenes from *Brasilia sickii*. Phytochemistry 22(1983): 1213-1218.
45. Rashid, M. A.; Gray, A. I.; and Watermann P. G. Aromadendrane sesquiterpene from *Phebalium*, *Drummondita* and *Eriostemon* species. J. Nat. Prod. 58(1995): 618.
46. Toyota, M.; Koyama, H.; Mizutani, M.; and Asakawa, Y. (-)-Ent-spathulenol isolated from liverworts is an artefact. Phytochemistry 41(1996): 1347-1350.

47. Gijssen, H. J. M.; Wijnberg, J. B. P.A.; Stork, G. A.; and Groot, A. The synthesis of Mono- and Dihydroxy Aromadendrane Sesquiterpenes, starting from Natural (+)-Aromadendrane-III. Tetrahedron 48(1992): 2465.
48. Cardóna, L.; Garcia, B.; Pedro, J. R.; and Perez, J. 6-Prenyloxy-7-methoxy coumarin, a coumarin-hemiterpene ether from *Carduus tenuiflorus*. Phytochemistry 31(1992): 3989-3991.
49. Gottlieb, H. E.; Lima, R. A.; and Monache, F. D., ¹³C Nuclear Magnetic Resonance Spectroscopy of 6-and 7-Substituted Coumarins; Correlation with Hammett Constants. J. Chem. Soc. Perkin Trans II (1979): 436-437.
50. Silvan, A. M.; Abad, M. J.; Bermejo, P.; Sollhuber, M.; Villar, A. Antiinflammatory activity of coumarins from *Santolina oblongifolia*. J. Nat.Prod. 59(1996): 1183.
51. Corbett, R. E.; Cumming, S. D., Lichens and Fungi. Part XI 14(-Taraxerane). J. Chem. Soc. Perkin Trans I (1972): 2827.
52. Chavasiri, W. Chemical constituents and biological activities of *Rhizophora apiculata* Bl. Master's Thesis, Department of Chemistry, Graduate school, Chulalongkorn University, (1988).
53. Yang, L.; YaPing, S.; Zhong-Jian, J.; Sadiq, S.; and Jamil, L. Four esters of a new pentacyclic diterpenoid of the myrsinol type from *Euphorbia aleppica*. J. Nat. Prod. 58(1995): 1883-1888.
54. Rustaiyan, A.; Mosslemin-kupaii, M. H.; Papastergiou, F.; and Jakupovic, J. Persianone, a diameric diterpene from *Ballota aucheri*. Phytochemistry 40(1995): 875-879.

VITA

Miss Butsarakham Supudompol was born on July 13, 1973 in Ubonrachathanee province, Thailand. She received a Bachelor Degree of Science in Chemistry at Ramkhamhaeng University in 1994. In 1996, she has been a graduate student studying Organic Chemistry at Chulalongkorn University. During her study towards the Master Degree, she was awarded a teacher assistantship by the Faculty of Science of Chulalongkorn University and was also supported a research grant for her Master degrees's thesis by the Graduate School of Chulalongkorn University.