

REFERENCES

- [1] Haxon, E., Melhus, A., Hylander, L.D. Silver – a toxic threat to our health and environment. Uppsala University. in Dept. of Clinical Microbiology. 2007.
- [2] Sangsuk, S. Center of excellence on hazardous substance management-HSM. 2007. Available from: <http://www.chemtrack.org/News-Detail.asp?TID=5&ID=2> [20 Dec 2013]
- [3] Senjen, R. Nanotechnology Project - Friends of the Earth Australia. 2007. Available from: <http://nano.foe.org.au/node/190> [10 Jan 2014]
- [4] Benn, T.M. and Westerhoff, P. cNanoparticle silver released into water from commercially available sock fabrics. Environmental Science & Technology 42(11) (2008): 4133-4139.
- [5] Farkas, J., et al. Characterization of the effluent from a nanosilver producing washing machine. Environment International 37(6) (2011): 1057-1062.
- [6] Echegoyen, Y. and Nerin, C. Nanoparticle release from nano-silver antimicrobial food containers. Food and Chemical Toxicology 62 (2013): 16-22.
- [7] Kulthong K1, S.S., Boonpavanitchakul K, Kangwansupamonkon W, Maniratanachote R. Determination of silver nanoparticle release from antibacterial fabrics into artificial sweat. particle and fibre toxicology 7 (2010): 8.
- [8] Radulescu, M.C., Chira, A., Radulescu, M., Bucur, B., Bucur, M.P., and Radu, G.L. Determination of silver(I) by differential pulse voltammetry using a glassy carbon electrode modified with synthesized N-(2-aminoethyl)-4,4'-bipyridine. Sensors (Basel, Switzerland) 10(12) (2010): 11340-11351.
- [9] Koh, T. and Sugimoto, T. Extractive spectrophotometric determination of silver(I) at the 10^{-7} M level using 1,10-phenanthroline and tetrabromophenolphthalein ethyl ester. Analytica Chimica Acta 333(1-2) (1996): 167-173.
- [10] Manzoori, J.L., Abdolmohammad-Zadeh, H., and Amjadi, M. Ultra-trace determination of silver in water samples by electrothermal atomic absorption spectrometry after preconcentration with a ligand-less cloud point extraction methodology. Journal of Hazardous Materials 144(1-2) (2007): 458-463.
- [11] Rahman, M.A., Kaneko, S., Amin, M.N., Suzuki, T., and Ohta, K. Determination of silver in environmental samples by tungsten wire preconcentration



- method-electrothermal atomic absorption spectrometry. *Talanta* 62(5) (2004): 1047-1050.
- [12] Argekar, A.A., Kulkarni, M.J., Mathur, J.N., Page, A.G., and Iyer, R.H. ICP-AES determination of silver after chemical separation from uranium matrix. *Talanta* 42(12) (1995): 1937-1942.
- [13] Luo, Y., Dabek-Zlotorzynska, E., Celo, V., Muir, D.C.G., and Yang, L. Accurate and precise determination of silver isotope fractionation in environmental samples by multicollector-ICPMS. *Analytical Chemistry* 82(9) (2010): 3922-3928.
- [14] Al-Andis, N.M. Determination of lead, cadmium and silver in industrial wastewater of Riyadh, Saudi Arabia by SIA/ICP/MS technique. *Journal-King Saud University Science* 10(1) (1998): 85-92.
- [15] Bilberg, K., Hovgaard, M.B., Besenbacher, F., and Baatrup, E. In vivo toxicity of silver nanoparticles and silver ions in Zebrafish (*Danio rerio*). *Journal of Toxicology* 2012 (2012).
- [16] Ngeontae, W., Janrungroatsakul, W., Morakot, N., Aeungmaitrepirom, W., and Tuntulani, T. New silver selective electrode fabricated from benzothiazole calix[4]arene: Speciation analysis of silver nanoparticles. *Sensors and Actuators B: Chemical* 134(2) (2008): 377-385.
- [17] Chen, L., Ju, H., Zeng, X., He, X., and Zhang, Z. Silver ion-selective electrodes based on novel containing benzothiazolyl calix[4]arene. *Analytica Chimica Acta* 437(2) (2001): 191-197.
- [18] Mahajan, R.K. and Parkash, O. Silver(I) ion selective PVC membrane based on bis-pyridine tetramide macrocycle. *Talanta* 52(4) (2000): 691-693.
- [19] Xu, D. and Katsu, T. O,O,O-trialkyl phosphorothioates as simple and effective ionophores for silver ion-selective membrane electrodes. *Analytica Chimica Acta* 443(2) (2001): 235-240.
- [20] Zhao, Q., Chai, Y., Yuan, R., Zhang, T., and Yang, C. A new silver(I)-selective electrode based on derivatized MWCNTs@SiO₂ nanocomposites as a neutral carrier. *Materials Science and Engineering: C* 32(6) (2012): 1352-1357.
- [21] Bualom, C., et al. Bulk optode sensors for batch and flow-through determinations of lead ion in water samples. *Talanta* 82(2) (2010): 660-667.
- [22] A., B. *Physical Methods in Chemistry and Nano Science* Connexions Website Available from: <http://cnx.org/content/col10699/1.14/>. [22 Jan 2014]
- [23] Dybko, A. Errors in chemical sensor measurements. *Sensors* 1(1) (2001): 29-37.
- [24] Bradl, H.B. Heavy metal in the environment. *Elsevier* 6 (2004).



- [25] Järup, L. Hazards of heavy metal contamination. *British Medical Bulletin*. 68(1) (2003): 167-182.
- [26] Pollution Control Department, Bangkok, Thailand. 2013. Available from: http://www.pcd.go.th/info_serv/reg_std_water01.html [12 Nov 2013]
- [27] Industrial Estate Authority of Thailand. 2011. Available from: <http://www.ieat.go.th/ieat/index.php/th/> [5 Dec 2013]
- [28] Mayr, T. Optical sensors for the determination of heavy metal ions. (2002).
- [29] Deng, G. Principles of chemical and biological sensors. *Materials and Manufacturing Processes* 14(4) (1999): 623-625.
- [30] Anna Grazia, M. and Francesco, B. Fibre-optic sensors in health care. *Physics in Medicine and Biology* 42(5) (1997): 967-979.
- [31] Chan, D.W. 21 - Automation of immunoassays. in Diamandis, E.P. and Christopoulos, T.K. (eds.),Immunoassay, pp. 483-504. San Diego: Academic Press, 1996.
- [32] Wolfbeis, O.S., et al. Set of luminescence decay time based chemical sensors for clinical applications. *Sensors and Actuators B: Chemical* 51(1-3) (1998): 17-24.
- [33] Huber, C., Klimant, I., Krause, C., and Wolfbeis, O.S. Dual Lifetime Referencing as Applied to a Chloride Optical Sensor. *Analytical Chemistry* 73(9) (2001): 2097-2103.
- [34] Krause, C., Werner, T., Huber, C., and Wolfbeis, O.S. Emulsion-based fluorosensors for potassium featuring improved stability and signal Change. *Analytical Chemistry* 71(23) (1999): 5304-5308.
- [35] Huber, C., Klimant, I., Krause, C., Werner, T., Mayr, T., and Wolfbeis, O.S. Optical sensor for seawater salinity. *Fresenius' Journal of Analytical Chemistry* 368(2-3) (2000): 196-202.
- [36] Lerchi, M., Reitter, E., Simon, W., Pretsch, E., Chowdhury, D.A., and Kamata, S. Bulk optodes based on neutral dithiocarbamate ionophores with H]high selectivity and sensitivity for silver and mercury cations. *Analytical Chemistry* 66(10) (1994): 1713-1717.
- [37] Silk, M.G. Review of progress in quantitative nondestructive evaluation: Edited by D.O. Thompson and D.E. Chimenti Plenum Press, New York, USA, 1988 NDT International 21(6) (1988): 469-470.
- [38] Bakker, E., Bühlmann, P., and Pretsch, E. Carrier-based Ion-selective electrodes and bulk Optodes. 1. General characteristics. *Chemical Reviews* 97(8) (1997): 3083-3132.



36228910427

- [39] Shamsipur, M., Sadeghi, M., Alizadeh, K., Sharghi, H., and Khalifeh, R. An efficient and selective fluorescent optode membrane based on 7-[(5-chloro-8-hydroxy-7-quinoliny) methyl]-5,6,7,8,9,10-hexahydro-2H-1,13,4,7,10-benzodioxatriazacyclopentadecine-3,11(4H,12H)-dione as a novel fluoroionophore for determination of cobalt(II) ions. *Analytica Chimica Acta* 630(1) (2008): 57-66.
- [40] Shamsipur, M., Poursaberi, T., Avanes, A., and Sharghi, H. Copper(II)-selective fluorimetric bulk optode membrane based on a 1-hydroxy-9,10-anthraquinone derivative having two propenyl arms as a neutral fluorogenic ionophore. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* 63(1) (2006): 43-48.
- [41] He, C.-L., Ren, F.-L., Zhang, X.-B., and Han, Z.-X. A fluorescent chemical sensor for Hg(II) based on a corrole derivative in a PVC matrix. *Talanta* 70(2) (2006): 364-369.
- [42] Kuswandi, B., Nuriman, Dam, H.H., Reinhoudt, D.N.; and Verboom, W. Development of a disposable mercury ion-selective optode based on trityl-picolinamide as ionophore. *Analytica Chimica Acta* 591(2) (2007): 208-213.
- [43] Amini, M.K., Khezri, B., and Firooz, A.R. Development of a highly sensitive and selective optical chemical sensor for batch and flow-through determination of mercury ion. *Sensors and Actuators B: Chemical* 131(2) (2008): 470-478.
- [44] Seiler K, W.K., Bakker E, Morf WE, Rusterholz B, Spichiger UE, Simon W. Characterization of sodium-selective optode membranes based on neutral ionophores and assay of sodium in plasma. *Clin Chem.* 37(8) (1991): 1350-1355.
- [45] Yang, X., Wang, K., and Guo, C. A fluorescent optode for sodium ion based on the inner filter effect. *Analytica Chimica Acta* 407(1-2) (2000): 45-52.
- [46] Albero, M.I., Ortúño, J.A., García, M.S., Sánchez-Pedreño, C., and Expósito, R. Determination of zinc (II) in pharmaceuticals based on a flow-through bulk optode. *Journal of Pharmaceutical and Biomedical Analysis* 29(5) (2002): 779-786.
- [47] Shamsipur, M., Rouhani, S., Mohajeri, A., and Ganjali, M. PVC-membrane ion-selective bulk optode for Ag⁺ ion based on hexathia-18-crown-6 and 1,2-benzo-3-octadecanoylimino-7-diethylaminophenoxyazine. *Analytical and Bioanalytical Chemistry* 375(5) (2003): 692-697.
- [48] Shamsipur, M., Alizadeh, K., Hosseini, M., Caltagirone, C., and Lippolis, V. A selective optode membrane for silver ion based on fluorescence quenching



3622910427

- of the dansylamidopropyl pendant arm derivative of 1-aza-4,7,10-trithiacyclododecane ([12]aneNS₃). *Sensors and Actuators B: Chemical* 113(2) (2006): 892-899.
- [49] Wygladacz, K., Radu, A., Xu, C., Qin, Y., and Bakker, E. Fiber-optic microsensor array based on fluorescent bulk optode microspheres for the trace analysis of silver ions. *Analytical Chemistry* 77(15) (2005): 4706-4712.
- [50] Lerchi, M., Reitter, E., Simon, W., Pretsch, E., Chowdhury, D.A., and Kamata, S. Bulk optodes based on neutral dithiocarbamate ionophores with high selectivity and sensitivity for silver and mercury cations. *Analytical Chemistry* 66(10) (1994): 1713-1717.
- [51] Lerchi, M., Orsini, F., Cimerman, Z., Pretsch, E., Chowdhury, D.A., and Kamata, S. Selective optical sensing of silver ions in drinking water. *Analytical Chemistry* 68(18) (1996): 3210-3214.
- [52] Murkovic, I., Oehme, I., Mohr, G., Ferber, T., and Wolfbeis, O. Optode membrane for continuous measurement of silver ions. *Microchimica Acta* 121(1-4) (1995): 249-258.
- [53] Firooz, A.R., Ensafi, A.A., Kazemifard, N., and Sharghi, H. A highly sensitive and selective bulk optode based on benzimidazol derivative as an ionophore and ETH5294 for the determination of ultra trace amount of silver ions. *Talanta* 101 (2012): 171-176.
- [54] Firooz, A.R., Ensafi, A.A., Kazemifard, N., and Khalifeh, R. Development of a highly sensitive and selective optical sensor for determination of ultra-trace amount of silver ions. *Sensors and Actuators B: Chemical* 176 (2013): 598-604.
- [55] Kometani, N., Tsubonishi, M., Fujita, T., Asami, K., and Yonezawa, Y. Preparation and optical absorption spectra of Dye-Coated Au, Ag, and Au/Ag colloidal nanoparticles in aqueous solutions and in alternate assemblies. *Langmuir* 17(3) (2001): 578-580.
- [56] Bühlmann, P. and Chen, L.D. Ion-selective electrodes with ionophore-doped sensing membranes. in *Supramolecular Chemistry*: John Wiley & Sons, Ltd, 2012.
- [57] Yew, P.L. and Heng, L.Y. A reflectometric ion sensor for potassium based on acrylic microspheres. *Sensors and Actuators B: Chemical* 191 (2014): 719-726.
- [58] Seiler, K. and Simon, W. Theoretical aspects of bulk optode membranes. *Analytica Chimica Acta* 266(1) (1992): 73-87.
- [59] Alberto, R., et al. Silver(I) complexes of the derivatized crown thioether ligands 3,6,9,12,15,18-hexathianonadecanol and 3,6,9,13,16,19-



- hexathiaicosanol. determination of stability constants and the crystal structures of $[\text{Ag}(19\text{-aneS}_6\text{-OH})][\text{CF}_3\text{SO}_3]$ and $[\text{Ag}(20\text{-aneS}_6\text{-OH})][\text{BF}_4]$. Inorganic Chemistry 35(11) (1996): 3420-3427.
- [60] Capitán-Vallvey, L.F., Fernández Ramos, M.D., and Alvarez de Cienfuegos-Gálvez, P. Optical test strip for calcium determination based on a neutral ionophore. Analytica Chimica Acta 451(2) (2002): 231-241.
- [61] Capitán-Vallvey, L.F., Lapresta-Fernández, A., Fernández-Ramos, M.D., and Cuadros-Rodríguez, L. Establishment of wide-range linear response curves in bulk optode sensors for cations based on ion exchange. Sensors and Actuators B: Chemical 117(1) (2006): 27-34.
- [62] Shamsipur, M., et al. Development of a new fluorimetric bulk optode membrane based on 2,5-thiophenylbis(5-tert-butyl-1,3-benzoxazole) for nickel(II) ions. Analytica Chimica Acta 501(1) (2004): 55-60.
- [63] Huber, L. Validation and qualification in analytical laboratories. Illinois: Interharm Press, 1999.



VITA

Miss Rawiwan Wattanayon was born on January 6, 1988 in Narathiwat, Thailand. She received her Bachelor's Degree of Science from Department of Chemistry, Faculty of Science, Chulalongkorn University in 2009. After that, she continues to study further as a graduate student at the Department of Chemistry, Faculty of Science, Chulalongkorn University and become one of the members of Environmental Analysis Research Unit (EARU). The present address is 80/2 Jaturonggrassamee Road, Bangnak, Muang Narathiwat, Narathiwat, Thailand, 96000. Contact number is 086-5125862.

3622910427

