

References

- Aggarwal, K. 1988. Incidence of *Tropilaelaps clareae* on three *Apis* species in Hisar (India). In G.R. Needham, R. E. Page, M. Delfinado-Baker and C. E. Bowman (eds.), Africanized honey bees and bee mites, New York, pp. 396-403.
- Aggarwal, K., and Kapil, R. P. 1986. Seasonal population dynamics of *Tropilaelaps clareae* (Acar: Laelapidae) in *Apis dorsata* colonies. In G. P. Channabasavanna and C. A. Viraktamath (eds.), Progress in acarology, New Delhi.
- Akratanakul, P. 1979. Occurrence of a parasitic bee mite *Tropilaelaps clareae* (Acar: Gamasida) in Thailand. BRL's Technical Report No.1/1979.
- Akratanakul, P. 1987. Honeybee diseases and enemies in Asia: a practical guild. FAO Agricultural services Bulletin 68/5, pp. 17-31.
- Anderson, D. L. 1994. Non-reproduction of *Varroa jacobsoni* in *Apis mellifera* colonies in Papua New Guinea and Indonesia. Apidologie 25: 412-421.
- Atwal, A. S., and Goyal, N. P. 1971. Infestation of honey bee colonies with *Tropilaelaps* and its control. J. Apic. Res. 10: 137-142.
- Bharadwaj, R. K. 1968. A new record of the mite *Tropilaelaps clareae* from *Apis dorsata* colonies. Bee Wld. 49(3): 115.
- Boecking, O., and Drescher, W. 1990. The reaction of worker bees in different *Apis mellifera* colonies to *Varroa* infested brood cells. Proceedings of the international symposium on recent research on bee pathology, Gent, Belgium, pp. 41-42.
- Boecking, O., and Drescher, W. 1991. Response of *Apis mellifera* L. colonies infested with *Varroa jacobsoni* Oud. Apidologie 22: 237-241.
- Boecking, O., Rath, W., and Drescher, W. 1992. *Apis mellifera* removes *Varroa jacobsoni* Oud. and *Tropilaelaps clareae* Delfinado & Baker from sealed brood in the tropics. Am. Bee J. 132: 732-734.
- Borchers, H. A. 1964. Effects of certain factors on the nest-cleaning behavior of honey bees (*Apis mellifera*). Master's Thesis. Ohio State University.
- Buchler, R., and Drescher, W. 1990. Variance and heritability of the capped developmental stage in European *Apis mellifera* L. colonies and its correlation with increased *Varroa jacobsoni* Oud. infestation. J. Apic. Res. 29: 172-176.

- Buchler, R., Drescher, W., and Tornier, I. 1992. Grooming behavior of *Apis cerana*, *Apis mellifera* and *Apis dorsata* and its effect on the parasitic mites *Varroa jacobsoni* and *Tropilaelaps clareae*. Exp. Appl. Acarol. 16, 313-319.
- Burgett, D. M., and Akratanakul, P. 1985. *Tropilaelaps clareae* the little known honey bee brood mite, Am. Bee J. 125, 112-114.
- Burgett, D. M., Akratanakul, P., and Morse R. A. 1983. *Tropilaelaps clareae*: a parasite of honeybees in southeast Asia, Bee Wld. 64, 25-28.
- Burgett, D. M., and Burikam, I. 1985. Number of adult honey bees (Hymenoptera: Apidae) occupying a comb: A standard for estimating colony populations. J. Econ. Entomol. 78: 1154-1156.
- Burgett, D. M., and Kitprasert, C. 1989. *Tropilaelaps clareae* Delfinado and Baker parasitism in relationship with its natural host *Apis dorsata* F. Proceedings of the first Asia-Pacific conference of Entomology, Chiang Mai, Thailand, p. 282.
- Burgett, D. M., and Krantz, G. W. 1984. The future of the European honey bee (*Apis mellifera* L.) in Southeast Asia: constraints of parasitism. Proceedings of the expert consultation on beekeeping with *Apis mellifera* in tropical and sub-tropical asia. Chiang Mai, Thailand. pp. 34-43.
- Burgett, D. M., and Rossignol, P. A. 1990. A model of dispersion and regulation of brood mite (*Tropilaelaps clareae*) parasitism on the giant honey bee (*Apis dorsata*). Can. J. Zool. 68: 1423-1427.
- Chen, Y. 1993. Apiculture in China. Beijing: Agricultural publishing housing.
- Crane, E. 1968. Mites infesting honeybees in Asia. Bee Wld. 49(3): 113-114.
- Crane, E. 1978. Beekeeping round the world. Bee Wld. 59(4): 164-167.
- Crane, E. 1990. Bees and beekeeping. Ithaca, New York: Comstock Publishing Associates
- Danka, R.G., Rinderer,T.E., Kuznetsov,V.N., and Delatte, G.T. 1995. A USDA-ARS project to evaluate resistance to *Varroa jacobsoni* by honey bees of Far-eastern Russia. Am. Bee J. 135(11): 746-748.
- De Guzman, L. I., Rinderer, T. E., Delatte, G. T., and Macchiavelli, R. E. 1996. *Varroa jacobsoni* Oudemans tolerance in selected stocks of *Apis mellifera* L. Apidologie 27: 193-210.

- De Guzman, L. I., Rinderer, T. E., Delatte, G. T., Stelzer, J. A., Beaman, L. and Harper, C. 2001. An evaluation of Far-eastern Russia honey bees another methods for the control of Tracheal mites. Am. Bee J. 141 (10): 737-741.
- De Guzman, L. I., Rinderer, T. E., Stelzer, J. A., Beaman, L., Delatte, G. and Harper, C. 2001. Hygienic behavior by honey bees from Far-eastern Russia. Am. Bee J. 141: 58-60.
- Delfinado-Baker, M. 1963. Mites of the honeybee in south-east Asia. J. Apic. Res. 2: 113-114.
- Delfinado-Baker, M., and Aggarwal, K. 1987. A new *Varroa* (Acari: Varroidae) from the nest of *Apis cerana* (Apidae). Int. J. Acarol. 13: 233-237.
- Delfinado-Baker, M. and Baker, E. W. 1961. *Tropilaelaps*, a new genus of mite from the Philippines (Laelapidae: Acarina). Fieldiana Zoology 44(7): 53-54.
- Delfinado-Baker, M. and Baker, E. W. 1982. A new species of *Tropilaelaps* parasitic on honey bees. Am. Bee J. 122: 416-417.
- Delfinado-Baker, M., Baker, E. W., and Phoon, A. C. G. 1989. Mites (Acari) associated with bees (Apidae) in Asia, with description of a new species, Am. Bee J. 129, 609-610, 612-613.
- Delfinado-Baker, M., and Peng, C. Y. S. 1995. *Varroa jacobsoni* and *Tropilaelaps clareae*: A perspective of life history and why Asian bee-mites preferred European honey bees. Am. Bee J. 135: 415-420.
- Delfinado-Baker, M., Rath, W., and Boecking, O. 1992. Phoretic bee mites and honeybee grooming behavior. Int. J. Acarol. 18(4): 315-322.
- Delfinado-Baker, M., Underwood, B. A., and Baker, E.W. 1985. The occurrence of *Tropilaelaps* mites in brood nests of *Apis dorsata* and *A. laboriosa* in Nepal, with descriptions of the nymphal stages.¹ Am. Bee J.: 703-706.
- De Jong, D., Morse, R .A., and Eickwort, G .C. 1982. Mite pests of honey bees. Ann. Rev. Entomol. 27: 229-252.
- Diwan, V. V., and Salvi, S.R. 1965. Some interesting behavioral features of *Apis dorsata* Fab. Indian Bee J. 27: 52.
- Dyer, F. C. 1985. Nocturnal orientation by the Asian honeybee; *Apis dorsata*. Anim. Behav. 33: 769-779.

- Gautam, S. K. 1984. Country status report on beekeeping in Nepal (discussion). Proc. Expert Consult Bangkok, FAO Rome: 198-202.
- Gilliam, M., Taber, S. I., and Richardson, G. V. 1983. Hygienic behavior of honey bees in relation to chalkbrood disease. Apidologie 14: 29-39.
- Hanel, H. 1983. Effect of JH-III on the reproduction of *Varroa jacobsoni*. Apidologie 14:137-142.
- Hanel, H. 1986. Effect of Juvenile hormone (III) from the host *Apis mellifera* (Insecta: Hymenoptera) on the neurosecretion of the parasite *Varroa jacobsoni* (Acari: Mesostigmata). Exp. Appl. Acarol. 2: 257-271.
- Hanel, H., and Koeniger, N. 1986. Possible regulation of the reproduction of the honey bee mite *Varroa jacobsoni* (Mesostigmata: Acari) by a host's hormone: juvenile hormone III. J. Insect Physiol. 32: 791-798.
- Harbo, J. R., and Harris, J. W. 1999. Heritability in honey bees (Hymenoptera: Apidae) of characteristics associated with resistance to *Varroa jacobsoni* (Mesostigmata: Varroidae). J. Econ. Entomol. 92(2): 261-265.
- Harris, J. W., and Harbo, J. R. 2001. Natural & suppressed reproduction of Varroa. Bee Culture. : 34-38.
- Hoppe, H., Stephen, E., and Muhammad, F. 1989. Formic acid; A highly efficient treatment against the external parasitic brood mite *Tropilaelaps clareae*. Proc. 4 int. Conf. Apic. Climates, Cairo, 1988: p. 369.
- Kallapur, S. K. 1950. An experiment in the collection of wild honey. Indian Bee J. 12: 122-124.
- Kavinseksan, B., Wongsiri S., de Guzman L. I., and Rinderer T. E. 2003. Absence of *Tropilaelaps* infestation from recent swarms of *Apis dorsata* in Thailand. J. Apic. Res. 49-50.
- Kavinseksan, B., Wongsiri, S., de Guzman, L. I., and Rinderer, T. E. (in Press b) Population and sex ratio of *Tropilaelaps clareae* in debris and on adult bees of *Apis dorsata* in Thailand. Apidologie.
- Kavinseksan, B., Wongsiri, S., Rinderer, T. E., and de Guzman, L. I. (in Press a) Comparison of the hygienic behavior of Thai commercial and ARS Russian honey bees. Science Asia.

- Kitprasert, C. 1984. Biology and systematics of the parasitic bee mite, *Tropilaelaps clareae* Delfinado and Baker (Acarina : Laelapidae). M. S. thesis, Kasetsart University, Thailand. (In Thai)
- Krantz, G. W. 1978. A manual of Acarology. Second edition. Oregon State University Book Stores INC., USA.
- Kraus, B., and Velthuis, H. H. W. 1997. High humidity in the honey bee (*Apis mellifera* L.) brood nest limits reproduction of the parasitic mite *Varroa jacobsoni* Oud. Naturwissenschaften 84: 217-218.
- Koeniger, G., Koeniger, N., Anderson, D.L., Lekprayon, C., and Tingek, S. 2002. Mites from debris and sealed brood cells of *Apis dorsata* colonies in Sabah (Borneo) Malaysia, including a new haplotype of *Varroa jacobsoni*. Apidologie 33: 15-24.
- Koeniger, N., and Koeniger, G. 1980. Observations and experiments on migration and dance communication of *Apis dorsata* in Sri Lanka. J. Apic. Res. 19(1): 21-34.
- Koeniger, N., Koeniger, G., Mardan, M., and Wongsiri, S. 1993. Possible effects of regular treatments of varroatosis on the host-parasite relationship between *Apis mellifera* and *Varroa jacobsoni*. In L. J. Connor., T. Rinderer., H. A. Sylvester and S. Wongsiri (eds.), Asian apiculture. Cheshire, Connecticut. pp. 541-550.
- Koeniger, N., and Muzaffar, N. 1988. Lifespan of the parasitic honeybee mite *Tropilaelaps clareae*, on *Apis cerana*, *dorsata* and *mellifera*. J. Apic. Res. 27: 207-212.
- Laigo, F. M. and Morse, R. 1968. The mite *Tropilaelaps clareae* in *Apis dorsata* colonies in the Philipines. Bee Wld. 49(3): 116-118.
- Le Conte, Y., Arnold, G., and Desenfant, P. 1990. Influence of brood temperature and hygrometry variation on the development of the honey bee ectoparasite *Varroa jacobsoni* (Mesostigmata: Varroidae). Environmental Entomology 19: 1780-1785.
- Lensky, Y., Dechmani, U., Jirasavetakul, U., and Tiewtrakul, S. 2001. The effect of flualinate on *Varroa jacobsoni* (Oud.) and *Tropilaelaps clareae* (Delfinado & Baker) in *Apis mellifera* colonies in north Thailand. Proceedings of the third asian apicultural association conference on bee research and beekeeping development, Hanoi, Vietnam, 1996: pp. 176-182.

- Lindauer, M. 1961. Communication among social bees. Harvard University Press. Cambridge.
- Maa, T. C. 1953. An inquiry into the systematics of the Tribus Apidini or honey bees (Hymenoptera). Treubia 21: 525-640.
- Mardan, M. 1989. Thermoregulation in the Asiatic giant honeybee *Apis dorsata* F (Hymenopter: Apidae). Ph.D. dissertation. University of Guelp, In G. W. Otis (ed.), 1991. A review of the diversity of species within Apis. pp. 29-47.
- Martin, S. J. 1995. Reproduction of *Varroa jacobsoni* in cells of *Apis mellifera* containing one or more mother mites and the distribution of these cells. J. Apic. Res. 34: 187-196.
- McDonald, J. L. 1971. Beekeeping in Taiwan, Republic of China. Am. Bee J. 111(2) : 58-60.
- Momot, J. P., and Rothenbuhler, W. C. 1971. Behaviour genetics of nest cleaning in honey bees. VI. Interactions of age and genotype of bees, and nectar flow. J. Apic. Res. 10: 11-21.
- Moritz, R. F. A., Kryger, P., Koeniger, N., Estoup, A., and Tingek, S. 1995 High degree of polyandry in *Apis dorsata* queens detected by DNA microsatellite variability. Behav. Ecol. Sociobiol. 37: 357-363.
- Morse, R. A., and Laigo, F. M. 1969. Apis dorsata in the Philippines. Monogr. Philipp. Associa. Entomol. 1: pp.1-96.
- Morse, R. A., Shearer, D. A., Boch, R., and Benton, A. W. 1969. Observations on alarm substances in the genus *Apis*. J. Apic Res. 6: 113-118.
- Muttoo, R. N. 1956. Facts about beekeeping in India. Bee Wld. 37: 125-133, 154-157.
- Nixon, M. 1983. World maps of *Varroa jacobsoni* and *Tropilaelaps clareae*, with additional records for honeybee diseases and parasites previously mapped. Bee Wld. 64(3): 124-131.
- Nyein, M. M., and Zmarlicki, C. 1982. Control of mites in European bees in Burma. Am. Bee J. 122: 638-639.
- Oldroyd, B., Rinderer, T. E., and Wongsiri, S. (1992). Pollen resource partitioning by *Apis dorsata*, *A. cerana*, *A. andreniformis* and *A. florea* in Thailand. J. Apic. Res. 31: 3-7.

- Otis, W. G. and Kralj, J. 2001. Parasitic brood mites not present in North America. In C. W. Thomas and S. D. Keith (eds.), Mites of the Honey Bee. Hamilton, Illinois: Dadant & Sons, 280 pp.
- Park, O. W. 1937. Testing for resistance to American foulbrood in honey bees. J. Econ. Entomol. 30(3): 504-512.
- Peng, Y. S., Fang, Y., Xu, S., and Ge, L. 1987. The resistance mechanism of the Asian honey bee, *Apis cerana* Fabr. to an ectoparasite, *Varroa jacobsoni* Oudemans. J. Invertebr. Pathol. 49: 54-60.
- Pianka, E. R. 1983. Evolutionary ecology. Third edition. New York: Harper & Row, Publishers.
- Qayyum, H. A., and Nabi, A. 1968. Biology of *Apis dorsata* F. Pak. J. Sci. 19: 109-113.
- Rajesh, G., Sharma, O. P., and Dogra, G. S. 1984. Formic acid: an effective acaricide against *Tropilaelaps clareae* Delfinado and Baker (Laelaptidae: Acarina) and its effect on the brood and longevity of honey bees. Am. Bee J.: 736-738.
- Rath, W. 1991. Investigation on the parasite mites *Varroa jacobsoni* Oud. and *Tropilaelaps clareae* Delfinado & Baker and their hosts *Apis cerana* Fabr., *Apis dorsata* Fabr. and *Apis mellifera* L., Doctoral Thesis, Matematisch Naturwissen. Fakultat der Rheinsischen Friedrich Wilhelms Universitat Bonn, Germany, 148 pp.
- Rath, W., and Delfinado-Bake, M. 1990. Analysis of *Tropilaelaps clareae* populations from the debris of *Apis dorsata* and *Apis mellifera* in Thailand. Proceedings of the apimondia symposium recent research on bee pathology, Gent, Belgium, pp. 86-89.
- Rath, W., Delfinado-Baker, M., and Drescher, W. 1991. Observations on the mating behavior, sex ratio, phoresy and dispersal of *Tropilaelaps clareae* (Acari: Laelapidae). Int. J. Acarol. 17: 201-208.
- Rembold, H., Kremer, J. P., and Ulrich, G. M. 1980. Charaterization of postembryonic developmental stages of the female castes of the honey bee, *Apis mellifera* L. Apidologie 11(1): 29-38.
- Rinderer, T. E, de Guzman, L. I., Harris, J., Kuznetsov, V., Delatte, G. T., Stelzer, J. A., and Beaman, L. 2000. The release of ARS Russian honey bees. Am. Bee J. 140 (4): 305-307.

- Rinderer, T. E., Delatte, G. T., De Guzman, L. I., Williams, J., Stelzer, J., and Kuznetsov, V. N. 1999. Evaluations of the *Varroa*-resistance of the honey bees imported from Far-Eastern Russia. Am. Bee J. 139(4): 287-290.
- Rinderer, T. E., Kuznetsov, V. N., Danka, R. G. and Detatte, G. T. 1997. An importation of potentially *Varroa* - resistant honey bees from Far-Eastern Russia. Am. Bee J. 137(11): 787-789.
- Rinderer, T. E., Oldroyd, P. B., Lekprayoon, C., Wongsiri, S., Boonthai, C., and Thapa, R. 1994. Extended survival of the parasitic honey bee mite *Tropilaelaps clareae* on adult workers of *Apis mellifera* and *Apis dorsata*. J. Apic. Res. 33(3): 171-174.
- Rinderer, T. E. et al. 2001. Multi-state field trials of ARS Russian honey bees: 1. Responses to *Varroa destructor* 1999, 2000. Am. Bee J. 141(9): 658-661.
- Rittter, W. 1993. What future for biotechnical control? In: A. Matheson (ed.), Living with *Varroa*. Proceeding of an IBRA symposium, London, 21 November 1992 : pp. 25-31.
- Ritter, W., and Schneider-Ritter, U. 1988. Difference in biology and means of controlling *Varroa jacobsoni* and *Tropilaelaps clareae*, two novel parasitic mites of *Apis mellifera*, In G.R. Needham, R. E. Page, M. Delfinado-Baker and C. E. Bowman (eds.), Africanized honey bees and bee mites, New York, pp. 387-395.
- Rosenkranz, P., and Engels, W. 1994. Infertility of *Varroa jacobsoni* females aftter invasion into *Apis mellifera* worker brood as a tolerance factor against varroatosis. Apidologie 25: 402-411.
- Rosenkranz, P., Rachinsky, A., Strambi, C., Strambi, A., Schricker, B., Ropstorf, P., and Paulino, S. Z. L. 1989. Juvenile hormone titer in capped L5 larvae of various races of honeybee. Apidologie 20: 524-526.
- Rosenkranz, P., Tewarson, N. C., Rachinsky, A., Strambi, A., Strambi, C., and Engels, W. 1993. Juvenile hormone titer and reproduction of *Varroa jacobsoni* in capped brood stages of *Apis cerana indica* in comparison to *Apis mellifera ligustica*. Apidologie 24: 375-382.
- Rothenbuhler, W. C. 1964. Behavior genetics of nest cleaning in honey bees: IV. Responses of F₁ and backcross generations to disease-killed brood. Am. Zoologist 4: 111-123.

- Ruttner, F. 1988. Biogeography and taxonomy of honeybees. Berlin: Springer, 284 pp.
- Ruttner, F., and Hanel, H. 1992. Active defense against *Varroa* mites in a carniolan strain of honeybee (*Apis mellifera carnica* Pollman). Apidologie 23: 173-178.
- Sakagami, Sh. F., Matsumura, T., and Ito, K. 1980. *Apis laboriosa* in Himalayas. The little know Worlds' largest honey bee (Hymenoptera: Apidae). Insecta Matsumurana 19: 47-77.
- Seeley, T. D., Seeley, R. H., and Akratanakul, P. 1982. Colony defense strategies of the honey bee in Thailand. Ecol. Monogr. 52: 43-63.
- Spivak, M., and Gilliam, M. 1993. Facultative expression of hygienic behavior in honey bees in relation to disease resistance. J. Apic. Res. 32: 147-157.
- Spivak, M., and Reuter, G. S. 1998. Honey bee hygienic behavior. Am. Bee J. 138(4): 283-286.
- Spivak, M., and Reuter, G. S. 2001. Comparing hygienic bees. Canadian Honey Council 14(3): 25.
- Stephen, W. A. 1968. Mites: A beekeeping problem in Vietnam and India. Bee Wld. 49: 119-120.
- Tangkanasing, P., Wongsiri, S., and Vongsamanod, S. 1988. Integrated control of *Varroa jacobsoni* and *Tropilaelaps clareae* in bee hives in Thailand. In G. R. Needham, R. E. Page, M. Delfinado-Baker and C. E. Bowman (eds.), Africanized Honey Bee and Bee Mites. New York: John Wiley & Sons, pp. 409-412.
- Thakar, C. V., and Tonapi, K. V. 1961. Nesting behavior of Indian honey bees I. Differentiation of worker, queen and drone cells of the combs of *Apis dorsata* Fab. Bee Wld. 42: 61-62,72.
- Thapa, R. B. 1998. Colony Migration of the Giant Honeybee, Apis dorsata Fabr. Ph.D.dissertation. Chulalongkorn University. 102 pp.
- Thapa, R. B., and Wongsiri, S. 2003. Flying predators of the giant honey bees; *Apis dorsata* and *Apis laboriosa* in Nepal. Am. Bee J. 143(7) : 540-542.
- Thomas, H., U. 1997. Practical aspects of alternative *Varroa* control methods. In: P. Munn and R. Jones (eds.), *Varroa! Fight The Mite*. International Bee Research Association, Cardiff: 22-30 pp.

- Thrybom, B. and Fries, I. 1991. Development of infestations by *Varroa jacobsoni* in hybrid colonies of *Apis mellifera monticola* and *Apis mellifera ligustica*. J. Apic.Res. 30:151-155.
- Trump, R. F., Thomson, V. C., and Rothenbuhler, W. C. 1967. Behaviour genetics of nest cleaning in honey bees. V. Effect of previous experience and composition of mixed colonies on response to disease-killed brood. J. Apic. Res. 6(3): 127-131.
- Tubbs, H., Harper, C., Bigalk, M., Bernard, S. J., Delatte, G. T., Sylvester, H. A., and Rinderer, T. E. 2003. Commercial management of ARS Russian honey bees. Am. Bee J. 144: 819-820.
- Underwood, B. A. 1986. The natural history of Apis laboriosa Smith in Nepal. M.S. thesis, Cornell University, Ithaca, New York.
- Wilson, E. O. 1971. The Insect Societies. Cambridge, Mass: Harvard University Press.
- Wongsiri, S., and Chen, P. 1995. Effects of agricultural development on honey bees in Thailand. Bee Wld. 76(1): 3-5.
- Wongsiri, S., and Tangkanasing, P. 1987. Bee mites control in Thailand. In Workshop on parasitic bee mites and their control, Pulawy, Poland, 27 August-1 September 1987.
- Wongsiri, S., Tangkanasing, P., and Sylvester, H. A. 1989. The resistance behavior of *Apis cerana* against *Tropilaelaps clareae*. Proceedings of the First Asia-Pacific Conference of Entomology, Chaing Mai, Thailand: pp. 828-836.
- Wongsiri, S., Tangkanasing, P., and Vongsamanodes, S. 1987a. Effectiveness of Asuntol® (coumaphos), Perizin® (coumaphos), Mitac® (amitraz) and powder of sulphur with naphthalene for the control of bee mites(*Varroa jacobsoni* and *Tropilaelaps clareae*) in Thailand. In Proceedings of the XXXI st International Apicultural Congress of Apicultural Congress of APIMONDIA, Warsaw, Poland, pp. 322-325.
- Wongsiri, S., Tangkanasing, P., and Sylvester, H. A. 1987b. Mites, pests and beekeeping with *Apis cerana* and *Apis mellifera* in Thailand. Am. Bee J. 127(7): 500-503.
- Wongsiri, S., Thapa, R., and Kongpitak, P. 1998. Longan: a major honey plant in Thailand. Bee Wld. 79(1): 23 -28.

- Wongsiri, S., Thapa, R., Oldroyd, B. P., and Burgett, M. D. 1996. A magic bee tree: Home of *Apis dorsata*. Am. Bee J. 136 (11): 196-199.
- Woo, K. S., and Lee, J. H. 2001. Current status of honey bee mites in Korea. Proceedings of the Third Asian Apicultural Association Conference on Bee Research and Beekeeping Development, Hanoi, Vietnam, 1996: pp. 168-171.
- Woyke, J. 1984. Survival and prophylactic control of *Tropilaelaps clareae* infesting *Apis mellifera* colonies in Afghanistan. Apidologie 15(4): 421-434.
- Woyke, J. 1985a. *Tropilaelaps clareae*, a serious pest of *Apis mellifera* in the Tropics but not dangerous for apiculture in temperate zones. Am. Bee J. 125(7): 497-499.
- Woyke, J. 1985b. Further investigations into control of the parasite bee mite *Tropilaelaps clareae* without medication. J. Apic. Res. 24(4): 250-254.
- Woyke, J. 1987a. Length of successive stages in the development of the mite *Tropilaelaps clareae* in relation to honeybee brood age. J. Apic. Res. 26(2): 110-114.
- Woyke, J. 1987b. Length of stay of the parasitic mite *Tropilaelaps clareae* outside sealed honeybee brood cells as a basis for its effective control. J. Apic. Res. 26 (2): 104-109.
- Woyke., J. 1989. Change in shape of *Tropilaelaps clareae* females and the onset of egg laying. J. Apic. Res. 28(4): 196-200.
- Woyke, J. 1990. Biology and control of the parasitic bee mite *Tropilaelaps clareae*, Proceedings of the apimondia symposium recent research on bee pathology, Gent, Belgium, pp. 90-99.
- Woyke, J., Wilde, J., and Wilde, M. 2000. Swarming, migrating and absconding of *Apis dorsata* colonies. Proceedings of the seventh international conference on tropical bees: management and diversity & fifth asian apicultural association conference. pp. 183-188.
- Woyke, J. 2001. Different reaction of *Apis dorsata* and *Apis mellifera* to brood infestation by parasite mites. Proc. 3rd AAA Conf. on Bee Res. and Beekeeping Dev., Hanoi, Vietnam, 1996, pp. 172-175.

Wrensch D. L., and Young, S. S. Y. 1983. Relationship between primary and tertiary sex ratio in the two-spotted spider mite (Acarina: Tetranychidae). Ann. Entomol. Soc. Am. 76: 786-789.



ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย



APPENDIX

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Absence of *Tropilaelaps* infestation from recent swarms of *Apis dorsata* in Thailand



B KAVINSEKSA¹, *S WONGSIRI¹, L I DE GUZMAN² AND T E RINDERER²

¹Bee Biology Research Unit, Department of Biology, Faculty of Science, Chulalongkorn University, Phayathai Rd., Phatumwan, Bangkok 10330, Thailand

²USDA-ARS, Honey Bee Breeding, Genetics and Physiology Laboratory, 1157 Ben Hur Road, Baton Rouge, Louisiana 70820-5502, USA

Received 22 April 2002, accepted subject to revision 24 June 2002, accepted for publication 9 June 2003

Keywords: *Tropilaelaps clareae*, *Apis dorsata*, honey bees, parasitic mites, Thailand

Tropilaelaps clareae is a brood parasite of *Apis dorsata*. *A. dorsata* colonies tend to migrate during dearth periods and stop brood rearing in preparation for migration. This means that during migrations there is a period of broodlessness. *T. clareae* cannot survive more than 3 days on adult bees of *A. dorsata* (Rinderer et al., 1994), thus the origin of infestations in colonies that have recently undergone migration is as yet unknown. Presumably populations of *T. clareae* gradually build up in new *A. dorsata* nests after re-infestation. To test this hypothesis, we evaluated *Tropilaelaps* infestations in new and established nests of *A. dorsata* in Samut Songkhram, Thailand.

Five new (swarms or colonies with the first generation of brood) and five established (with more than the first generation of brood) colonies of *A. dorsata* were examined for the presence of *Tropilaelaps* between April 2000 and September 2002. For the new colonies, the whole nests (adult bees and brood combs containing unsealed and sealed brood) were collected and examined. We inferred the age of the new colonies from the stage of the brood present. Colony 1 was a swarm (without comb),

which had arrived at the site about eight days before being sampled. Colonies 2 and 3 were nine- and 13-days-old respectively, while colonies 4 and 5 were both 14-days-old. The established colonies were all more than two months old. The presence of *Tropilaelaps* in these established colonies was determined by cutting a brood comb section (about 10 × 18 cm) from each of the colonies. Both unsealed and sealed brood within the cut sections was examined (table 1). The entire nest of colonies 4 and 5 were collected and examined for the presence of mites.

A sample of adult bees ($n = 57\text{--}9782$) was obtained from each colony. These were washed in 70% ethanol to remove any mites present. The bees were then removed with a sieve, and mites counted in the alcohol.

T. clareae was not found in the five newly founded colonies (table 1). This observation suggests that *A. dorsata* colonies start with uninfested bees. This mite-free condition is probably due to broodlessness during the swarming events that interrupt the mite's life cycle. This period can be very lengthy. Koeniger & Koeniger (1980) reported that *A. dorsata* swarms fly in several

*Corresponding author: kavinseksan@yahoo.com

TABLE 1. Numbers of *Tropilaelaps clareae* adults in new and established colonies of *Apis dorsata* in Samut Songkram, Thailand.

Colony type	Colony number	No. of unsealed brood examined	No. of sealed brood examined	No. of adult bees examined
New	1	No comb	No comb	9 782 (0)
	2	1 262 (0)	No sealed brood	5 055 (0)
	3	1 440 (0)	620* (0)	5 181 (0)
	4	1 281 (0)	425* (0)	3 767 (0)
	5	394 (0)	No sealed brood	3 930 (0)
Total		4 377 (0)	1 045 (0)	27 715 (0)
Established	1	200 (0)	241 (16)	57 (0)
	2	173 (0)	313 (15)	521 (0)
	3	133 (0)	443 (13)	167 (0)
	4	341 (0)	3 366 (10)	1 232 (0)
	5	529 (0)	3 413 (5)	5 056 (9)
Total		1 376 (0)	7 776 (59)	7 033 (9)

Numbers inside () indicate numbers of detected mites. * Sealed larvae

stages and often rest between stages of the flight. They showed that swarms often rest one to three days without building any comb and can spend more than one month on their migration. In our study, colony 1 did not commence comb building for eight days after its arrival at the nest site, providing a further period of broodlessness. The extensive period of broodlessness suggests that any mites travelling phoretically on a swarm are likely to die during swarming events before having the chance to enter brood cells of the new nest. This suggestion is supported by the absence of *T. clareae* in the new colonies. In Borneo, Koeniger *et al.* (2002) observed infestations of *T. clareae* in a newly settled swarm that had originated from an infested colony nearby. This shows that *T. clareae* can spread via reproductive swarms, but probably not, as we have shown, in migratory swarms.

We confirmed that *T. clareae* populations gradually build up in new *A. dorsata* nests after re-infestation. The five established colonies had 5–16 mites in the sampled sealed brood, and nine mites were collected from adult bees of colony 5 only. Most often uninhabited colonies must acquire *T. clareae* from other *Apis* species or nests of *A. dorsata* present in the area. In south-east Asia *T. clareae* has proved to be a more serious pest of *A. mellifera* than either *Varroa destructor* or *V. jacobsoni*. Since brood production in *Apis mellifera* colonies is year-round in Thailand, these and non-migrating *A. dorsata* colonies may serve as constant reservoirs of *Tropilaelaps* for new colonies. *T. clareae* is also known to reproduce in *A. cerana* drone brood, and to be phoretic on adult bees of *A. dorsata*, *A. laboriosa*, *A. mellifera*, *A. florea* and *A. cerana* (Aggarwal 1988). In Samut Songkram, all of these bee species except *A. laboriosa* coexist. *A. cerana* and *A. dorsata* have been reported to forage simultaneously on the same inflorescence of king palm in Thailand (Oldroyd *et al.*, 1992). In Malaysia, *A. cerana* adult bees have been observed to rob *A. dorsata* colonies (Koeniger *et al.* 2002). Thus, foraging of different bee species in the same flower or robbing must be considered as a potential source of *T. clareae* infestation for uninhabited colonies.

Acknowledgements

We thank the Thailand Research Fund for the Royal Golden Jubilee PhD Program and National Science and Technology Development Agency of Thailand (grant number CO-B-07-22-09-005) for financial support. We also thank Associate Prof. Penrat Hongtayakorn and Surachai Leephitakrat for their assistance on research techniques and sample collection. This research was completed in co-operation with the Louisiana Agricultural Experiment Station.

REFERENCES

- AGGARWAL, K (1988). Incidence of *Tropilaelaps clareae* on three *Apis* species in Hisar (India). In Needham, G R; Page, R E; Delfinado-Baker, M; Bowman, C E (eds) *Africanized honey bees and bee mites*. Ellis Horwood, Chichester, UK; pp. 396–403.
- KOENIGER, N; KOENIGER, G (1980) Observations and experiments on migration and dance communication of *Apis dorsata* in Sri Lanka. *Journal of Apicultural Research* 19: 21–34.
- KOENIGER, G; KOENIGER, N; ANDERSON, D L; LEKPRAVON, C; TINGEK, S (2002) Mites from debris and sealed brood cells of *Apis dorsata* colonies in Sabah (Borneo) Malaysia, including a new haplotype of *Varroa jacobsoni*. *Apidologie* 33: 15–24.
- OLDROYD, B; RINDERER, T E; WONGSIRI, S (1992). Pollen resource partitioning by *Apis dorsata*, *A. cerana*, *A. andreniformis* and *A. florea* in Thailand. *Journal of Apicultural Research* 31: 3–7.
- RINDERER, T E; OLDROYD, P B; LEKPRAVON, C; WONGSIRI, S; BOONTHAI, C; THAPA, R (1994) Extended survival of the parasitic honey bee mite *Tropilaelaps clareae* on adult workers of *A. mellifera* and *A. dorsata*. *Journal of Apicultural Research* 33: 171–174.

Biography

Mr. Boonmee Kavinseksan was born on the 20th of July 1968. He graduated his Bachelor's Degree of Science in Agriculture (second class honors) in Agronomy major, from King Mongkut's Institute of Technology Chaohunthaharn Ladkrabang in 1991. In 1994, he received his Master's Degree of Science in Biotechnology program from Chulalongkorn University. His Ph.D. study in Biological Science program at the Faculty of Science, Chulalongkorn University was funded by the Royal Golden Jubilee Project of the Thailand Research Fund in 1999. At present, he is a lecturer at the Department of Biology, Faculty of Science and Technology, Bansomdejchaopraya Rajabhat University, Bangkok, Thailand.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย