

## BIBLIOGRAPHY

1. Beneke, E.S., and A.L. Rogers. 1971. Medical Mycology Manual. 3<sup>rd</sup> ed. Minneapolis Minn: Burgess Publishing Comp. 165 pp.
2. Beneke, E.S., S. Pichyangkura, and A.L. Rogers. 1975. Intracellular Soluble Protein, Enzyme and Isoenzyme. Patterns of some Dematiaceous Fungi. 75<sup>th</sup> International Society for Human and Animal Mycology Tokyo, Japan G(7)-5.
3. Benham, R.W., 1931. Certain Monilias Parasitic on Man. Their Identification by Morphology and by Agglutination. J. Infect. Dis. 49:183-215.
4. Berchev, K., and I. Izimirov. 1967. Isoenzyme of some Oxido-reductases in the Candida as a Basis of Species Identification after Electrophoresis. Experientia 23:961-962.
5. Bernfeld, P. 1951. Enzyme of Starch Degradation and Synthesis. Adv. in Enzymol. 12:380-428.
6. Brewbaker, J.L., M.O. Upadhy. 1968. Isoenzyme Polymorphism in Flowering Plants: II Gel. Electrophoresis and Applications. Physiol. Plant. 21:930-940.

7. Chang, L.O., A.M. Srb and F.C. Steward. 1962.  
Electrophoretic Separations of the Soluble  
Proteins of *Neurospora*. Nature. 193:756-759.
8. Chilgren, R.A., R. Hong, et. al. 1968. Human Serum  
Interactions with *Candida albicans*. J. Immunol.  
101:128-132.
9. Conant, N.F., et. al. 1971. Manual of Clinical  
Mycology. 3<sup>rd</sup> ed. Philadelphia: W.B. Saunders  
Comp. 325-364.
10. Dabrowa, N., D.H. Howard, J.W. Landau and Y. Shechter.  
1970. Synthesis of Nucleic Acids and Proteins  
in the Dimorphic Forms of *Candida albicans*.  
Sabouraudia 8:163-169.
11. Davis, B.J. 1964. Disc Electrophoresis II. Method  
and Application to Human Serum Proteins.  
Ann. New York Acad. Sci. 121:404-427.
12. Emmons, C.W. 1970. Medical Mycology. 2<sup>nd</sup> ed. Philadel-  
phia: Lea and Fibiger. 167-182.
13. Fairbanks G., T.L. Steck and D.F.H. Wallach. 1971.  
Electrophoretic Analysis of the Major Polypeptides  
of the Human Erythrocyte Membrane. Biochemistry.  
10 (13) : 2606-2616.

14. Franke, R.G. and J.A. Berry. 1972. Taxonomic Application of Isoenzymes Patterns Produced with Disc Electrophoresis of some Myxomycetes in the order Physarales. Mycologia 64:830-840.
15. Funder, S. 1968. Practical Mycology. New York: Hafner Publishing Comp., INC.
16. Gomori, G. 1955. Preparation of Buffers for use in Enzyme Studies. Vol. 1:138-148. Colowick, S.P. and Kaplan, N.O. Ed. in Methods in Enzymology. N.Y. Academic Press.
17. Gordon, M.A., E.G. Bradley and V.Q. Grant. 1952. The Influence of Different Types of Corn Meal Agar upon Chlamydospores Production by Candida albicans. J. Lab. & Clin. Med. 40:316-320.
18. Gordon, M.A., J.C. Elliott et. al. 1967. Identification of Candida albicans, other Candida species and Torulopsis glabrata by means of Immunofluorescence. Sabouraudia 5:323-328.
19. Gunsalus, C.I. 1955. Extraction of Enzymes from Microorganism (Bacteria and Yeast). Method of Enzymology. Academic Press. New York: Vol. I. pp. 51-64.
20. Hasenclever, H.F. 1959. Comparative Pathogenicity of Candida albicans for Mice and Rabbits. J.Bacteriol. 78:105-109.

21. Hasenclever, H.F. and W.O. Mitchell. 1961. Antigenic Studies of Candida I. Observation of two Antigenic Groups in Candida albicans. J. Bacteriol. 82:570-573.
22. Hasenclever, H.F. and W.O. Mitchell. 1961. Antigenic Studies of Candida II Antigenic Relation of C. albicans Group A and Group B to C. stellatoidea and C. tropicalis. J. Bacteriol. 82:574-577.
23. Hasenclever, H.F. 1963-64. Antigenic Studies of Candida IV. The Relationship of the Antigenic Groups of Candida albicans to their isolation from various clinical specimens. Sabouraudia 2:201-204.
24. Hasenclever, H.F. and W.O. Mitchell. 1964. Immunochemical Studies on Polysaccharides of Yeasts. J. of Immu. 93:763-771
25. Hurley, R. and V.C. Stanley. 1969. Cytopathic Effects of Pathogenic and Nonpathogenic Species of Candida on Culture Mouse Epithelial Cells: Relation to Growth Rate and Morphology of the Fungi. J. Med. Microbiol. 2:63-74.
26. Kapica, L. and A. Clifford. 1968. Quantitative Determination of Yeasts in Sputum. Mycopath. et. Mycol. Appl. 34:27-32.

27. Kligman, A.M. 1950. Aids in Technique in the Identification of Candida albicans. J. Invest. Dermatol. 14: 173-176.
28. Landau, J.W., Y. Shechter and V.D. Newcomer. 1968. Biochemical Taxonomy of the Dermatophyte II. Numerical Analysis of Electrophoretic Protein Patterns. J. Invest. Dermatol. 51:170-176.
29. Lodder, J. and N.J.W. Kreger-Van Rij. 1952. The Yeasts, A Taxonomic Study. Amsterdam, North Holland Publishing Comp. 713 p.
30. Louria, D.B. and R.G. Brayton. 1964. The Behavior of Candida Cells within Leucocytes. Proc. Soc. Exp. Biol. Med. 115:93-98.
31. Lowry, O.H., N.J. Rosebrough, A.L. Farr and R.J. Randall. 1951. Protein Measurement with the Folin Phenol Reagent. J. Biol. Chem. 193:265-275.
32. Mahgoub, E.S. 1968. The Incidence of Yeasts in Cases of Vaginitis and Children with Oral Thrush in the Sudan. Mycopath. et. Mycol. Appl. 36:145-149.
33. Maibach, H.I. and A.M. Kligman. 1962. The Biology of Experimental Human Cutaneous Moniliasis. Arch. Dermatol. 85:233-257.

34. Montes, L. and W.H. Wilborn. 1968. Ultrastructural Features of Host-parasite Relationship in Oral Candidiasis. J. Bacteriol. 96:1349-1356.
35. Nickerson, W.J. and Z.T. Mankowski. 1953. A Polysaccharide Medium of Known Composition Favoring Chlamydospore Formation in Candida albicans. J. Infect. Dis. 92 : 20-25.
36. Rippon, J.W. 1974. Medical Mycology. The Pathogenic Fungi and The Pathogenic Actinomycetes. Philadelphia. W.B. Saunders Comp! 175-204.
37. Shechter, Y., J.W. Landau, N. Dabrowa and V.D. Newcomer. 1966. Comparative Disc Electrophoretic Studies of Proteins from Dermatophytes. Sabouraudia 3: 144-149.
38. Shechter, Y., J.W. Landau and V.D. Newcomer. 1967. Disc Electrophoretic Studies of Proteins from Coccidioides immitis. J. Invest. Dermatol. 48:119-123.
39. Shechter, Y., J.W. Landau, N. Dabrows and V.D. Newcomer. 1968. Biochemical Taxonomy of the Dermatophytes I. Comparative Disc Electrophoresis of Culture Filtrate Proteins. J. Invest. Dermatol 51:165-169.

40. Shechter, Y., J.W. Landau, N. Dabrowa and V.D. Newcomer. 1968. Disc Electrophoretic Studies of Intra-specific Variability of Proteins from Dermato-phytes. Sabouraudia 6:133-137.
41. Shechter, Y., J.W. Landau and N. Dabrowa. 1972. Comparative Electrophoresis and Numerical Taxonomy of some Candida species. Mycologia 64:841-853.
42. Simonetti, N. and V. Strippoli. 1971. A New Simple Method for Rapid Acquisition of Chlamyospores in Candida albicans. Experientia 27/8 : 985.
43. Soles, P., L.Y. Lim et. al. 1967. Active Immunity in Experimental Candidiasis in Mice. Sabouraudia 5:315-322.
44. Sukroongreung, S. 1971. A Glutinous Rice Culture Medium for Demonstration of Chlamyospores of Candida albicans. Mycopath. et. Mycol. Appl. 43:329-335.
45. Taschdjian, C.L., J.J. Burchall and P.J. Kozinn. 1960. Rapid Identification of Candida albicans by Filamentation on Serum and Serum Substitutes. A.M.A.J. Dis. Child. 99:212-215.
46. Taschdjian, C.L., M.S. Manuel, et.al. 1969. A Modified Antigen for Serodiagnosis of Systemic Candidiasis. The Amer. J. of Clin. Pathol. 52:468-471.

47. Taschdjian, C.L., P.J. Kozinn, et.al. 1969. Post Mortem Studies of Systemic Candidiasis Sabouraudia 7:110-117.
48. Taylor, R.L., et. al. 1968. Occurrence of Dermatophytes in Bangkok, Thailand. Sabouraudia 6: 307-311.
49. Tsuchiya, T., M. Miyasaki, et. al. 1955. Studies on the Classification of the Genus Candida Comparison of Antigenic Structures of Standard and other Strains. Jap. J. Exp. Med. 25:15-21.



APPENDIX

Mounting medium, 10-20% KOH, was used in direct examination of infected tissue under microscope.

Mycosel agar (Sabouraud plus antibacterial and antifungal) was used in isolation the pathogenic fungi from clinical materials.

Mycosel agar formula:-

Bactopeptone	10	gm.
Dextrose	40	gm.
Agar	15	gm.
Chloramphenicol	0.05	gm.
Cycloheximide	0.5	gm.
Distilled water	1000	ml.

Sabouraud Dextrose Agar was used in order to maintain and cultivation of C. albicans, throughout this experiment.

Sabouraud dextrose formular -

Bactopeptone	10	gm.
Dextrose	40	gm.
Agar	15	gm.
Distilled water	1000	ml.

Corn meal agar (Benham, 1931) and Rice agar (Beneke, 1971) were used to study chlamydospore production in Candida albicans.

Horse serum and physiological saline (0.85% NaCl) were used on the purpose of studying germ tube (Taschdjian, 1960).