

การคัดแยกและลักษณะสมบัติของแบคทีเรียที่ทนต่อตัวทำละลายอินทรีย์

นางสาว อจิรากรณ์ คงผล

วิทยานิพนธ์นี้เป็นส่วนหนึ่งของการศึกษาตามหลักสูตรปริญญาวิทยาศาสตรมหาบัณฑิต
สาขาวิชาเทคโนโลยีชีวภาพ
คณะวิทยาศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย
ปีการศึกษา 2550
ลิขสิทธิ์ของจุฬาลงกรณ์มหาวิทยาลัย



**ISOLATION AND CHARACTERIZATION OF ORGANIC-SOLVENT
TOLERANT BACTERIA**

Miss Ajiraporn Kongpol

**A Thesis Submitted in Partial Fulfillment of the Requirements
for the Degree of Master of Science Program in Biotechnology**

Faculty of Science

Chulalongkorn University

Academic year 2007

Copyright of Chulalongkorn University

502103

อภิการณ์ คงผล : การคัดแยกและลักษณะสมบัติของแบคทีเรียที่ทนต่อตัวทำละลายอินทรีย์。
 (ISOLATION AND CHARACTERIZATION OF ORGANIC-SOLVENT TOLERANT
 BACTERIA) อ.ที่ปรึกษา : ผศ. ดร. อลิสา วงศ์วัน, 150 หน้า.

การคัดแยกแบคทีเรียที่อุณหภูมิ 45 องศาเซลเซียส จากแหล่งดิน 13 แหล่ง สามารถคัดแยก
 แบคทีเรียได้ 36 สายพันธุ์ ซึ่งมีความสามารถในการทนต่อตัวทำละลายอินทรีย์ได้แตกต่างกัน โดย
 แบ่งออกได้เป็น 3 กลุ่ม คือ 1) กลุ่มที่ทนต่อตัวทำละลายอินทรีย์ที่มีค่า $\log P_{ow}$ ในช่วงกว้าง ได้แก่ *Deinococcus geothermalis* T27 และ *Bacillus cereus* 4/1 โดยเฉพาะ *D. geothermalis* T27
 สามารถทนต่อเอทิลอะซิตेटได้ดีกว่าแบคทีเรียนิดอื่นในความเข้มข้น 20% (v/v) และพบว่าเมื่อ¹
 เลี้ยงภายใต้ภาวะที่มีเอทิลอะซิตेटจะมีผลทำให้เซลล์มีขนาดเล็กลง แต่ไม่พบการเปลี่ยนแปลงของ
 ชนิดและปริมาณกรดไขมันในเยื่อหุ้มเซลล์ ในขณะเดียวกัน *D. geothermalis* T27 ยังสามารถใช้
 เอทิลอะซิตेटเป็นแหล่งคาร์บอนและพลังงานได้ โดยตรวจพบว่ามีการเหนี่ยวนำให้เซลล์ผลิต
 เอสเทอเรสขึ้น ในขณะที่แบคทีเรีย *B. cereus* 4/1 สามารถทนต่อโกลูอีนซึ่งเป็นตัวทำละลายอินทรีย์
 ที่มีความเป็นพิษสูงได้ที่ความเข้มข้นสูงถึง 20% (v/v) และยังสามารถใช้โกลูอีนเป็นแหล่งคาร์บอน
 และพลังงานได้ 2) กลุ่มที่สามารถทนตัวทำละลายอินทรีย์ที่มีค่า $\log P_{ow}$ สูงกว่า 3 คือ *Bacillus subtilis* 45 เมื่อเลี้ยงเซลล์ในภาวะที่มีอีนเดคเคนไม่พบการเปลี่ยนแปลงขนาดของเซลล์และชนิด
 ของกรดไขมันในเยื่อหุ้มเซลล์ แบคทีเรียสายพันธุ์นี้สามารถทนต่ออีนเดคเคน และ ไซโคลเอกเซน
 ได้ในความเข้มข้นสูงถึง 20% (v/v) และยังใช้เป็นแหล่งคาร์บอนและพลังงานได้อีกด้วย 3) กลุ่มที่ทน
 ตัวทำละลายอินทรีย์ที่มีค่า $\log P_{ow}$ ต่ำกว่า 2 เช่น บิวทิลอะซิตेट บิวทานอล และ เอทิลอะซิตेट
 ได้แก่ *Brevibacillus agri* 13 เมื่อเลี้ยงแบคทีเรียสายพันธุ์นี้ในภาวะที่มีเอทิลอะซิตेट พบร่วงขนาด
 ของเซลล์ไม่มีการเปลี่ยนแปลงแต่มีการเพิ่มขึ้นของกรดไขมันในเยื่อหุ้มเซลล์ แบคทีเรียสายพันธุ์นี้
 สามารถใช้บิวทานอลเป็นแหล่งคาร์บอนและพลังงาน พร้อมทั้งตรวจพบแอลกอฮอล์ดีไฮดรอเจนส์
 เมื่อเลี้ยงในภาวะที่มีบิวทานอล นอกจากนี้ยังสามารถทนต่อ เอทิลอะซิตेट ได้ที่ความเข้มข้น 20%
 (v/v) แต่ไม่สามารถใช้เป็นแหล่งคาร์บอนหรือพลังงานได้ ลักษณะสมบัติของแบคทีเรียนต่อตัวทำ
 ละลายอินทรีย์เหล่านี้ แสดงถึงศักยภาพในการนำไปใช้ในกระบวนการผลิตสารผลิตภัณฑ์ทาง
 เทคโนโลยีชีวภาพ

สาขาวิชา เทคโนโลยีชีวภาพ..... ลายมือชื่อนักศึกษา..... ๑๗๖๘

ปีการศึกษา 2550 ลายมือชื่ออาจารย์ที่ปรึกษา..... 

4772550423: MAJOR BIOTECHNOLOGY

KEY WORD: ORGANIC-SOLVENT TOLERANT BACTERIA/ SCREENING/ ISOLATION

AJIRAPORN KONGPOL: ISOLATION AND CHARACTERIZATION OF ORGANIC-SOLVENT TOLERANT BACTERIA. THESIS ADVISOR: ASST. PROF. ALISA VANGNAI, Ph.D., 150 pp.

Thirty six organic solvent tolerant bacteria were isolated at 45°C from 13 soil samples. The bacterial isolates could be divided into 3 groups. 1) Bacteria which tolerate to organic solvents with a broad rang of $\log P_{ow}$ value consist of *Deinococcus geothermalis* T27 and *Bacillus cereus* 4/1. *D. geothermalis* T27 demonstrated high tolerance to 20% (v/v) ethyl acetate. *D. geothermalis* T27 cells in a direct contact to ethyl acetate appeared to have a smaller size but no significant change in the level of fatty acid composition. Cells could also degrade ethyl acetate and induce esterase when exposed to ethyl acetate. For *B. cereus* 4/1, it had high tolerance to 20% (v/v) toluene and could degrade toluene. 2) Bacteria which tolerate to organic solvents with $\log P_{ow} \geq 3.0$ is *Bacillus subtilis* 45. *B. subtilis* 45 cells changed neither cell size nor the concentration of fatty acid composition in *n*-decane exposure. *B. subtilis* 45 tolerated to 20% (v/v) of *n*-decane and cyclohexane and could degrade both of them. 3) Bacteria which tolerate to organic solvents with $\log P_{ow} \leq 2.0$ is *Brevibacillus agri* 13. Cells of *Brevibacillus agri* 13 did not change cell size after ethyl acetate exposure. However, there is a change in a fatty acid composition. It could degrade *n*-butanol and induced alcohol dehydrogenase when exposed to with *n*-butanol. *Brevibacillus agri* 13 could tolerate to 20% (v/v) ethyl acetate but could not degrade ethyl acetate. Intriguing characteristics of organic solvent tolerant bacteria may demonstrate their potential use in biotechnological processes.

Field of Study: Biotechnology.....Student's Signature: ...Ajiraporn Kongpol...

Academic Year: 2007.....Advisor's Signature: ...Alisa Vangnai...

Thesis Title ISOLATION AND CHARACTERIZATION OF ORGANIC-SOLVENT TOLERANT BACTERIA
By Miss Ajiraporn Kongpol
Filed of study Biotechnology
Thesis Advisor Assistant Professor Alisa Vangnai, Ph.D.

Accepted by the Faculty of Science, Chulalongkorn University in Partial Fulfillment of the Requirements for the Master's Degree


..... Dean of the Faculty of Science
(Professor Supot Hannongbua, Ph.D.)

THESIS COMMITTEE


..... Chairman
(Professor Aran Incharoensakdi, Ph.D.)


..... Thesis Advisor
(Assistant Professor Alisa Vangnai, Ph.D.)


..... Member
(Associate Professor Warawut Chulalaksananukul, Ph.D.)


..... Member
(Assistant Professor Kanoktip Packdibamrung, Ph.D.)

ACKNOWLEDGEMENTS

I would like to express my deepest gratitude to my advisor, Assistant Professor Alisa Vangnai for her excellent instruction, guidance, encouragement, attention and support throughout this thesis. Without her kindness and understanding, this work could not be accomplished.

My gratitude is also extended to Professor Aran Incharoensakdi, Associate Professor Warawut Chulaluksananukul and Assistant Professor Kanoktip Packdibamrung for serving as thesis committee, for valuable comments and also for useful suggestions.

There are so many friends to mention from so little space to fill. I would like to give a deep thank to all staff members of room 604 and friends in the Department of Biochemistry and Program in Biotechnology for their friendship, kind assistance and support. I would like to express thanks to Miss Kanokwan Chodchey and Professor Cornelis Verduyn for FAME analysis.

Finally, the greatest gratitude is expressed to my lovely family for their infinite love, willpower, support, understanding, encouragement and everything giving to my life.

This work was financially supported by 90th anniversary of Chulalongkorn University Fund, Graduate School, and Program in Biotechnology, Faculty of Science, Chulalongkorn University.

CONTENTS

	Page
ABSTRACT (THAI).....	iv
ABSTRACT (ENGLISH).....	v
ACKNOWLEDGEMENTS.....	vi
CONTENTS.....	vii
LIST OF TABLES.....	xiv
LIST OF FIGURES.....	xvi
NOMENCLATURE.....	xix
CHAPTER I INTRODUCTION.....	1
1.1 Statement of problem	1
1.2 Objectives and expected results.....	3
1.3 Hypotheses.....	4
1.4 Scopes of the study.....	4
1.5 The organization.....	5
CHAPTER II LITERATURE REVIEW.....	6
2.1 Organic solvent.....	6
2.2 Use of organic solvent.....	7
2.3 Organic solvent tolerant bacteria.....	8

	Page
2.4 Physiological basis of organic solvent toxicity	
and the concept of organic solvent tolerance	9
2.5 Isolation of organic solvent tolerant bacteria	10
2.6 Adaptive mechanisms of organic-solvent tolerance	14
2.6.1 Morphological adaptation	15
2.6.2 Changes of the energetic status	15
2.6.3 Adaptation of the cell membrane	16
2.6.4 Changes in the cell wall and outer membrane.....	17
2.6.5 Modifying surface properties (charge and hydrophobicity)	17
2.6.6 Transformation or degradation of the organic solvent.....	18
2.6.7 Active transport of solvents from the membrane into the environment by energy-consuming efflux systems	18
2.6.8 Modifying membrane proteins.....	19
2.7 Application of organic solvent-tolerant bacteria.....	19
2.7.1 Biotechnology application	20
2.7.2 Bioremediation	21

	Page
CHAPTER III METHODOLOGY.....	23
3.1 Laboratory equipments, chemicals	23
3.1.1 Laboratory equipments.....	23
3.1.2 Laboratory chemicals.....	24
3.2 Culture medium.....	27
3.3 Methodologies	29
3.3.1 Screening and isolation of organic solvent tolerant bacteria using enrichment technique with toluene and cyclohexane vapor.....	30
3.3.2 Primary test of the isolates for solvent utilization and tolerance	30
3.3.3 Secondary test of the isolates for solvent utilization and tolerance	31
3.3.4 Identification of the organic solvent-tolerant bacteria	31
3.3.5 Characterization of the organic solvent-tolerant bacteria.....	33
3.3.5.1 Effect of types and concentrations of organic solvent on cell growth and tolerance	33
3.3.5.2 Effect of organic solvent on cell morphology ...	34
3.3.5.3 Determination of cell fatty acid composition	36
3.3.5.4 Organic solvent utilization using resting cell technique	37

	Page
3.3.5.5 Determination of enzymatic activity involving organic solvent utilization	38
3.3.6 Factors involving organic solvent tolerance of the bacteria	42
3.3.6.1 Effect of divalent ion on cell growth and organic solvent tolerant	42
3.3.6.2 Effect of nutrient on cell growth and organic solvent tolerance	42
CHAPTER IV RESULTS	43
4.1 Screening, isolation and identification of organic solvent-tolerant bacteria	44
4.1.1 Primary test of the isolates for organic solvent utilization and tolerance.....	44
4.1.2 Secondary test of the isolates for organic solvent utilization and tolerance	45
4.1.3 Identification of organic solvent-tolerant bacteria	59
4.1.3.1 The morphological characteristics	59
4.1.3.2 The biochemical test	59
4.1.3.3 16S ribosomal DNA gene	59

	Page
4.3.2.2 Effect of nutrient on growth and tolerance	87
4.4 <i>Bacillus subtilis</i> strain 45	89
4.4.1 Characterization of organic-solvent tolerant bacteria	89
4.4.1.1 Effect of type and concentration of organic solvent on growth and tolerance.....	89
4.4.1.2 Effect of organic solvent on cell morphology	91
4.4.1.3 Effect of organic solvent on fatty acid composition	92
4.4.1.4 Organic solvent utilization	93
4.4.2 Factors involving of organic-solvent tolerance	94
4.4.2.1 Effect of ions on growth and tolerance	94
4.4.2.2 Effect of nutrient on growth and tolerance	98
4.5 <i>Brevibacillus agri</i> strain 13	101
4.5.1 Characterization of organic-solvent tolerant bacteria ...	101
4.5.1.1 Effect of type and concentration of organic solvent on growth and tolerance	101
4.5.1.2 Effect of organic solvent on cell morphology ...	102
4.5.1.3 Effect of organic solvent on fatty acid composition	104
4.5.1.4 Organic solvent utilization	104
4.5.1.5 Enzymatic activity involving organic solvent utilization.....	106

	Page
4.5.2 Factors involving of organic-solvent tolerance	108
4.5.2.1 Effect of ions on growth and tolerance	108
4.5.2.2 Effect of nutrient on growth and tolerance.....	112
CHAPTER V DISCUSSION.....	114
Discussion	114
CHAPTER VI CONCLUSIONS.....	124
Conclusions.....	124
REFERENCES.....	126
APPENDICES.....	135
Appendix A.....	135
Appendix B	146
Appendix C	148
BIOGRAPHY.....	149

LIST OF TABLES

Table	Page
2.1 Chemical classification of organic solvents	6
2.2 Popular solvents in solvent-containing products	8
2.3 Organic solvents and their log P_{ow} values	10
2.4 Examples of organic solvent tolerant bacteria	12
2.5 Solvent tolerance of Gram-negative bacteria	13
2.6 Solvent tolerance of Gram-positive bacteria	13
4.1 Source of soil sample used for screening and bacterial isolates obtained ...	46
4.2 Primary test for solvent tolerance and solvent utilization on agar overlaid with solvent	47
4.3 Organic solvent tolerant bacterial isolates chosen for further study	52
4.4 Morphological characteristic of bacteria	61
4.5 The biochemical test results of four bacterial isolates were chosen	63
4.6 Comparison of 16S ribosomal DNA gene for bacterial identification	64
4.7 Cell size of <i>D. geothermalis</i> T27 exposed and not exposed ethyl acetate ...	68
4.8 Fatty acid composition of <i>D. geothermalis</i> T27 non-exposed and exposed to ethyl acetate	70
4.9 Esterase activity of <i>D. geothermalis</i> T27	72
4.10 Cell size of <i>B. cereus</i> strain 4/1 exposed and not exposed to chloroform ...	81
4.11 Fatty acid composition of <i>B. cereus</i> strain 4/1 non-exposed and exposed to chloroform	82
4.12 Cell size of <i>B. subtilis</i> strain 45 grown with and without <i>n</i> -decane	91

	Page
4.13 Fatty acid composition of <i>B. subtilis</i> strain 45 non-exposed and exposed to <i>n</i> -decane	93
4.14 Cell size of <i>Brevibacillus agri</i> strain 13 exposed with ethyl acetate.....	103
4.15 Fatty acid composition of <i>Brevibacillus agri</i> strain 13 non-exposed and exposed to ethyl acetate	104
4.16 Esterase activity of <i>Brevibacillus agri</i> strain 13	106

LIST OF FIGURES

Figure	Page
2.1 Adaptive mechanisms protecting cells against toxic effects of organic solvents	14
3.1 A flow chart of experimental methods of the thesis	29
3.2 Characteristics of colonies	32
4.1 Secondary test of bacterial isolates in group I	53
4.2 Secondary tests of bacterial isolates in group II.....	55
4.3 Secondary tests of bacterial isolates in group IV	57
4.4 Growth inhibitions of <i>D. geothermalis</i> T27	67
4.5 Cell morphology of <i>D. geothermalis</i> T27	69
4.6 Utilization of ethyl acetate by <i>D. geothermalis</i> T27	71
4.7 Effect of divalent ion on toluene tolerance in <i>D. geothermalis</i> T27	74
4.8 Effect of divalent ion on ethyl acetate tolerance in <i>D. geothermalis</i> T27	75
4.9 Growth of <i>D. geothermalis</i> T27 in various types of nutrients	77
4.10 Effect of nutrient on cyclohexane; toluene; benzene; ethyl acetate tolerance in <i>D. geothermalis</i> T27	78
4.11 Growth inhibition of <i>B. cereus</i> strain 4/1	80
4.12 Cell morphology of <i>B. cereus</i> strain 4/1	81

	Page
4.13 Utilization of toluene by <i>B. cereus</i> strain 4/1	83
4.14 Effect of divalent ion on styrene tolerance in <i>B. cereus</i> strain 4/1	85
4.15 Effect of divalent ion on toluene tolerance in <i>B. cereus</i> strain 4/1	86
4.16 Growth of <i>B. cereus</i> strain 4/1 in various types of nutrients.....	87
4.17 Effect of nutrient on styrene; toluene; chloroform tolerance in <i>B. cereus</i> strain 4/1	88
4.18 Growth inhibition of <i>B. subtilis</i> strain 45	90
4.19 Cell morphology of <i>B. subtilis</i> strain 45	92
4.20 Utilization of <i>n</i> -decane by <i>B. subtilis</i> strain 45	93
4.21 Utilization of cyclohexane by <i>B. subtilis</i> strain 45	94
4.22 Effect of divalent ion on <i>n</i> -heptane tolerance in <i>B. subtilis</i> strain 45	95
4.23 Effect of divalent ion on cyclohexane tolerance in <i>B. subtilis</i> strain 45	96
4.24 Effect of divalent ion on ethylbenzene tolerance in <i>B. subtilis</i> strain 45	97
4.25 Growth of <i>B. subtilis</i> strain 45 in varies type of nutrients	98
4.26 Effect of nutrient on <i>n</i> -decane; <i>n</i> -heptane; <i>n</i> -hexane; cyclohexane; ethylbenzene tolerance in <i>B. subtilis</i> strain 45	99
4.27 Growth inhibition of <i>Brevibacillus agri</i> strain 13	102
4.28 Cell morphologies of <i>Brevibacillus agri</i> strain 13	103

	Page
4.29 Utilization of <i>n</i> -butanol by <i>Brevibacillus agri</i> strain 13	105
4.30 Utilization of ethyl acetate by <i>Brevibacillus agri</i> strain 13	105
4.31 Native- polyacrylamind gel electrophorsis with activity staining of butanol dehydrogenase	107
4.32 Effect of divalent ion on butyl acetate tolerance in <i>Brevibacillus agri</i> strain 13	109
4.33 Effect of divalent ion on <i>n</i> -butanol tolerance in <i>Brevibacillus agri</i> strain 13	110
4.34 Effect of divalent ion on ethyl acetate tolerance in <i>Brevibacillus agri</i> strain 13	111
4.35 Growth of <i>Brevibacillus agri</i> strain 13 in various types of nutrients. Symbols of organic solvents	112
4.36 Effect of nutrient on butyl acetate; <i>n</i> -butanol; ethyl acetate tolerance in <i>Brevibacillus agri</i> strain 13	113

NOMENCLATURE

CFU	=	Colony forming unit
ml	=	Milliliter
µl	=	Microliter
µg	=	microgram
O.D.	=	Optical density
rpm	=	Revolution per minut
min	=	Minute
°C	=	Degree of Celsius
cm	=	centimeter
<i>et al.</i>	=	ET. Alii (latin), and others
g	=	Gravitation acceleration
h	=	hour
l	=	liter
M	=	molar
mM	=	millimolar
mg	=	milligram
min	=	minute
MW	=	molecular weight