

## REFERENCES

- [1] Zhorov, B. S.; Bregestovski, P. D. Chloride channels of glycine and GABA receptors with blockers: Monte Carlo minimization and structure-activity relationships. *Biophys. J.* **2000**, *78*, 1786-1803.
- [2] Livingston, S. Antiepileptic drugs. *Am. J. Nurs.* **1963**, *63*, 103-107.
- [3] Fiedler, D.; Leung, D. H.; Bergman, R. G.; Raymond, K. N. Selective molecular recognition, C-H bond activation, and catalysis in nanoscale reaction vessels. *Acc. Chem. Res.* **2005**, *38*, 349-358.
- [4] Lehn, J. M. *Supramolecular Chemistry*; VCH: Weinheim, 1995.
- [5] Trapani, G.; Laquintana, V.; Latrofa, A.; Ma, J.; Reed, K.; Serra, M.; Biggio, G.; Liso, G.; Gallo, J. M. Peripheral benzodiazepine receptor ligand-melphalan conjugates for potential selective drug delivery to brain tumors. *Bioconjugate Chem.* **2003**, *14*, 830-839.
- [6] Cram, D. J.; Cram, J. M. Host-guest chemistry: Complexes between organic compounds simulate the substrate selectivity of enzymes. *Science* **1974**, *183*, 803-809.
- [7] Fitzmaurice, R. J.; Kyne, G. M.; Douheret, D.; Kilburn, J. D. Synthetic receptors for carboxylic acids and carboxylates. *J. Chem. Soc., Perkin Trans. 1* **2002**, *7*, 841-846.
- [8] Linton, B.; Hamilton, A. D. Formation of artificial receptors by metal-templated self-assembly. *Chem. Rev.* **1997**, *97*, 1669-1680.
- [9] Schmidtchen, F. P.; Berger, M. Artificial organic host molecules for anions. *Chem. Rev.* **1997**, *97*, 1609-1646.
- [10] Martinez-Manez, R.; Sancenon, F. Fluorogenic chromogenic chemosensors and reagents for anions. *Chem. Rev.* **2003**, *103*, 4419-4476.
- [11] Beer, P. D. Redox responsive macrocyclic receptor molecules containing transition metal redox centers. *Chem. Soc. Rev.* **1989**, *18*, 409-450.
- [12] Suksai, C.; Tuntulani, T. Chromogenic anion sensors. *Chem. Soc. Rev.* **2003**, *32*, 192-202.
- [13] Suksai, C.; Tuntulani, T. Chromogenic anion sensors. *Top. Curr. Chem.* **2005**, *255*, 163-198.
- [14] Lee, D. H.; Lee, H. Y.; Hong, J.-I. Anion sensor based on the indoaniline-thiourea system. *Tetrahedron Lett.* **2002**, *43*, 7273-7276.


- [15] Beer, P. D.; Hayes, E. J. Transition metal and organometallic anion complexation agents. *Coord. Chem. Rev.* **2003**, *240*, 167-189.
- [16] Fages, F.; Desvergne, J.-P.; Bouas-Laurent, H. Synthesis, structural, spectroscopic, and alkali-metal cations complexation studies of a bis-anthracenediyl macrotricyclic ditopic receptor. *J. Org. Chem.* **1994**, *59*, 5264-5271.
- [17] Prez-Inestrosa, E. Desvergne, J.-P.; Bouas-Laurent, H.; Rayez, J.-C.; Rayez, M.-T.; Cotrait, M.; Marsau, P. Barium cation complexation by flexible mono- and ditopic receptors, studied by UV absorption and fluorescence spectroscopy. *Eur. J. Org. Chem.* **2002**, 331-334.
- [18] Arduini, A.; Brindani, E.; Giorgi, G.; Pochini, A.; Secchi, A. Recognition of guests bearing donor and receptor hydrogen bonding groups by heteroditopic calix[4]arene receptors. *Tetrahedron* **2003**, *59*, 7587-7594.
- [19] Pelizzi, N.; Casnati, A.; Friggeri, A.; Ungaro, R. Synthesis and properties of new calixarene-based ditopic receptors for the simultaneous complexation of cations and carboxylate anions. *J. chem. Soc. Perkin Trans.* **1998**, *2*, 1307-1311.
- [20] Schreeder, J.; van Duynhover, J. P. M.; Engbersen, J. F. J.; Reinhoudt, D. N. Solubilization of NaX salts in chloroform by bifunctional receptors. *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 1090-1093.
- [21] Deetz, M.J.; Shang, M.; Smith, B. D. A macrobicyclic receptor with versatile recognition properties: simultaneous binding of an ion pair and selective complexation of dimethylsulfoxide. *J. Am. Chem. Soc.* **2000**, *122*, 6201-6207.
- [22] Mahoney, J. M.; Beatty, A. M.; Smith, B. D. Selective recognition of an alkali halide contact ion-pair. *J. Am. Chem. Soc.* **2001**, *123*, 5847-5848.
- [23] Cooper, J. B.; Drew, M. G. B.; Beer, P. D. Heteroditopic rhenium(I) and ruthenium(II) bipyridyl calix[4]arene receptors for binding cation-anion ion pairs. *J. Chem. Soc., Dalton Trans.* **2001**, 392-401.
- [24] Beer, P. D.; Drew, M. G. B.; Knubley, R. J.; Ogden, M. I. Synthesis and coordination chemistry of a novel bis(benzo crown ether) substituted calix[4]arene that can simultaneously complex cations and anions. *J. Chem. Soc., Dalton Trans.* **1995**, 3117-3124.

- [25] Miyaji, H.; Collinson, S. R.; Proke, I.; Tucker, J. H. R. A ditopic ferrocene receptor for anions and cations that functions as a chromogenic molecular switch. *Chem. Commun.* **2003**, 64-65.
- [26] Suksai, C.; Leeladee, P.; Jainuknan, D.; Tuntulani, T.; Muangsin, N.; Chailapakul, O.; Kongsaree, P.; Pakavatchai, C. A new heteroditopic receptor and sensor highly selective for bromide in the presence of a bound cation. *Tetrahedron Lett.* **2005**, 46, 2765-2769.
- [27] Knobloch, F. W.; Rauscher, W. H. Condensation polymers of ferrocene derivatives. *J. Polymer Sci.* **1961**, 54, 651-656.
- [28] Macomber, R. S. An introduction to NMR titration for studying rapid reversible complexation. *J. Chem. Edu.* **1992**, 69, 375-377.
- [29] Lockhart, J. C.; Robson, A. C.; Thompson, M. E.; Tyson, P. D.; Wallace, I. H. M. Ligands for the alkali metals. Part 4. Nuclear magnetic resonance of crown ethers with alkali metal ions. *J. Chem. Soc., Dalton Trans.* **1978**, 611-617.
- [30] Shriver, D. F.; Atkins, P. W. *Inorganic Chemistry*. English: Oxford University Press, 1994.
- [31] Hynes, M. J. EQNMR: a computer program for the calculation of stability constants from nuclear magnetic resonance chemical shift data. *J. Chem. Soc., Dalton Trans.* **1993**, 311-312.
- [32] Al-Sayah, M. H.; Branda, N. R. Protons as the triggers to regulate hydrogen-bonding receptors. *Org. Lett.* **2002**, 4, 881-884.
- [33] Tumcharern, G.; Tuntulani, T.; Coles, S. J.; Hursthouse, M. B.; Kilburn, J. D. A novel ditopic receptor and reversal of anion binding selectivity in the presence and absence of bound cation. *Org. Lett.* **2003**, 5, 4971-4974.
- [34] Beer, P. D.; Gale, P. A.; Chen, G. Z. Mechanisms of electrochemical recognition of cations, anions and neutral guest species by redox-active receptor molecules. *Coord. Chem. Rev.* **1999**, 185-186, 3-36.
- [35] Medina, J. C.; Goodnow, T. T.; Rojas, M. T.; Atwood, J. L.; Lynn, B. C.; Kaifer, A. E.; Gokel, G. W. Ferrocenyl iron as a donor group for complexed silver in ferrocenyldimethyl[2.2]cryptand: a redox-switched receptor effective in water. *J. Am. Chem. Soc.* **1992**, 114, 10583-10595.


- [36] Beer, P. D.; Gale, P. A.; Chen, G. Z. Electrochemical molecular recognition: pathways between complexation and signalling. *J. Chem. Soc., Dalton Trans.* **1999**, 1897-1909.
- [37] Saji, T.; Kinoshita, J. Electrochemical ion transport with ferrocene functionalized crown ether. *J. Chem. Soc., Chem. Commun.* **1986**, 716-717.
- [38] Beer, P. D. Anion selective recognition and optical/electrochemical sensing by novel transition-metal receptor systems. *Chem. Commun.* **1996**, 689-696.
- [39] Medina, J. C.; Goodnow, T. T.; Bott, S.; Atwood, J. L.; Kaifer, A. E.; Gokel, G. W. Ferrocenyldimethyl-[2.2]-cryptand: Solid state structure of the external hydrate and alkali and alkali-earth-dependent electrochemical behavior. *J. Chem. Soc., Chem. Commun.* **1991**, 290-292.
- [40] Miller, S. R.; Gustowski, D. A.; Chen, Z. H.; Gokel, G. W.; Echegoyen, L.; Kaifer, A. E. Rationalization of the unusual electrochemical behavior observed in lariat ethers and other reducible macrocyclic systems. *Anal. Chem.* **1988**, *60*, 2021-2024.
- [41] Scherer, M.; Sessler, J. L.; Gebauer, A.; Lynch, V. A bridged pyrrolic *ansa*-ferrocene. A new type of anion receptor. *Chem. Commun.* **1998**, 85-86.
- [42] Gale, P. A.; Hursthouse, M. B.; Light, M. E.; Sessler, J. L.; Warriner, C. N.; Zimmerman, R. S. Ferrocene-substituted calix[4]pyrrole: a new electrochemical sensors for anions involving CH—anion hydrogen bonds. *Tetrahedron Lett.* **2001**, *42*, 6759-6762.




## **APPENDIX**




**Figure A.1**  $^1\text{H-NMR}$  spectrum of compound **1a** in  $\text{CDCl}_3$



**Figure A.2**  $^{13}\text{C-NMR}$  spectrum of compound **1a** in  $\text{CDCl}_3$



**Figure A.3**  $^1\text{H-NMR}$  spectrum of compound **1b** in  $\text{CDCl}_3$



**Figure A.4**  $^{13}\text{C-NMR}$  spectrum of compound **1b** in  $\text{CDCl}_3$




Figure A.5  $^1\text{H-NMR}$  spectrum of compound **1** in  $\text{CDCl}_3$




Figure A.6  $^{13}\text{C-NMR}$  spectrum of compound **1** in  $\text{CDCl}_3$






Figure A.7  $^1\text{H}$ - $^1\text{H}$  COSY spectrum of compound **1** in  $\text{CDCl}_3$






Figure A.8  $^1\text{H}$ - $^{13}\text{C}$  HMQC spectrum of compound **1** in  $\text{CDCl}_3$



**Figure A.9**  $^1\text{H-NMR}$  spectrum of compound **2a** in  $\text{CDCl}_3$



**Figure A.10**  $^{13}\text{C-NMR}$  spectrum of compound **2a** in  $\text{CDCl}_3$




Figure A.11  $^1\text{H-NMR}$  spectrum of compound **2b** in  $\text{CDCl}_3$




Figure A.12  $^{13}\text{C-NMR}$  spectrum of compound **2b** in  $\text{CDCl}_3$




Figure A.13  $^1\text{H-NMR}$  spectrum of compound **2** in  $\text{CDCl}_3$




Figure A.14  $^{13}\text{C-NMR}$  spectrum of compound **2** in  $\text{CDCl}_3$




Figure A.15  $^1\text{H}$ - $^1\text{H}$  COSY spectrum of compound **2** in  $\text{CDCl}_3$






Figure A.16  $^1\text{H}$ - $^{13}\text{C}$  HMQC spectrum of compound **2** in  $\text{CDCl}_3$






**Figure A.17**  $^1\text{H-NMR}$  titration of receptor **1** with  $\text{NaPF}_6$  in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where ▼ is NH protons of the complex  $[\mathbf{1}.\text{Na}^+]$  and ◆ is NH proton of the free receptor **1**




**Figure A.18**  $^1\text{H-NMR}$  titration of receptor **1** with  $\text{KPF}_6$  in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where ▼ is NH protons of the complex  $[\mathbf{1}.\text{K}^+]$  and ◆ is NH proton of the free receptor **1**




**Figure A.19**  $^1\text{H-NMR}$  titration of receptor **2** with  $\text{NaPF}_6$  in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where ▼ is NH protons of the complex  $[\mathbf{2}.\text{Na}^+]$  and ◆ is NH proton of the free receptor **1**




**Figure A.20**  $^1\text{H-NMR}$  titration of receptor **2** with  $\text{KPF}_6$  in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where ▼ is NH protons of the complex  $[\mathbf{2}.\text{K}^+]$  and ◆ is NH proton of the free receptor **1**




**Figure A.21**  $^1\text{H-NMR}$  titration of receptor **1** with benzoate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the receptor **1**.




**Figure A.22**  $^1\text{H-NMR}$  titration of complex  $[1.\text{Na}^+]$  with benzoate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the complex  $[1.\text{Na}^+]$  and  $\blacktriangledown$  is NH protons of the free receptor **1**.




**Figure A.23**  $^1\text{H-NMR}$  titration of receptor **1** with acetate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the receptor **1**.



**Figure A.24**  $^1\text{H-NMR}$  titration of complex  $[\mathbf{1}.\text{Na}^+]$  with acetate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the complex  $[\mathbf{1}.\text{Na}^+]$  and  $\blacktriangledown$  is NH protons of the free receptor **1**.




**Figure A.25**  $^1\text{H-NMR}$  titration of receptor **1** with dihydrogenphosphate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the receptor **1**.




**Figure A.26**  $^1\text{H-NMR}$  titration of complex  $[\mathbf{1}.\text{Na}^+]$  with dihydrogenphosphate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the complex  $[\mathbf{1}.\text{Na}^+]$  and  $\blacktriangledown$  is NH protons of the free receptor **1**.







**Figure A.27**  $^1\text{H-NMR}$  titration of receptor **1** with chloride in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the receptor **1**.




**Figure A.28**  $^1\text{H-NMR}$  titration of complex  $[1.\text{Na}^+]$  with chloride in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the complex  $[1.\text{Na}^+]$  and  $\blacktriangledown$  is NH protons of the free receptor **1**.




**Figure A.29**  $^1\text{H-NMR}$  titration of receptor **1** with bromide in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the receptor **1**.




**Figure A.30**  $^1\text{H-NMR}$  titration of complex  $[\mathbf{1}.\text{Na}^+]$  with bromide in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the complex  $[\mathbf{1}.\text{Na}^+]$ .




**Figure A.31**  $^1\text{H-NMR}$  titration of receptor **1** with iodide in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where ◆ is NH protons of the receptor **1**.




**Figure A.32**  $^1\text{H-NMR}$  titration of complex  $[1.\text{Na}^+]$  with iodide in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where ◆ is NH protons of the complex  $[1.\text{Na}^+]$ .




**Figure A.33**  $^1\text{H-NMR}$  titration of receptor **2** with benzoate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the receptor **2**.



**Figure A.34**  $^1\text{H-NMR}$  titration of complex  $[2.\text{Na}^+]$  with benzoate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the complex  $[2.\text{Na}^+]$  and  $\blacktriangledown$  is NH protons of the free receptor **2**.




**Figure A.35**  $^1\text{H-NMR}$  titration of receptor **2** with acetate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the receptor **2**.




**Figure A.36**  $^1\text{H-NMR}$  titration of complex  $[2.\text{Na}^+]$  with acetate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the complex  $[2.\text{Na}^+]$  and  $\blacktriangledown$  is NH protons of the free receptor **2**.







**Figure A.37**  $^1\text{H-NMR}$  titration of receptor **2** with dihydrogenphosphate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the receptor **2**.




**Figure A.38**  $^1\text{H-NMR}$  titration of complex  $[2.\text{Na}^+]$  with dihydrogenphosphate in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the complex  $[2.\text{Na}^+]$  and  $\blacktriangledown$  is NH protons of the free receptor **2**.




**Figure A.39**  $^1\text{H-NMR}$  titration of receptor **2** with chloride in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the receptor **2**.




**Figure A.40**  $^1\text{H-NMR}$  titration of complex  $[2.\text{Na}^+]$  with chloride in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the complex  $[2.\text{Na}^+]$  and  $\blacktriangledown$  is NH protons of the free receptor **2**.




**Figure A.41**  $^1\text{H-NMR}$  titration of receptor **2** with bromide in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the receptor **2**.




**Figure A.42**  $^1\text{H-NMR}$  titration of complex  $[2.\text{Na}^+]$  with bromide in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where  $\blacklozenge$  is NH protons of the complex  $[2.\text{Na}^+]$ .




**Figure A.43**  $^1\text{H-NMR}$  titration of receptor **2** with iodide in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where ◆ is NH protons of the receptor **2**.



**Figure A.44**  $^1\text{H-NMR}$  titration of complex  $[2.\text{Na}^+]$  with iodide in 5%  $\text{CD}_3\text{CN}:\text{CDCl}_3$  where ◆ is NH protons of the complex  $[2.\text{Na}^+]$ .




**Figure A.45** CV titration of receptor **1** with acetate in 40%  $\text{CH}_3\text{CN}:\text{CH}_2\text{Cl}_2$  with 0.1M  $\text{TBAPF}_6$  at scan rate 100 mV/s.




**Figure A.46** CV titration of receptor **1** with chloride in 40%  $\text{CH}_3\text{CN}:\text{CH}_2\text{Cl}_2$  with 0.1M  $\text{TBAPF}_6$  at scan rate 100 mV/s.







**Figure A.47** CV titration of receptor **1** with bromide in 40%  $\text{CH}_3\text{CN}:\text{CH}_2\text{Cl}_2$  with 0.1M  $\text{TBAPF}_6$  at scan rate 100 mV/s.




**Figure A.48** CV titration of complex  $[1.\text{Na}^+]$  with acetate in 40%  $\text{CH}_3\text{CN}:\text{CH}_2\text{Cl}_2$  with 0.1M  $\text{TBAPF}_6$  at scan rate 100 mV/s.




**Figure A.49** CV titration of complex  $[1.Na^+]$  with chloride in 40%  $CH_3CN:CH_2Cl_2$  with 0.1M  $TBAPF_6$  at scan rate 100 mV/s.




**Figure A.50** CV titration of receptor **2** with benzoate in 40%  $CH_3CN:CH_2Cl_2$  with 0.1M  $TBAPF_6$  at scan rate 100 mV/s.




**Figure A.51** CV titration of receptor **2** with acetate in 40%  $\text{CH}_3\text{CN}:\text{CH}_2\text{Cl}_2$  with 0.1M  $\text{TBAPF}_6$  at scan rate 100 mV/s.




**Figure A.52** CV titration of receptor **2** with chloride in 40%  $\text{CH}_3\text{CN}:\text{CH}_2\text{Cl}_2$  with 0.1M  $\text{TBAPF}_6$  at scan rate 100 mV/s.



**Figure A.53** CV titration of receptor **2** with bromide in 40%  $\text{CH}_3\text{CN}:\text{CH}_2\text{Cl}_2$  with 0.1M  $\text{TBAPF}_6$  at scan rate 100 mV/s.



**Figure A.54** CV titration of complex  $[2.\text{Na}^+]$  with acetate in 40%  $\text{CH}_3\text{CN}:\text{CH}_2\text{Cl}_2$  with 0.1M  $\text{TBAPF}_6$  at scan rate 100 mV/s.



**Figure A.55** CV titration of complex [2.Na<sup>+</sup>] with chloride in 40% CH<sub>3</sub>CN:CH<sub>2</sub>Cl<sub>2</sub> with 0.1M TBAPF<sub>6</sub> at scan rate 100 mV/s.



## VITAE

- Name:** Miss Nisachol Nerngchamnong
- Born:** May 22<sup>nd</sup>, 1984
- Education:** 2005 B. Sc. (Chemistry), Chulalongkorn University, Bangkok, Thailand.  
2008 M. Sc. (Inorganic Chemistry), Chulalongkorn University, Bangkok, Thailand