

REFERENCES



- Aimes, R. T., and Quigley, J. P. (1995). Matrix metalloproteinase-2 is an interstitial collagenase. Inhibitor-free enzyme catalyzes the cleavage of collagen fibrils and soluble native type I collagen generating the specific 3/4- and 1/4-length fragments. J Biol Chem 270, 5872-6.
- Albrechtsen, R., Nielsen, M., Wewer, U., Engvall, E., and Ruoslahti, E. (1981). Basement membrane changes in breast cancer detected by immunohistochemical staining for laminin. Cancer Res 41, 5076-81.
- Allan, J. A., Docherty, A. J., Barker, P. J., Huskisson, N. S., Reynolds, J. J., and Murphy, G. (1995). Binding of gelatinases A and B to type-I collagen and other matrix components. Biochem J 309, 299-306.
- Arenas-Huertero, F. J., Herrera-Goepfert, R., Delgado-Chavez, R., Zinser-Sierra, J. W., De la Garza-Salazar, J. G., Herrera-Gomez, A., and Perez-Cardenas, E. (1999). Matrix metalloproteinases expressed in squamous cell carcinoma of the oral cavity: correlation with clinicopathologic features and neo-adjuvant chemotherapy response. J Exp Clin Cancer Res 18, 279-84.
- Atkinson, S. J., Crabbe, T., Cowell, S., Ward, R. V., Butler, M. J., Sato, H., Seiki, M., Reynolds, J. J., and Murphy, G. (1995). Intermolecular autolytic cleavage can contribute to the activation of progelatinase A by cell membranes. J Biol.Chem. 270, 30479-30485.
- Autio-Harmainen, H., Karttunen, T., Hurskainen, T., Hoyhtya, M., Kauppila, A., and Tryggvason, K. (1993). Expression of 72 kilodalton type IV collagenase (gelatinase A) in benign and malignant ovarian tumors. Lab Invest 69, 312-21.

- Azzam, H. S., Arand, G., Lippman, M. E., and Thompson, E. W. (1993). Association of MMP-2 activation potential with metastatic progression in human breast cancer cell lines independent of MMP- 2 production. J.Natl.Cancer Inst. 85, 1758-1764.
- Azzam, H. S., and Thompson, E. W. (1992). Collagen-induced activation of the Mr 72,000 type IV collagenase in normal and malignant human fibroblastoid cells. Cancer Res. 52, 4540-4544.
- Bachinger, H. P., Bruckner, P., Timpl, R., Prockop, D. J., and Engel, J. (1980). Folding mechanism of the triple helix in type-III collagen and type-III pN-collagen. Role of disulfide bridges and peptide bond isomerization. Eur J Biochem 106, 619-32.
- Baramova, E. N., Bajou, K., Remacle, A., L'Hoir, C., Krell, H. W., Weidle, U. H., Noel, A., and Foidart, J. M. (1997). Involvement of PA/plasmin system in the processing of pro-MMP-9 and in the second step of pro-MMP-2 activation. FEBS Lett 405, 157-62.
- Barsky, S. H., and Gopalakrishna, R. (1987). Increased invasion and spontaneous metastasis of BL6 melanoma with inhibition of the desmoplastic response in C57 BL/6 mice. Cancer Res 47, 1663-7.
- Barsky, S. H., Green, W. R., Grotendorst, G. R., and Liotta, L. A. (1984). Desmoplastic breast carcinoma as a source of human myofibroblasts. Am J Pathol 115, 329-33.
- Barsky, S. H., Siegal, G. P., Jannotta, F., and Liotta, L. A. (1983). Loss of basement membrane components by invasive tumors but not by their benign counterparts. Lab Invest 49, 140-7.

- Bateman, J. F., Pillow, J. J., Mascara, T., Medvedec, S., Ramshaw, J. A., and Cole, W. G. (1987). Cell-layer-associated proteolytic cleavage of the telopeptides of type I collagen in fibroblast culture. Biochem J 245, 677-82.
- Belaouaj, A., Shipley, J. M., Kobayashi, D. K., Zimonjic, D. B., Popescu, N., Silverman, G. A., and Shapiro, S. D. (1995). Human macrophage metalloelastase. Genomic organization, chromosomal location, gene linkage, and tissue-specific expression. J Biol Chem 270, 14568-75.
- Belien, A. T., Paganetti, P. A., and Schwab, M. E. (1999). Membrane-type 1 matrix metalloprotease (MT1-MMP) enables invasive migration of glioma cells in central nervous system white matter. J Cell Biol 144, 373-84.
- Belkhir, A., Richards, C., Whaley, M., McQueen, S. A., and Orr, F. W. (1997). Increased expression of activated matrix metalloproteinase-2 by human endothelial cells after sublethal H₂O₂ exposure. Lab Invest 77, 533-9.
- Benaud, C., Dickson, R. B., and Thompson, E. W. (1998). Roles of the matrix metalloproteinases in mammary gland development and cancer. J Cell Physiol 50, 97-116.
- Berg, R. A., and Prockop, D. J. (1973). The thermal transition of a non-hydroxylated form of collagen. Evidence for a role for hydroxyproline in stabilizing the triple-helix of collagen. Biochem Biophys Res Commun 52, 115-20.
- Berger, J., Tazawa, K., and Prockop, D. J. (1985). Sequential cleavage of type I procollagen by procollagen N-proteinase. An intermediate containing an uncleaved pro alpha 1(I) chain. Biochemistry 24, 600-5.
- Bigg, H. F., Shi, Y. E., Liu, Y. E., Steffensen, B., and Overall, C. M. (1997). Specific, high affinity binding of tissue inhibitor of metalloproteinases-4 (TIMP-4) to the COOH-terminal hemopexin-like domain of human gelatinase A. TIMP-4

- binds progelatinase A and the COOH-terminal domain in a similar manner to TIMP-2. J Biol Chem 272, 15496-500.
- Birembaut, P., Caron, Y., Adnet, J. J., and Foidart, J. M. (1985). Usefulness of basement membrane markers in tumoural pathology. J Pathol 145, 283-96.
- Birk, D. E., Zycband, E. I., Winkelmann, D. A., and Trelstad, R. L. (1990). Collagen fibrillogenesis in situ. Discontinuous segmental assembly in extracellular compartments. Ann N Y Acad Sci 580, 176-94.
- Birkedal-Hansen, H. (1995). Proteolytic remodeling of extracellular matrix. Curr Opin Cell Biol. 7, 728-735.
- Birkedal-Hansen, H., Moore, W. G., Bodden, M. K., Windsor, L. J., Birkedal-Hansen, B., De Carlo, A., and Engler, J. A. (1993). Matrix metalloproteinases: A review. Crit.Rev.Oral Biol.Med. 4, 197-250.
- Biswas, C. (1984). Collagenase stimulation in cocultures of human fibroblasts and human tumor cells. Cancer Lett 24, 201-7.
- Bode, W. (1995). A helping hand for collagenases: the haemopexin-like domain. Structure 3, 527-30.
- Boyd, R. S., and Balkwill, F. R. (1999). MMP-2 release and activation in ovarian carcinoma: the role of fibroblasts. Br J Cancer 80, 315-21.
- Brass, L. F., and Bensusan, H. B. (1976). On the role of the collagen carbohydrate residues in the platelet. Collagen interaction. Biochim Biophys Acta 444, 43-52.
- Brodsky, B., and Eikenberry, E. F. (1982). Characterization of fibrous forms of collagen. Methods Enzymol 82, 127-74.

- Brooks, P. C., Silletti, S., von Schalscha, T. L., Friedlander, M., and Cheresch, D. A. (1998). Disruption of angiogenesis by PEX, a noncatalytic metalloproteinase fragment with integrin binding activity. Cell 92, 391-400.
- Brooks, P. C., Stromblad, S., Sanders, L. C., von Schalscha, T. L., Aimes, R. T., Stetler-Stevenson, W. G., Quigley, J. P., and Cheresch, D. A. (1996). Localization of matrix metalloproteinase MMP-2 to the surface of invasive cells by interaction with integrin alpha v beta 3. Cell 85, 683-93.
- Brown, P. D., Bloxidge, R. E., Stuart, N. S., Gatter, K. C., and Carmichael, J. (1993). Association between expression of activated 72-kilodalton gelatinase and tumor spread in non-small-cell lung carcinoma. J Natl Cancer Inst 85, 574-8.
- Brown, P. D., Levy, A. T., Margulies, I. M., Liotta, L. A., and Stetler-Stevenson, W. G. (1990). Independent expression and cellular processing of Mr 72,000 type IV collagenase and interstitial collagenase in human tumorigenic cell lines. Cancer Res 50, 6184-91.
- Bruckner, P., Eikenberry, E. F., and Prockop, D. J. (1981). Formation of the triple helix of type I procollagen in cellulose. A kinetic model based on cis-trans isomerization of peptide bonds. Eur J Biochem 118, 607-13.
- Bruns, R. R., Hulmes, D. J., Therrien, S. F., and Gross, J. (1979). Procollagen segment-long-spacing crystallites: their role in collagen fibrillogenesis. Proc Natl Acad Sci U S A 76, 313-7.
- Butler, G. S., Apte, S. S., Willenbrock, F., and Murphy, G. (1999). Human tissue inhibitor of metalloproteinases 3 interacts with both the N- and C-terminal domains of gelatinases A and B. Regulation by polyanions. J Biol Chem 274, 10846-51.

- Butler, G. S., Butler, M. J., Atkinson, S. J., Will, H., Tamura, T., van, W. S., Crabbe, T., Clements, J., d'Ortho, M. P., and Murphy, G. (1998). The TIMP2 membrane type 1 metalloproteinase "receptor" regulates the concentration and efficient activation of progelatinase A. A kinetic study. J Biol Chem 273, 871-880.
- Cao, J., Drews, M., Lee, H. M., Conner, C., Bahou, W. F., and Zucker, S. (1999). The 9-kDa N-terminal propeptide domain of MT1-MMP is required for the activation of progelatinase A. Ann N Y Acad Sci 878, 710-2.
- Cao, J., Drews, M., Lee, H. M., Conner, C., Bahou, W. F., and Zucker, S. (1998). The propeptide domain of membrane type 1 matrix metalloproteinase is required for binding of tissue inhibitor of metalloproteinases and for activation of progelatinase A. J Biol Chem 273, 34745-52.
- Cao, J., Hymowitz, M., Conner, C., Bahou, W. F., and Zucker, S. (2000). The propeptide domain of membrane type 1-matrix metalloproteinase acts as an intramolecular chaperone when expressed in trans with the mature sequence in COS-1 cells. J Biol Chem 275, 29648-53.
- Cao, J., Rehmulla, A., Bahou, W., and Zucker, S. (1996). Membrane type matrix metalloproteinase 1 activates pro-gelatinase A without furin cleavage of the N-terminal domain. J Biol Chem 271, 30174-80.
- Cao, J., Sato, H., Takino, T., and Seiki, M. (1995). The C-terminal region of membrane type matrix metalloproteinase is a functional transmembrane domain required for pro-gelatinase A activation. J Biol Chem 270, 801-5.
- Capaldi, M. J., and Chapman, J. A. (1982). The C-terminal extrahelical peptide of type I collagen and its role in fibrillogenesis in vitro. Biopolymers 21, 2291-313.

- Cardinale, G. J., and Udenfriend, S. (1974). Prolyl hydroxylase. Adv Enzymol Relat Areas Mol Biol 41, 245-300.
- Caterina, J. J., Yamada, S., Caterina, N. C., Longenecker, G., Holmback, K., Shi, J., Yermovsky, A. E., Engler, J. A., and Birkedal-Hansen, H. (2000). Inactivating mutation of the mouse tissue inhibitor of metalloproteinases-2(Timp-2) gene alters ProMMP-2 activation [In Process Citation]. J Biol Chem 275, 26416-22.
- Caudroy, S., Polette, M., Tournier, J. M., Burlet, H., Toole, B., Zucker, S., and Birembaut, P. (1999). Expression of the extracellular matrix metalloproteinase inducer (EMMPRIN) and the matrix metalloproteinase-2 in bronchopulmonary and breast lesions. J Histochem Cytochem 47, 1575-80.
- Chambers, A. F., and Matrisian, L. M. (1997). Changing views of the role of matrix metalloproteinases in metastasis. J Biol Chem 272, 1260-1270.
- Chan, D., Lamande, S. R., Cole, W. G., and Bateman, J. F. (1990). Regulation of procollagen synthesis and processing during ascorbate-induced extracellular matrix accumulation in vitro. Biochem J 269, 175-81.
- Clavel, C., Doco, M., Lallemand, A., Laurent, M., and Birembaut, P. (1989). Detection by in situ hybridization of messenger RNAs of collagen types I and IV in murine mammary cancer. Int J Cancer 44, 548-53.
- Cohen-Solal, L., Castanet, J., Meunier, F. J., and Glimcher, M. J. (1986). Proline hydroxylation of collagens synthesized at different temperatures in vivo by two poikilothermic species. Comp Biochem Physiol B 83, 483-6.
- Corcoran, M. L., Hewitt, R. E., Kleiner, D. E., Jr., and Stetler-Stevenson, W. G. (1996). MMP-2: expression, activation and inhibition. Enzyme Protein 49, 7-19.

- Cowell, S., Knauper, V., Stewart, M. L., D'Ortho, M. P., Stanton, H., Hembry, R. M., Lopez-Otin, C., Reynolds, J. J., and Murphy, G. (1998). Induction of matrix metalloproteinase activation cascades based on membrane-type 1 matrix metalloproteinase: associated activation of gelatinase A, gelatinase B and collagenase 3. Biochem J 331, 453-8.
- De Nichilo, M. O., and Yamada, K. M. (1996). Integrin alpha v beta 5-dependent serine phosphorylation of paxillin in cultured human macrophages adherent to vitronectin. J Biol Chem 271, 11016-22.
- Deak, S. B., Glaug, M. R., Pierce, R. A., Bancila, E., Amenta, P., Mackenzie, J. W., Greco, R. S., and Boyd, C. D. (1991). Desmoplasia in benign and malignant breast disease is characterized by alterations in level of mRNAs coding for types I and III procollagen. Matrix 11, 252-8.
- Deryugina, E. I., Bourdon, M. A., Jungwirth, K., Smith, J. W., and Strongin, A. Y. (2000). Functional activation of integrin alpha V beta 3 in tumor cells expressing membrane-type 1 matrix metalloproteinase. Int J Cancer 86, 15-23.
- Deryugina, E. I., Bourdon, M. A., Luo, G. X., Reisfeld, R. A., and Strongin, A. (1997). Matrix metalloproteinase-2 activation modulates glioma cell migration. J Cell Sci 110, 2473-82.
- Doerge, K. J., and Fessler, J. H. (1986). Folding of carboxyl domain and assembly of procollagen I. J Biol Chem 261, 8924-35.
- d'Ortho, M. P., Stanton, H., Butler, M., Atkinson, S. J., Murphy, G., and Hembry, R. M. (1998). MT1-MMP on the cell surface causes focal degradation of gelatin films. FEBS Lett 421, 159-64.
- d'Ortho, M. P., Will, H., Atkinson, S., Butler, G., Messent, A., Gavrilovic, J., Smith, B., Timpl, R., Zardi, L., and Murphy, G. (1997). Membrane-type matrix

- metalloproteinases 1 and 2 exhibit broad-spectrum proteolytic capacities comparable to many matrix metalloproteinases. Eur J Biochem 250, 751-7.
- Du, B., Wang, P., and Guo, X. (1999). Expression of membrane type 1-matrix metalloproteinase in laryngeal carcinoma. Pathol Oncol Res 5, 214-7.
- Ellenrieder, V., Alber, B., Lacher, U., Hendler, S. F., Menke, A., Boeck, W., Wagner, M., Wilda, M., Friess, H., Buchler, M., Adler, G., and Gress, T. M. (2000). Role of MT-MMPs and MMP-2 in pancreatic cancer progression. Int J Cancer 85, 14-20.
- Ellerbroek, S. M., Fishman, D. A., Kearns, A. S., Bafetti, L. M., and Stack, M. S. (1999). Ovarian carcinoma regulation of matrix metalloproteinase-2 and membrane type 1 matrix metalloproteinase through beta1 integrin. Cancer Res 59, 1635-41.
- Emmert-Buck, M. R., Emonard, H. P., Corcoran, M. L., Krutzsch, H. C., Foidart, J. M., and Stetler-Stevenson, W. G. (1995). Cell surface binding of TIMP-2 and pro-MMP-2/TIMP-2 complex. FEBS Lett 364, 28-32.
- Eyre, D. R., Paz, M. A., and Gallop, P. M. (1984). Cross-linking in collagen and elastin. Annu Rev Biochem 53, 717-48.
- Fertala, A., Sieron, A. L., Ganguly, A., Li, S. W., Ala-Kokko, L., Anumula, K. R., and Prockop, D. J. (1994). Synthesis of recombinant human procollagen II in a stably transfected tumour cell line (HT1080). Biochem J 298, 31-7.
- Fessler, L. I., Duncan, K. G., Fessler, J. H., Salo, T., and Tryggvason, K. (1984). Characterization of the procollagen IV cleavage products produced by a specific tumor collagenase. J Biol Chem 259, 9783-9.

- Fitzgerald, J., Lamande, S. R., and Bateman, J. F. (1999). Proteasomal degradation of unassembled mutant type I collagen pro-alpha1(I) chains. J Biol Chem 274, 27392-8.
- Fleischmajer, R., Fisher, L. W., MacDonald, E. D., Jacobs, L., Perlish, J. S., and Termine, J. D. (1991). Decorin interacts with fibrillar collagen of embryonic and adult human skin. J Struct Biol 106, 82-90.
- Foda, H. D., George, S., Conner, C., Drews, M., Tompkins, D. C., and Zucker, S. (1996). Activation of human umbilical vein endothelial cell progelatinase A by phorbol myristate acetate: a protein kinase C-dependent mechanism involving a membrane-type matrix metalloproteinase. Lab Invest 74, 538-45.
- Franklin, T., Morris, W., Edwards, P., Large, M., and Stephenson, R. (2001). Inhibition of prolyl 4-hydroxylase *in vitro* and *in vivo* by members of a novel series of phenanthrourolines. Biochem. J. 53, 333-338.
- Fraser, R. D., and MacRae, T. P. (1979). The crystalline structure of collagen fibrils in tendon. J Mol Biol 127, 129-33.
- Fraser, R. D., MacRae, T. P., and Suzuki, E. (1979). Chain conformation in the collagen molecule. J Mol Biol 129, 463-81.
- Freiberger, H., Grove, D., Sivarajah, A., and Pinnell, S. R. (1980). Procollagen I synthesis in human skin fibroblasts: effect on culture conditions on biosynthesis. J Invest Dermatol 75, 425-30.
- Fridman, R., Bird, R. E., Hoyhtya, M., Oelkuct, M., Komarek, D., Liang, C. M., Berman, M. L., Liotta, L. A., Stetler-Stevenson, W. G., and Fuerst, T. R. (1993). Expression of human recombinant 72 kDa gelatinase and tissue inhibitor of metalloproteinase-2 (TIMP-2): characterization of complex and free enzyme. Biochem J 289, 411-6.

- Gilles, C., Bassuk, J. A., Pulyaeva, H., Sage, E. H., Foidart, J. M., and Thompson, E. W. (1998). SPARC/osteonectin induces matrix metalloproteinase 2 activation in human breast cancer cell lines. Cancer Res 58, 5529-36.
- Gilles, C., Polette, M., Piette, J., Munaut, C., Thompson, E. W., Birembaut, P., and Foidart, J. M. (1996). High level of MT-MMP expression is associated with invasiveness of cervical cancer cells. Int J Cancer 65, 209-13.
- Gilles, C., Polette, M., Seiki, M., Birembaut, P., and Thompson, E. W. (1997). Implication of collagen type I-induced membrane-type 1-matrix metalloproteinase expression and matrix metalloproteinase-2 activation in the metastatic progression of breast carcinoma. Lab Invest 76, 651-60.
- Goldberg, G. I., Strongin, A., Collier, I. E., Genrich, L. T., and Marmer, B. L. (1992). Interaction of 92-kDa type IV collagenase with the tissue inhibitor of metalloproteinases prevents dimerization, complex formation with interstitial collagenase, and activation of the proenzyme with stromelysin. J Biol Chem 267, 4583-91.
- Gomez, D. E., Alonso, D. F., Yoshiji, H., and Thorgeirsson, U. P. (1997). Tissue inhibitors of metalloproteinases: structure, regulation and biological functions. Eur J Cell Biol 74, 111-22.
- Goslen, J. B., Eisen, A. Z., and Bauer, E. A. (1985). Stimulation of skin fibroblast collagenase production by a cytokine derived from basal cell carcinomas. J Invest Dermatol 85, 161-4.
- Grant, M. E., and Prockop, D. J. (1972). The biosynthesis of collagen. 2. N Engl J Med 286, 242-9.

- Greene, J., Wang, M., Liu, Y. E., Raymond, L. A., Rosen, C., and Shi, Y. E. (1996). Molecular cloning and characterization of human tissue inhibitor of metalloproteinase 4. J Biol Chem 271, 30375-80.
- Guo, H., Zucker, S., Gordon, M. K., Toole, B. P., and Biswas, C. (1997). Stimulation of matrix metalloproteinase production by recombinant extracellular matrix metalloproteinase inducer from transfected Chinese hamster ovary cells. J Biol Chem 272, 24-7.
- Haas, T. L., Davis, S. J., and Madri, J. A. (1998). Thource-dimensional type I collagen lattices induce coordinate expression of matrix metalloproteinases MT1-MMP and MMP-2 in microvascular endothelial cells. J Biol Chem 273, 3604-10.
- Haas, T. L., Stitelman, D., Davis, S. J., Apte, S. S., and Madri, J. A. (1999). Egr-1 mediates extracellular matrix-driven transcription of membrane type 1 matrix metalloproteinase in endothelium. J Biol Chem 274, 22679-85.
- Hanemaaijer, R., Verheijen, J. H., Maguire, T. M., Visser, H., Toet, K., McDermott, E., O'Higgins, N., and Duffy, M. J. (2000). Increased gelatinase-A and gelatinase-B activities in malignant vs. benign breast tumors. Int J Cancer 86, 204-7.
- Hattori, S., Adachi, E., Ebihara, T., Shirai, T., Someki, I., and Irie, S. (1999). Alkali-treated collagen retained the triple helical conformation and the ligand activity for the cell adhesion via alpha2beta1 integrin. J Biochem (Tokyo) 125, 676-84.
- Heinegard, D., and Oldberg, A. (1989). Structure and biology of cartilage and bone matrix noncollagenous macromolecules. Faseb J 3, 2042-51.

- Helseth, D. L., Jr., and Veis, A. (1981). Collagen self-assembly in vitro. Differentiating specific telopeptide-dependent interactions using selective enzyme modification and the addition of free amino telopeptide. J Biol Chem 256, 7118-28.
- Heppner, K. J., Matrisian, L. M., Jensen, R. A., and Rodgers, W. H. (1996). Expression of most matrix metalloproteinase family members in breast cancer represents a tumor-induced host response. Am J Pathol 149, 273-82.
- Hernandez-Barrantes, S., Toth, M., Bernardo, M. M., Yurkova, M., Gervasi, D. C., Raz, Y., Sang, Q. A., and Fridman, R. (2000). Binding of active (57 kDa) membrane type 1-matrix metalloproteinase (MT1-MMP) to tissue inhibitor of metalloproteinase (TIMP)-2 regulates MT1-MMP processing and pro-MMP-2 activation. J Biol Chem 275, 12080-9.
- Hibbs, M. S., Hasty, K. A., Seyer, J. M., Kang, A. H., and Mainardi, C. L. (1985). Biochemical and immunological characterization of the secreted forms of human neutrophil gelatinase. J Biol Chem 260, 2493-500.
- Himelstein, B. P., Canete-Soler, R., Bernhard, E. J., and Muschel, R. J. (1994). Induction of fibroblast 92 kDa gelatinase/type IV collagenase expression by direct contact with metastatic tumor cells. J Cell Sci 107, 477-86.
- Hojima, Y., van der Rest, M., and Prockop, D. J. (1985). Type I procollagen carboxyl-terminal proteinase from chick embryo tendons. Purification and characterization. J Biol Chem 260, 15996-6003.
- Holliday, L. S., Welgus, H. G., Fliszar, C. J., Veith, G. M., Jeffrey, J. J., and Gluck, S. L. (1997). Initiation of osteoclast bone resorption by interstitial collagenase. J Biol Chem 272, 22053-8.

- Holmbeck, K., Bianco, P., Caterina, J., Yamada, S., Kromer, M., Kuznetsov, S. A., Mankani, M., Robey, P. G., Poole, A. R., Pidoux, I., Ward, J. M., and Birkedal-Hansen, H. (1999). MT1-MMP-deficient mice develop dwarfism, osteopenia, arthouritis, and connective tissue disease due to inadequate collagen turnover. Cell 99, 81-92.
- Hotary, K., Allen, E., Punturieri, A., Yana, I., and Weiss, S. J. (2000). Regulation of cell invasion and morphogenesis in a thouree-dimensional type I collagen matrix by membrane-type matrix metalloproteinases 1, 2, and 3. J Cell Biol 149, 1309-23.
- Howard, E. W., and Banda, M. J. (1991). Binding of tissue inhibitor of metalloproteinases 2 to two distinct sites on human 72-kDa gelatinase. Identification of a stabilization site. J Biol Chem 266, 17972-7.
- Howard, E. W., Bullen, E. C., and Banda, M. J. (1991). Preferential inhibition of 72- and 92-kDa gelatinases by tissue inhibitor of metalloproteinases-2. J Biol Chem 266, 13070-5.
- Howard, E. W., Bullen, E. C., and Banda, M. J. (1991). Regulation of the autoactivation of human 72-kDa progelatinase by tissue inhibitor of metalloproteinases-2. J Biol Chem 266, 13064-9.
- Hudson, B. G., Reeders, S. T., and Tryggvason, K. (1993). Type IV collagen: structure, gene organization, and role in human diseases. Molecular basis of Goodpasture and Alport syndromes and diffuse leiomyomatosis. J Biol Chem 268, 26033-6.
- Hulmes, D. J. (1992). The collagen superfamily--diverse structures and assemblies. Essays Biochem 27, 49-67.

- Hulmes, D. J., Bruns, R. R., and Gross, J. (1983). On the state of aggregation of newly secreted procollagen. Proc Natl Acad Sci U S A 80, 388-92.
- Hulmes, D. J., Kadler, K. E., Mould, A. P., Hojima, Y., Holmes, D. F., Cummings, C., Chapman, J. A., and Prockop, D. J. (1989). Pleomorphism in type I collagen fibrils produced by persistence of the procollagen N-propeptide. J Mol Biol 210, 337-45.
- Imai, K., Ohuchi, E., Aoki, T., Nomura, H., Fujii, Y., Sato, H., Seiki, M., and Okada, Y. (1996). Membrane-type matrix metalloproteinase 1 is a gelatinolytic enzyme and is secreted in a complex with tissue inhibitor of metalloproteinases 2. Cancer Res 56, 2707-10.
- Ito, A., Mukaiyama, A., Itoh, Y., Nagase, H., Thogersen, I. B., Enghild, J. J., Sasaguri, Y., and Mori, Y. (1996). Degradation of interleukin 1beta by matrix metalloproteinases. J Biol Chem 271, 14657-60.
- Itoh, Y., Ito, A., Iwata, K., Tanzawa, K., Mori, Y., and Nagase, H. (1998). Plasma membrane-bound tissue inhibitor of metalloproteinases (TIMP)-2 specifically inhibits matrix metalloproteinase 2 (gelatinase A) activated on the cell surface. J Biol Chem 273, 24360-7.
- Itoh, Y., Kajita, M., Kinoh, H., Mori, H., Okada, A., and Seiki, M. (1999). Membrane type 4 matrix metalloproteinase (MT4-MMP, MMP-17) is a glycosylphosphatidylinositol-anchored proteinase. J Biol Chem 274, 34260-6.
- Jaenisch, R., Harbers, K., Schnieke, A., Lohler, J., Chumakov, I., Jahner, D., Grotkopp, D., and Hoffmann, E. (1983). Germline integration of moloney murine leukemia virus at the Mov13 locus leads to recessive lethal mutation and early embryonic death. Cell 32, 209-16.

- Jamall, I. S., Finelli, V. N., and Que Hee, S. S. (1981). A simple method to determine nanogram levels of 4-hydroxyproline in biological tissues. *Anal Biochem* 112, 70-5.
- Jimenez, S., Harsch, M., and Rosenbloom, J. (1973). Hydroxyproline stabilizes the triple helix of chick tendon collagen. *Biochem Biophys Res Commun* 52, 106-14.
- Kadler, K. E., Hojima, Y., and Prockop, D. J. (1987). Assembly of collagen fibrils de novo by cleavage of the type I pC-collagen with procollagen C-proteinase. Assay of critical concentration demonstrates that collagen self-assembly is a classical example of an entropy-driven process. *J Biol Chem* 262, 15696-701.
- Kadler, K. E., Hojima, Y., and Prockop, D. J. (1990). Collagen fibrils in vitro grow from pointed tips in the C- to N-terminal direction. *Biochem J* 268, 339-43.
- Kadler, K. E., Holmes, D. F., Trotter, J. A., and Chapman, J. A. (1996). Collagen fibril formation. *Biochem J* 316, 1-11.
- Kao, R. T., Hall, J., and Stern, R. (1986). Collagen and elastin synthesis in human stroma and breast carcinoma cell lines: modulation by the extracellular matrix. *Connect Tissue Res* 14, 245-55.
- Kataoka, H., DeCastro, R., Zucker, S., and Biswas, C. (1993). Tumor cell-derived collagenase-stimulatory factor increases expression of interstitial collagenase, stromelysin, and 72-kDa gelatinase. *Cancer Res* 53, 3154-8.
- Kinoshita, T., Sato, H., Okada, A., Ohuchi, E., Imai, K., Okada, Y., and Seiki, M. (1998). TIMP-2 promotes activation of progelatinase A by membrane-type 1 matrix metalloproteinase immobilized on agarose beads. *J Biol Chem* 273, 16098-103.

- Kinoshita, T., Sato, H., Takino, T., Itoh, M., Akizawa, T., and Seiki, M. (1996). Processing of a precursor of 72-kilodalton type IV collagenase/gelatinase A by a recombinant membrane-type 1 matrix metalloproteinase. Cancer Res 56, 2535-8.
- Kivirikko, K. I., and Myllyla, R. (1979). Collagen glycosyltransferases. Int Rev Connect Tissue Res 8, 23-72.
- Kivirikko, K. I., and Myllyla, R. (1982). Posttranslational enzymes in the biosynthesis of collagen: intracellular enzymes. Methods Enzymol 82, 245-304.
- Knauper, V., Will, H., Lopez-Otin, C., Smith, B., Atkinson, S. J., Stanton, H., Hembry, R. M., and Murphy, G. (1996). Cellular mechanisms for human procollagenase-3 (MMP-13) activation. Evidence that MT1-MMP (MMP-14) and gelatinase a (MMP-2) are able to generate active enzyme. J Biol Chem 271, 17124-31.
- Koshihara, T., Hosotani, R., Wada, M., Miyamoto, Y., Fujimoto, K., Lee, J. U., Doi, R., Arai, S., and Imamura, M. (1998). Involvement of matrix metalloproteinase-2 activity in invasion and metastasis of pancreatic carcinoma. Cancer 82, 642-650.
- Koshikawa, N., Giannelli, G., Cirulli, V., Miyazaki, K., and Quaranta, V. (2000). Role of cell surface metalloprotease MT1-MMP in epithelial cell migration over laminin-5. J Cell Biol 148, 615-24.
- Krane, S. M., Byrne, M. H., Lemaitre, V., Henriot, P., Jeffrey, J. J., Witter, J. P., Liu, X., Wu, H., Jaenisch, R., and Eeckhout, Y. (1996). Different collagenase gene products have different roles in degradation of type I collagen. J Biol Chem 271, 28509-15.

- Kurschat, P., Zigrino, P., Nischt, R., Breitkopf, K., Steurer, P., Klein, C. E., Krieg, T., and Mauch, C. (1999). Tissue inhibitor of matrix metalloproteinase-2 regulates matrix metalloproteinase-2 activation by modulation of membrane-type 1 matrix metalloproteinase activity in high and low invasive melanoma cell lines. J Biol Chem 274, 21056-62.
- Lamberg, A., Helaakoski, T., Myllyharju, J., Peltonen, S., Notbohm, H., Pihlajaniemi, T., and Kivirikko, K. I. (1996). Characterization of human type III collagen expressed in a baculovirus system. Production of a protein with a stable triple helix requires coexpression with the two types of recombinant prolyl 4-hydroxylase subunit. J Biol Chem 271, 11988-95.
- Lambert, C. A., Soudant, E. P., Nusgens, B. V., and Lapiere, C. M. (1992). Pretranslational regulation of extracellular matrix macromolecules and collagenase expression in fibroblasts by mechanical forces. Lab Invest 66, 444-51.
- Lee, A. Y., Akers, K. T., Collier, M., Li, L., Eisen, A. Z., and Seltzer, J. L. (1997). Intracellular activation of gelatinase A (72-kDa type IV collagenase) by normal fibroblasts. Proc Natl Acad Sci U S A 94, 4424-9.
- Lee, S., Solow-Cordero, D. E., Kessler, E., Takahara, K., and Greenspan, D. S. (1997). Transforming growth factor-beta regulation of bone morphogenetic protein-1/procollagen C-proteinase and related proteins in fibrogenic cells and keratinocytes. J Biol Chem 272, 19059-66.
- Lee, S. T., Kessler, E., and Greenspan, D. S. (1990). Analysis of site-directed mutations in human pro-alpha 2(I) collagen which block cleavage by the C-proteinase. J Biol Chem 265, 21992-6.

- Leibovich, S. J., and Weiss, J. B. (1970). Electron microscope studies of the effects of endo- and exopeptidase digestion on tropocollagen. A novel concept of the role of terminal regions in fibrillogenesis. Biochim Biophys Acta 214, 445-54.
- Lepage, T., and Gache, C. (1990). Early expression of a collagenase-like hatching enzyme gene in the sea urchin embryo. Embo J 9, 3003-12.
- Levi, E., Fridman, R., Miao, H. Q., Ma, Y. S., Yayon, A., and Vlodavsky, I. (1996). Matrix metalloproteinase 2 releases active soluble ectodomain of fibroblast growth factor receptor 1. Proc Natl Acad Sci U S A 93, 7069-74.
- Liotta, L. A., Rao, C. N., and Barsky, S. H. (1983). Tumor invasion and the extracellular matrix. Lab Invest 49, 636-49.
- Liotta, L. A., Steeg, P. S., and Stetler-Stevenson, W. G. (1991). Cancer metastasis and angiogenesis: an imbalance of positive and negative regulation. Cell 64, 327-36.
- Liotta, L. A., Tryggvason, K., Garbisa, S., Hart, I., Foltz, C. M., and Shafie, S. (1980). Metastatic potential correlates with enzymatic degradation of basement membrane collagen. Nature 284, 67-8.
- Lohi, J., Lehti, K., Westermarck, J., Kahari, V. M., and Keski-Oja, J. (1996). Regulation of membrane-type matrix metalloproteinase-1 expression by growth factors and phorbol 12-myristate 13-acetate. Eur J Biochem 239, 239-47.
- Lohler, J., Timpl, R., and Jaenisch, R. (1984). Embryonic lethal mutation in mouse collagen I gene causes rupture of blood vessels and is associated with erythropoietic and mesenchymal cell death. Cell 38, 597-607.

- Long, L., Navab, R., and Brodt, P. (1998). Regulation of the Mr 72,000 type IV collagenase by the type I insulin-like growth factor receptor. Cancer Res 58, 3243-7.
- MacDougall, J. R., and Matrisian, L. M. (1995). Contributions of tumor and stromal matrix metalloproteinases to tumor progression, invasion and metastasis. Cancer Metastasis Rev 14, 351-62.
- Maquoi, E., Frankenne, F., Noel, A., Krell, H. W., Grams, F., and Foidart, J. M. (2000). Type IV collagen induces matrix metalloproteinase 2 activation in HT1080 fibrosarcoma cells. Exp Cell Res 261, 348-59.
- Maquoi, E., Noel, A., Frankenne, F., Angliker, H., Murphy, G., and Foidart, J. M. (1998). Inhibition of matrix metalloproteinase 2 maturation and HT1080 invasiveness by a synthetic furin inhibitor. FEBS Lett 424, 262-6.
- Martinez-Hernandez, A., Francis, D. J., and Silverberg, S. G. (1977). Elastosis and other stromal reactions in benign and malignant breast tissue: an ultrastructural study. Cancer 40, 600-6.
- Massova, I., Kotra, L. P., Fridman, R., and Mobashery, S. (1998). Matrix metalloproteinases: structures, evolution, and diversification. Faseb J 12, 1075-95.
- Matrisian, L. M. (1994). Matrix metalloproteinase gene expression. Ann N Y Acad Sci 732, 42-50.
- Matsumoto, S., Katoh, M., Saito, S., Watanabe, T., and Masuho, Y. (1997). Identification of soluble type of membrane-type matrix metalloproteinase-3 formed by alternatively spliced mRNA. Biochim Biophys Acta 1354, 159-70.
- Mazzieri, R., Masiero, L., Zanetta, L., Monea, S., Onisto, M., Garbisa, S., and Mignatti, P. (1997). Control of type IV collagenase activity by components of

the urokinase-plasmin system: a regulatory mechanism with cell-bound reactants. Embo J 16, 2319-32.

McKerrow, J. H., Bhargava, V., Hansell, E., Huling, S., Kuwahara, T., Matley, M., Coussens, L., and Warren, R. (2000). A functional proteomics screen of proteases in colorectal carcinoma. Mol Med 6, 450-60.

Minafra, S., Minafra, I. P., Tomasino, R. M., and Sciarrino, S. (1984). Collagen composition in the ductal infiltrating carcinoma of human breast. Cell Biol Int Rep 8, 79-85.

Minamoto, T., Ooi, A., Okada, Y., Mai, M., Nagai, Y., and Nakanishi, I. (1988). Desmoplastic reaction of gastric carcinoma: a light- and electron-microscopic immunohistochemical analysis using collagen type-specific antibodies. Hum Pathol 19, 815-21.

Miyahara, M., Njieha, F. K., and Prockop, D. J. (1982). Formation of collagen fibrils in vitro by cleavage of procollagen with procollagen proteinases. J Biol Chem 257, 8442-8.

Murad, S., Grove, D., Lindberg, K. A., Reynolds, G., Sivarajah, A., and Pinnell, S. R. (1981). Regulation of collagen synthesis by ascorbic acid. Proc Natl Acad Sci U S A 78, 2879-82.

Murphy, G., Segain, J. P., O'Shea, M., Cockett, M., Ioannou, C., Lefebvre, O., Chambon, P., and Basset, P. (1993). The 28-kDa N-terminal domain of mouse stromelysin-3 has the general properties of a weak metalloproteinase. J Biol Chem 268, 15435-41.

Murphy, G., Stanton, H., Cowell, S., Butler, G., Knauper, V., Atkinson, S., and Gavrilovic, J. (1999). Mechanisms for pro matrix metalloproteinase activation. Apmis 107, 38-44.

- Murphy, G., Willenbrock, F., Ward, R. V., Cockett, M. I., Eaton, D., and Docherty, A. J. (1992). The C-terminal domain of 72 kDa gelatinase A is not required for catalysis, but is essential for membrane activation and modulates interactions with tissue inhibitors of metalloproteinases [published erratum appears in *Biochem J* 1992 Jun 15;284(Pt 3):935]. *Biochem J* 283, 637-41.
- Nagase, H. (1997). Activation mechanisms of matrix metalloproteinases. *Biol Chem* 378, 151-60.
- Nagase, H., Ogata, Y., Suzuki, K., Enghild, J. J., and Salvesen, G. (1991). Substrate specificities and activation mechanisms of matrix metalloproteinases. *Biochem Soc Trans* 19, 715-8.
- Nagase, H., and Woessner, J. F., Jr. (1999). Matrix metalloproteinases. *J Biol Chem* 274, 21491-4.
- Nakahara, H., Howard, L., Thompson, E. W., Sato, H., Seiki, M., Yeh, Y., and Chen, W. T. (1997). Transmembrane/cytoplasmic domain-mediated membrane type 1-matrix metalloprotease docking to invadopodia is required for cell invasion. *Proc Natl Acad Sci U S A* 94, 7959-64.
- Nakamura, T., Jamieson, G. A., Okuma, M., Kambayashi, J., and Tandon, N. N. (1998). Platelet adhesion to native type I collagen fibrils. Role of GPVI in divalent cation-dependent and -independent adhesion and thromboxane A₂ generation. *J Biol Chem* 273, 4338-44.
- Nakamura, T., Kambayashi, J., Okuma, M., and Tandon, N. N. (1999). Activation of the GP IIb-IIIa complex induced by platelet adhesion to collagen is mediated by both $\alpha 2\beta 1$ integrin and GP VI. *J Biol Chem* 274, 11897-903.
- Nguyen, M., Arkell, J., and Jackson, C. J. (2000). Activated protein C directly activates human endothelial gelatinase A. *J Biol Chem* 275, 9095-8.

- Nguyen, M., Arkell, J., and Jackson, C. J. (2000). Thourree-dimensional collagen matrices induce delayed but sustained activation of gelatinase A in human endothelial cells via MT1-MMP. Int J Biochem Cell Biol 32, 621-31.
- Nguyen, M., Arkell, J., and Jackson, C. J. (1999). Thourombin rapidly and efficiently activates gelatinase A in human microvascular endothelial cells via a mechanism independent of active MT1 matrix metalloproteinase. Lab Invest 79, 467-75.
- Njieha, F. K., Morikawa, T., Tuderman, L., and Prockop, D. J. (1982). Partial purification of a procollagen C-proteinase. Inhibition by synthetic peptides and sequential cleavage of type I procollagen. Biochemistry 21, 757-64.
- Nomura, H., Sato, H., Seiki, M., Mai, M., and Okada, Y. (1995). Expression of membrane-type matrix metalloproteinase in human gastric carcinomas. Cancer Res. 55, 3263-3266.
- Nusgens, B. V., Verellen-Dumoulin, C., Hermanns-Le, T., De Paepe, A., Nuytinck, L., Pierard, G. E., and Lapiere, C. M. (1992). Evidence for a relationship between Ehlers-Danlos type VII C in humans and bovine dermatosparaxis. Nat Genet 1, 214-7.
- Oberg, A., Hoyhtya, M., Tavelin, B., Stenling, R., and Lindmark, G. (2000). Limited value of preoperative serum analyses of matrix metalloproteinases (MMP-2, MMP-9) and tissue inhibitors of matrix metalloproteinases (TIMP-1, TIMP-2) in colorectal cancer. Anticancer Res 20, 1085-91.
- Ogata, Y., Itoh, Y., and Nagase, H. (1995). Steps involved in activation of the pro-matrix metalloproteinase 9 (progelatinase B)-tissue inhibitor of metalloproteinases-1 complex by 4-aminophenylmercuric acetate and proteinases. J Biol Chem 270, 18506-11.

- Ohtani, H., Kuroiwa, A., Obinata, M., Ooshima, A., and Nagura, H. (1992). Identification of type I collagen-producing cells in human gastrointestinal carcinomas by non-radioactive in situ hybridization and immunoelectron microscopy. J Histochem Cytochem 40, 1139-46.
- Ohuchi, E., Imai, K., Fujii, Y., Sato, H., Seiki, M., and Okada, Y. (1997). Membrane type 1 matrix metalloproteinase digests interstitial collagens and other extracellular matrix macromolecules. J Biol Chem 272, 2446-51.
- Okada, Y., Morodomi, T., Enghild, J. J., Suzuki, K., Yasui, A., Nakanishi, I., Salvesen, G., and Nagase, H. (1990). Matrix metalloproteinase 2 from human rheumatoid synovial fibroblasts. purification and activation of the precursor and enzymic properties. Eur.J.Biochem. 194, 721-730.
- Okumura, Y., Sato, H., Seiki, M., and Kido, H. (1997). Proteolytic activation of the precursor of membrane type 1 matrix metalloproteinase by human plasmin. A possible cell surface activator. FEBS Lett 402, 181-4.
- Overall, C. M., King, A. E., Sam, D. K., Ong, A. D., Lau, T. T., Wallon, U. M., DeClerck, Y. A., and Atherstone, J. (1999). Identification of the tissue inhibitor of metalloproteinases-2 (TIMP-2) binding site on the hemopexin carboxyl domain of human gelatinase A by site-directed mutagenesis. The hierarchical role in binding TIMP-2 of the unique cationic clusters of hemopexin modules III and IV. J Biol Chem 274, 4421-9.
- Overall, C. M., Wrana, J. L., and Sodek, J. (1991). Transcriptional and post-transcriptional regulation of 72-kDa gelatinase/type IV collagenase by transforming growth factor-beta 1 in human fibroblasts. Comparisons with collagenase and tissue inhibitor of matrix metalloproteinase gene expression. J Biol Chem 266, 14064-71.

- Pak, J. H., Liu, C. Y., Huangpu, J., and Graham, J. S. (1997). Construction and characterization of the soybean leaf metalloproteinase cDNA. FEBS Lett 404, 283-8.
- Payne, K. J., King, T. A., and Holmes, D. F. (1986). Collagen fibrillogenesis in vitro: an investigation of the thermal memory effect and of the early events occurring during fibril assembly using dynamic light scattering. Biopolymers 25, 1185-207.
- Pei, D. (1999). Identification and characterization of the fifth membrane-type matrix metalloproteinase MT5-MMP. J Biol Chem 274, 8925-32.
- Pei, D., Majmudar, G., and Weiss, S. J. (1994). Hydrolytic inactivation of a breast carcinoma cell-derived serpin by human stromelysin-3. J Biol Chem 269, 25849-55.
- Pei, D., and Weiss, S. J. (1995). Furin-dependent intracellular activation of the human stromelysin-3 zymogen. Nature 375, 244-7.
- Pei, D., and Weiss, S. J. (1996). Transmembrane-deletion mutants of the membrane-type matrix metalloproteinase-1 process progelatinase A and express intrinsic matrix-degrading activity. J Biol Chem 271, 9135-40.
- Pilcher, B. K., Dumin, J. A., Sudbeck, B. D., Krane, S. M., Welgus, H. G., and Parks, W. C. (1997). The activity of collagenase-1 is required for keratinocyte migration on a type I collagen matrix. J Cell Biol 137, 1445-57.
- Polette, M., and Birembaut, P. (1996). Matrix metalloproteinases in breast cancer. The Breast Journal 2, 209-220.
- Poulsom, R., Pignatelli, M., Stetler-Stevenson, W. G., Liotta, L. A., Wright, P. A., Jeffery, R. E., Longcroft, J. M., Rogers, L., and Stamp, G. W. (1992). Stromal

expression of 72 kda type IV collagenase (MMP-2) and TIMP-2 mRNAs in colorectal neoplasia. 141, 389-396.

Preaux, A. M., Mallat, A., Nhieu, J. T., D'Ortho, M. P., Hembry, R. M., and Mavier, P. (1999). Matrix metalloproteinase-2 activation in human hepatic fibrosis regulation by cell-matrix interactions. Hepatology 30, 944-50.

Prockop, D. J., and Fertala, A. (1998). Inhibition of the self-assembly of collagen I into fibrils with synthetic peptides. Demonstration that assembly is driven by specific binding sites on the monomers. J Biol Chem 273, 15598-604.

Prockop, D. J., and Kivirikko, K. I. (1995). Collagens: molecular biology, diseases, and potentials for therapy. Annu Rev Biochem 64, 403-34.

Prockop, D. J., Kivirikko, K. I., Tuderman, L., and Guzman, N. A. (1979). The biosynthesis of collagen and its disorders (second of two parts). N Engl J Med 301, 77-85.

Pucci Minafra, I., Luparello, C., Sciarrino, S., Tomasino, R. M., and Minafra, S. (1985). Quantitative determination of collagen types present in the ductal infiltrating carcinoma of human mammary gland. Cell Biol Int Rep 9, 291-6.

Puente, X. S., Pendas, A. M., Llano, E., Velasco, G., and Lopez-Otin, C. (1996). Molecular cloning of a novel membrane-type matrix metalloproteinase from a human breast carcinoma. Cancer Res 56, 944-9.

Pulyaeva, H., Bueno, J., Polette, M., Birembaut, P., Sato, H., Seiki, M., and Thompson, E. W. (1997). MT1-MMP correlates with MMP-2 activation potential seen after epithelial to mesenchymal transition in human breast carcinoma cells [published erratum appears in Clin Exp Metastasis 1997 May;15(3):338]. Clin Exp Metastasis 15, 111-20.

- Pyke, C., Ralfkiaer, E., Huhtala, P., Hurskainen, T., Dano, K., and Tryggvason, K. (1992). Localization of messenger rna for mr 72,000 and 92,000 type iv collagenases in human skin cancers by in situ hybridization. J Biol Chem 267, 1336-1341.
- Reiser, K., McCormick, R. J., and Rucker, R. B. (1992). Enzymatic and nonenzymatic cross-linking of collagen and elastin. Faseb J 6, 2439-49.
- Romanic, A. M., Adachi, E., Hojima, Y., Engel, J., and Prockop, D. J. (1992). Polymerization of pNcollagen I and copolymerization of pNcollagen I with collagen I. A kinetic, thermodynamic, and morphologic study. J Biol Chem 267, 22265-71.
- Ruangpanit, N., Chan, D., Holmbeck, K., Birkedal-Hansen, H., Polarek, J., Yang, C., Bateman, J. F., and Thompson, E. W. (2001). Gelatinase A (MMP-2) activation by skin fibroblasts: Dependence on MT1-MMP expression and fibrillar collagen form. Matrix Biology, in press.
- Salo, T., Liotta, L. A., and Tryggvason, K. (1983). Purification and characterization of a murine basement membrane collagen-degrading enzyme secreted by metastatic tumor cells. J Biol Chem 258, 3058-63.
- Santoro, S. A., Rajpara, S. M., Staatz, W. D., and Woods, V. L., Jr. (1988). Isolation and characterization of a platelet surface collagen binding complex related to VLA-2. Biochem Biophys Res Commun 153, 217-23.
- Sasaguri, Y., Komiya, S., Sugama, K., Suzuki, K., Inoue, A., Morimatsu, M., and Nagase, H. (1992). Production of matrix metalloproteinases 2 and 3 (stromelysin) by stromal cells of giant cell tumor of bone. Am J Pathol 141, 611-21.
- Sato, H., Kinoshita, T., Takino, T., Nakayama, K., and Seiki, M. (1996). Activation of a recombinant membrane type 1-matrix metalloproteinase (MT1-MMP) by

- furin and its interaction with tissue inhibitor of metalloproteinases (TIMP)-2. FEBS Lett 393, 101-4.
- Sato, H., and Seiki, M. (1996). Membrane-type matrix metalloproteinases (MT-MMPs) in tumor metastasis. J Biochem (Tokyo) 119, 209-15.
- Sato, H., Takino, T., Kinoshita, T., Imai, K., Okada, Y., Stetler Stevenson, W. G., and Seiki, M. (1996). Cell surface binding and activation of gelatinase A induced by expression of membrane-type-1-matrix metalloproteinase (MT1-MMP). FEBS Lett 385, 238-40.
- Sato, H., Takino, T., Okada, Y., Cao, J., Shinagawa, A., Yamamoto, E., and Seiki, M. (1994). A matrix metalloproteinase expressed on the surface of invasive tumour cells. Nature 370, 61-65.
- Sato, M., Ishikawa, O., Abe, M., and Miyachi, Y. (1998). Opposite effects of tumour necrosis factor-alpha on type I and III collagen gene expression by human dermal fibroblasts in monolayer and thource-dimensional cultures. Br J Dermatol 138, 118-21.
- Sato, T., Kondo, T., Fujisawa, T., Seiki, M., and Ito, A. (1999). Furin-independent pathway of membrane type 1-matrix metalloproteinase activation in rabbit dermal fibroblasts. J Biol Chem 274, 37280-4.
- Schnieke, A., Dziadek, M., Bateman, J., Mascara, T., Harbers, K., Gelinas, R., and Jaenisch, R. (1987). Introduction of the human pro alpha 1(I) collagen gene into pro alpha 1(I)-deficient Mov-13 mouse cells leads to formation of functional mouse-human hybrid type I collagen. Proc Natl Acad Sci U S A 84, 764-8.

- Schnieke, A., Harbers, K., and Jaenisch, R. (1983). Embryonic lethal mutation in mice induced by retrovirus insertion into the alpha 1(I) collagen gene. Nature 304, 315-20.
- Scott, J. E. (1988). Proteoglycan-fibrillar collagen interactions. Biochem J 252, 313-23.
- Seidah, N. G., Chouretien, M., and Day, R. (1994). The family of subtilisin/kexin like pro-protein and pro-hormone convertases: divergent or shared functions. Biochimie 76, 197-209.
- Seiki, M. (1999). Membrane-type matrix metalloproteinases. Apmis 107, 137-43.
- Seltzer, J. L., Lee, A. Y., Akers, K. T., Sudbeck, B., Southon, E. A., Wayner, E. A., and Eisen, A. Z. (1994). Activation of 72-kDa type IV collagenase/gelatinase by normal fibroblasts in collagen lattices is mediated by integrin receptors but is not related to lattice contraction. Exp Cell Res 213, 365-74.
- Shaw, L. M., and Olsen, B. R. (1991). FACIT collagens: diverse molecular bridges in extracellular matrices. Trends Biochem Sci 16, 191-4.
- Siegal, G. P., Barsky, S. H., Terranova, V. P., and Liotta, L. A. (1981). Stages of neoplastic transformation of human breast tissue as monitored by dissolution of basement membrane components. An immunoperoxidase study. Invasion Metastasis 1, 54-70.
- Smith, L. T., Wertelecki, W., Milstone, L. M., Petty, E. M., Seashore, M. R., Braverman, I. M., Jenkins, T. G., and Byers, P. H. (1992). Human dermatosparaxis: a form of Ehlers-Danlos syndrome that results from failure to remove the amino-terminal propeptide of type I procollagen. Am J Hum Genet 51, 235-44.

- Soini, Y., Hurskainen, T., Hoyhtya, M., Oikarinen, A., and Autio-Harminen, H. (1994). 72 KD and 92 KD type IV collagenase, type IV collagen, and laminin mRNAs in breast cancer: a study by in situ hybridization. J Histochem Cytochem 42, 945-51.
- Soini, Y., Paakko, P., and Autio-Harminen, H. (1993). Genes of laminin B1 chain, alpha 1 (IV) chain of type IV collagen, and 72-kd type IV collagenase are mainly expressed by the stromal cells of lung carcinomas. Am J Pathol 142, 1622-30.
- Southby, J., O'Keeffe, L. M., Martin, T. J., and Gillespie, M. T. (1995). Alternative promoter usage and mRNA splicing pathways for parathyroid hormone-related protein in normal tissues and tumours. Br J Cancer 72, 702-7.
- Stanton, H., Gavrilovic, J., Atkinson, S. J., d'Ortho, M. P., Yamada, K. M., Zardi, L., and Murphy, G. (1998). The activation of ProMMP-2 (gelatinase A) by HT1080 fibrosarcoma cells is promoted by culture on a fibronectin substrate and is concomitant with an increase in processing of MT1-MMP (MMP-14) to a 45 kDa form. J Cell Sci 111, 2789-98.
- Steffensen, B., Wallon, U. M., and Overall, C. M. (1995). Extracellular matrix binding properties of recombinant fibronectin type II-like modules of human 72-kDa gelatinase/type IV collagenase. High affinity binding to native type I collagen but not native type IV collagen. J Biol Chem 270, 11555-66.
- Sternlicht, M. D., and Bergers, G. (2000). Matrix metalloproteinases as emerging targets in anticancer therapy: status and prospects. Emerging Therapeutic Targets 4, 609 - 633.

- Stetler-Stevenson, W. G., Aznavoorian, S., and Liotta, L. A. (1993). Tumor cell interactions with the extracellular matrix during invasion and metastasis. *Journal of Cellular Biochemistry* 9, 541-573.
- Stetler-Stevenson, W. G., Kruttsch, H. C., Wachter, M. P., Margulies, I. M., and Liotta, L. A. (1989). The activation of human type IV collagenase proenzyme. Sequence identification of the major conversion product following organomercurial activation. *Journal of Biological Chemistry* 264, 1353-1356.
- Stocker, W., Grams, F., Baumann, U., Reinemer, P., Gomis-Ruth, F. X., McKay, D. B., and Bode, W. (1995). The metzincins--topological and sequential relations between the astacins, adamalysins, serralysins, and matrixins (collagenases) define a superfamily of zinc-peptidases. *Protein Sci* 4, 823-40.
- Strongin, A. Y., Collier, I., Bannikov, G., Marmer, B. L., Grant, G. A., and Goldberg, G. I. (1995). Mechanism of cell surface activation of 72-kDa type IV collagenase. Isolation of the activated form of the membrane metalloprotease. *Journal of Biological Chemistry* 270, 5331-5338.
- Strongin, A. Y., Marmer, B. L., Grant, G. A., and Goldberg, G. I. (1993). Plasma membrane-dependent activation of the 72-kDa type IV collagenase is prevented by complex formation with TIMP-2. *J Biol Chem* 268, 14033-9.
- Suzuki, N., Labosky, P. A., Furuta, Y., Hargett, L., Dunn, R., Fogo, A. B., Takahara, K., Peters, D. M., Greenspan, D. S., and Hogan, B. L. (1996). Failure of ventral body wall closure in mouse embryos lacking a procollagen C-proteinase encoded by *Bmp1*, a mammalian gene related to *Drosophila* *tolloid*. *Development* 122, 3587-95.
- Suzuki, Y., Someki, I., Adachi, E., Irie, S., and Hattori, S. (1999). Interaction of collagen molecules from the aspect of fibril formation: acid-soluble, alkali-

- treated, and MMP1-digested fragments of type I collagen. J Biochem (Tokyo) 126, 54-67.
- Takino, T., Sato, H., Shinagawa, A., and Seiki, M. (1995). Identification of the second membrane-type matrix metalloproteinase (MT-MMP-2) gene from a human placenta cDNA library. MT-MMPs form a unique membrane-type subclass in the MMP family. J Biol Chem 270, 23013-20.
- Templeton, N. S., and Stetler-Stevenson, W. G. (1991). Identification of a basal promoter for the human Mr 72,000 type IV collagenase gene and enhanced expression in a highly metastatic cell line. Cancer Res 51, 6190-3.
- Teti, A., Farina, A. R., Villanova, I., Tiberio, A., Tacconelli, A., Sciortino, G., Chambers, A. F., Gulino, A., and Mackay, A. R. (1998). Activation of MMP-2 by human GCT23 giant cell tumour cells induced by osteopontin, bone sialoprotein and GRGDSP peptides is RGD and cell shape change dependent. Int J Cancer 77, 82-93.
- Thompson, E. W., Yu, M., Bueno, J., Jin, L., Maiti, S. N., Palao-Marco, F. L., Pulyaeva, H., Tamborlane, J. W., Tirgari, R., and Wapnir, I. (1994). Collagen induced MMP-2 activation in human breast cancer. Breast Cancer Res.Treat. 31, 357-370.
- Tomasek, J. J., Halliday, N. L., Updike, D. L., Ahern-Moore, J. S., Vu, T. K., Liu, R. W., and Howard, E. W. (1997). Gelatinase A activation is regulated by the organization of the polymerized actin cytoskeleton. J Biol Chem 272, 7482-7.
- Trelstad, R. L., and Hayashi, K. (1979). Tendon collagen fibrillogenesis: intracellular subassemblies and cell surface changes associated with fibril growth. Dev Biol 71, 228-42.

- Tryggvason, K., Hoyhtya, M., and Salo, T. (1987). Proteolytic degradation of extracellular matrix in tumor invasion. Biochim Biophys Acta 907, 191-217.
- Tuderman, L., and Prockop, D. J. (1982). Procollagen N-proteinase. Properties of the enzyme purified from chick embryo tendons. Eur J Biochem 125, 545-9.
- Ueno, H., Nakamura, H., Inoue, M., Imai, K., Noguchi, M., Sato, H., Seiki, M., and Okada, Y. (1997). Expression and tissue localization of membrane-types 1, 2, and 3 matrix metalloproteinases in human invasive breast carcinomas. Cancer Res 57, 2055-60.
- van der Rest, M., Aubert-Foucher, E., Dublet, B., Eichenberger, D., Font, B., and Goldschmidt, D. (1991). Structure and function of the fibril-associated collagens. Biochem Soc Trans 19, 820-4.
- van der Rest, M., and Garrone, R. (1991). Collagen family of proteins. Faseb J 5, 2814-23.
- Van Wart, H. E., and Birkedal-Hansen, H. (1990). The cysteine switch: a principle of regulation of metalloproteinase activity with potential applicability to the entire matrix metalloproteinase gene family. Proc Natl Acad Sci U S A 87, 5578-82.
- Velasco, G., Cal, S., Merlos-Suarez, A., Ferrando, A. A., Alvarez, S., Nakano, A., Arribas, J., and Lopez-Otin, C. (2000). Human MT6-matrix metalloproteinase: identification, progelatinase A activation, and expression in brain tumors. Cancer Res 60, 877-82.
- Vogel, W. (1999). Discoidin domain receptors: structural relations and functional implications. Faseb J 13, S77-82.
- Vogel, W., Gish, G. D., Alves, F., and Pawson, T. (1997). The discoidin domain receptor tyrosine kinases are activated by collagen. Mol Cell 1, 13-23.

- Wada, K., Sato, H., Kinoh, H., Kajita, M., Yamamoto, H., and Seiki, M. (1998). Cloning of three *Caenorhabditis elegans* genes potentially encoding novel matrix metalloproteinases. Gene 211, 57-62.
- Wang, Z., Juttermann, R., and Soloway, P. D. (2000). TIMP-2 is required for efficient activation of proMMP-2 in vivo [In Process Citation]. J Biol Chem 275, 26411-5.
- Wapnir, I. L., Southard, H., Chen, G., Friedman, J., Boyd, C. D., and Amenta, P. S. (1996). Collagen gene expression in the neomatrix of carcinoma of the breast. Invasion Metastasis 16, 308-16.
- Will, H., Atkinson, S. J., Butler, G. S., Smith, B., and Murphy, G. (1996). The soluble catalytic domain of membrane type 1 matrix metalloproteinase cleaves the propeptide of progelatinase A and initiates autoproteolytic activation. Regulation by TIMP-2 and TIMP-3. 271, 17119-17123.
- Will, H., and Hinzmann, B. (1995). cDNA sequence and mRNA tissue distribution of a novel human matrix metalloproteinase with a potential transmembrane segment. Eur J Biochem 231, 602-8.
- Woessner, J. F., Jr. (1994). The family of matrix metalloproteinases. Ann N Y Acad Sci 732, 11-21.
- Wu, H., Byrne, M. H., Stacey, A., Goldring, M. B., Birkhead, J. R., Jaenisch, R., and Krane, S. M. (1990). Generation of collagenase-resistant collagen by site-directed mutagenesis of murine pro alpha 1(I) collagen gene. Proc Natl Acad Sci U S A 87, 5888-92.
- Yamagata, S., Yoshii, Y., Suh, J. G., Tanaka, R., and Shimizu, S. (1991). Occurrence of an active form of gelatinase in human gastric and colorectal carcinoma tissues. Cancer Lett 59, 51-5.

- Yamamoto, M., Sumiyoshi, H., Nakagami, K., Taniyama, K., and Tahara, E. (1984). Distribution of collagen types I and III and basal lamina in human gastric carcinoma: an immunohistochemical and electron microscopic study. Virchows Arch A Pathol Anat Histopathol 403, 313-22.
- Yoshiyama, Y., Sato, H., Seiki, M., Shinagawa, A., Takahashi, M., and Yamada, T. (1998). Expression of the membrane-type 3 matrix metalloproteinase (MT3-MMP) in human brain tissues. Acta Neuropathol (Berl) 96, 347-50.
- Yu, A. E., Hewitt, R. E., Kleiner, D. E., and Stetler-Stevenson, W. G. (1996). Molecular regulation of cellular invasion--role of gelatinase A and TIMP-2. Biochem Cell Biol 74, 823-31.
- Yu, M., Sato, H., Seiki, M., Spiegel, S., and Thompson, E. W. (1997). Calcium influx inhibits MT1-MMP processing and blocks MMP-2 activation. FEBS Lett 412, 568-72.
- Yu, M., Sato, H., Seiki, M., and Thompson, E. W. (1995). Complex regulation of membrane-type matrix metalloproteinase expression and matrix metalloproteinase-2 activation by concanavalin A in MDA-MB-231 human breast cancer cells. Cancer Res. 55, 3272-3277.
- Yurchenco, P. D., and Schittny, J. C. (1990). Molecular architecture of basement membranes. Faseb J 4, 1577-90.
- Zhou, Z., Apte, S. S., Soininen, R., Cao, R., Baaklini, G. Y., Rauser, R. W., Wang, J., Cao, Y., and Tryggvason, K. (2000). Impaired endochondral ossification and angiogenesis in mice deficient in membrane-type matrix metalloproteinase I. Proc Natl Acad Sci U S A 97, 4052-7.
- Zucker, S., Conner, C., DiMassmo, B. I., Ende, H., Drews, M., Seiki, M., and Bahou, W. F. (1995). Thrombin induces the activation of progelatinase A in

vascular endothelial cells. Physiologic regulation of angiogenesis. J Biol Chem 270, 23730-8.

Zucker, S., Drews, M., Conner, C., Foda, H. D., DeClerck, Y. A., Langley, K. E., Bahou, W. F., Docherty, A. J., and Cao, J. (1998). Tissue inhibitor of metalloproteinase-2 (TIMP-2) binds to the catalytic domain of the cell surface receptor, membrane type 1-matrix metalloproteinase 1 (MT1-MMP). J Biol Chem 273, 1216-22.

VITA

Miss Neeracha Ruangpanit was born on November 21, 1971 in Bangkok, Thailand. She received the degree of Doctor of Dental Surgery from Chulalongkorn University in 1994. After working as a lecturer in the Department of Anatomy, Faculty of Dentistry, Chulalongkorn University from June 1994 - June 1996, she started her graduate study for the Degree of Doctor of Philosophy in Oral Biology, Faculty of Dentistry, Chulalongkorn University. The research component of this degree was performed at St. Vincent's Institute of Medical Research, Melbourne, Australia, in the laboratory of Associate Professor Dr. Erik W. Thompson. At present she continues her work in the Department of Anatomy, Faculty of Dentistry, Chulalongkorn University, Bangkok, Thailand.