การดัดแปรผ้าฝ้ายด้วยแคตไอออนิกรีแอคทีฟพอลิเมอร์ในการฟอกเพื่อเพิ่มการรับสีย้อมรีแอคทีฟ

นางสาวพัชรี่ ลาภสุริยกุล



วิทยานิพนธ์นี้เป็นส่วนหนึ่งของการศึกษาตามหลักสูตรปริญญาวิทยาศาสตรมหาบัณฑิต สาขาวิชาวิทยาศาสตร์พอลิเมอร์ประยุกต์และเทคโนโลยีสิ่งทอ ภาควิชาวัสดุศาสตร์ คณะวิทยาศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย ปีการศึกษา 2543 ISBN 974-13-0462-5 ลิขสิทธิ์ของจุฬาลงกรณ์มหาวิทยาลัย

MODIFICATION OF COTTON FABRIC WITH A CATIONIC REACTIVE POLYMER IN BLEACHING TO IMPROVE REACTIVE DYEABILITY

Miss Patcharee Larpsuriyakul

A Thesis Submitted in Partial Fulfillment of the Requirements

for the Degree of Master of Science in Applied Polymer Science and Textile Technology

Department of Materials Science

Faculty of Science

Chulalongkorn University

Academic Year 2000

ISBN 974-13-0462-5

Thesis Title	Modification of Cotton Fabric with a Cationic Reactive Polymer in
	Bleaching to Improve Reactive Dyeability
Ву	Miss Patcharee Larpsuriyakul
Field of Study	Applied Polymer Science and Textile Technology
Thesis Advisor	Asst. Prof. Kawee Srikulkit, Ph.D.
Accer	oted by the Faculty of Science, Chulalongkorn University in Partial
Fulfillment of the Req	uirements for the Master 's Degree
	Wand: PLT Dean of Faculty of Science
	(Associate Professor Wanchai Phothiphichitr, Ph.D.)
THESIS COMMITTEE	
	8 Chuay julit Chairman
	(Associate Professor Saowaroj Chuayjuljit)
	K Snkulling Thesis Advisor
	(Assistant Professor Kawee Srikulkit, Ph.D.)
	Payan Santih Member
	(Associate Professor Paiparn Santisuk)
	Us Communication Member
	(Usa Sangwatanaroj, Ph.D)
	Shirtan Kuth'ngovarot Member
	(Siriwan Kittinaowarat, Ph.D.)

นางสาว พัชรี ลาภสุริยกุล : การดัดแปรผ้าฝ้ายด้วยแคตไอออนิกรีแอคทีฟพอลิเมอร์ในการฟอกเพื่อเพิ่มการรับสี ย้อมรีแอคทีฟ (MODIFICATION OF COTTON FABRIC WITH A CATIONIC REACTIVE POLYMER IN BLEACHING TO IMPROVE REACTIVE DYEABILITY) อ. ที่ปรึกษา : ผศ.ดร. กาวี ศรีกูลกิจ, 63 หน้า. ISBN 974-13-0462-5.

ผ้าฝ้ายที่ผ่านการกำจัดใชมันแล้วถูกนำมาใช้เพื่อศึกษาเกี่ยวกับการดัดแปรเล้นใยเซลลูโลสด้วยวิธีทาง เคมี เพื่อทำการย้อมผ้าฝ้ายด้วยสีรีแอคทีฟในภาวะไร้เกลือ โดยการเติม MAPTAC ซึ่งเป็นสารมอนอเมอร์ที่มีประจุ บวกเพื่อให้เกิดปฏิกิริยาพอลิเมอร์ไรเซชัน เข้าไปในโมเลกุลของเซลลูโลส ในระหว่างกระบวนการฟอกผ้า ซึ่งไม่เป็น การเพิ่มขั้นตอนให้กับกระบวนการผลิตเดิม หลังจากการดัดแปร ผ้าฝ้ายจะถูกนำไปวัดปริมาณของ MAPTAC ที่ สามารถยึดติดอยู่บนเส้นใยเซลลูโลส ซึ่งจะใช้วิธีการวิเคราะห์ธาตุในโตรเจนโดยอาศัยเครื่องมือวิเคราะห์ธาตุ และ นำไปย้อมสีรีแอคทีฟด้วยวิธีการย้อมแบบแข่ (exhaustion dyeing) โดยไม่ต้องใช้เกลือเป็นสารช่วยย้อม

ผลการทดลองแสดงให้เห็นว่า ประสิทธิภาพการฟอกชาวของสารไฮโดรเจนเปอร์ออกไซด์ ในขณะที่มี
MAPTAC อยู่ด้วย จะลดลงเล็กน้อย ส่วนการย้อมผ้าฝ้ายดัดแปรด้วยสีรีแอคทีฟในภาวะไร้เกลือ ทำให้ผ้าฝ้ายมี
การติดสีสูงขึ้นตามปริมาณ MAPTAC ที่ใช้และสูงกว่าผ้าฝ้ายที่ไม่ได้ผ่านการดัดแปร ทั้งนี้เนื่องจากประจุบวกที่อยู่
บน MAPTAC ซึ่งจะดึงคูดประจุลบของสีรีแอคทีฟได้ดี และจากผลการย้อมทำให้สรุปได้ว่า สมบัติการย้อมผ้าฝ้าย
ที่ผ่านการดัดแปรจะขึ้นกับประสิทธิภาพในการยึดติดของ MAPTAC บนเส้นใยเซลลูโลสในขั้นตอนการดัดแปร

ภาควิชาวัสดุศาสตร์ สาขาวิชาวิทยาศาสตร์พอลิเมอร์ประยุกต์ ฯ ปีการศึกษา2543 ลายมือชื่อนิสิต ทั่งรักษา My พาวิกา

: MAJOR APPLIED POLYMER SCIENCE AND TEXTILE TECHNOLOGYMAJOR # # 4272348623

KEY WORD: COTTON FABRIC / DYEABILITY MODIFICATION / BLEACHING / SINGLE-BATH

PATCHAREE LARPSURIYAKUL: MODIFICATION OF COTTON FABRIC WITH A CATIONIC

REACTIVE POLYMER IN BLEACHING TO IMPROVE REACTIVE DYEABILITY. THESIS

ADVISOR: ASST.PROF.KAWEE SRIKULKIT, Ph.D., 63 PAGES, ISBN 974-13-0462-5.

The modification of scoured cotton fabric by chemical treatment was studied. The study was

aimed to dye cotton fabric in the absence of salt. The principle method of modification was to graft

polymerization of the cationic based vinyl monomer. [3-(Methacryloylamino)

propyl]trimethylammonium chloride (MAPTAC), onto scoured cotton during bleaching process,

aiming at modifying dyeability of the fiber using single-bath. An elemental analyzer was employed to

measure the extent of MAPTAC fixation on cellulose. Modified cotton fabric was dyed with commercial

reactive dyes by exhaustion method without the requirement of salt.

The results showed that the ability of bleaching performance of H₂O₂ in the presence of the

modifying agent was slightly reduced. The dye uptake and color strength of dyed modified fabric was

markedly increased with an increase in the concentration of MAPTAC. In addition, those properties

obtained from modified cotton fabric were higher than those obtained from the dyeing of unmodified

cotton fabric. This was attributed to the presence of the cationic groups of the MAPTAC that played a

crucial role in attracting the anionic dyes from the dyebath. The finding results tended to suggest that

the dyeing properties of modified fabric were exactly dependent on the efficiency of MAPTAC fixation

on cellulose during concurrent modifying and bleaching of cotton.

Department Materials Science

Field of study Applied polymer science

Academic year 2000

Advisor's signature

Co-advisor's signature

Student's signature R. Langsoniyahol.

K. Sikuliya L.



Acknowledgement

I wish to express my deep gratitude to Asst. Prof. Dr. Kawee Srikulkit, my advisor, for his valuable guidance, advice and encouragement throughout this research work.

I am grateful to Assoc. Prof. Paipam Santisuk and Dr. Usa Sangwatanaroj for their comments and assistance for my study and the Department of Materials Science, Chulalongkom University for the graduate course.

I wish to thank my lecturers and friends, especially to Miss Pulsiri Rattananiyomkul for her kindly assistance, and Miss Nattaya Punrattanasin for her helpful encouragement.

My thanks are also extended to the administration staff of the Materials Science Department for their assistance.

Finally, I would like to express my deepest appreciation to my family for their love, encouragement, and worthy moral and financial support throughout my study at Chulalongkom University.

CONTENTS

	P	age
Abstract (Thai)	iv
Abstract (English)	٧
Acknowle	dgement	vi
List of Tab	ples	X
List of Fig	ures	хi
List of Sch	nemes	xiii
Chapter		
I	Introduction	
H	Literature Survey	3
	Basic Information about Cellulose	3
	Cotton	3
	Cotton Fiber Composition	6
	Properties of Cotton Fiber	12
	Structure of Cotton in Relation to Dyeing	17
	Bleaching of Cotton with Hydrogen peroxide	17
	Dyeing of Cotton with Reactive Dyes	19
	Problems of Reactive Dyeing	22
	Previous Research Works on Chemical Modification of Cellulose	
	Fibers to Improve Dyeability	23
	Concept of Modification and Bleaching of Cotton Fabric in	
	Single-bath Process	30
111	Experimental	33
	Materials	33
	Chemicals and Dyes	33
	Equipment	33
	Method of the Modification and Bleaching of Cotton Fabric in	
	Single-bath Process	34

Contents (continued)		Page	
	General Dyeing Procedures		
	Dyeing of Modified Fabric Using Exhaustion Method	35	
	Dyeing of Unmodified Cotton Fabric by Conventional		
	Exhaustion Method	36	
	Evaluation of Whiteness Property of Modified Bleached Cotton		
	Fabric	. 37	
	Measurement of Dye Exhaustion	. 38	
	Measurement of Dye Fixation	. 39	
	Total Nitrogen Content Determination	. 39	
	Image Microscopic Analysis	40	
	Measurement of Colorfastness	. 40	
IV	Results and Discussion	42	
	Effect of the Modifying Agent Concentration on Whiteness		
	Property	42	
	Total Nitrogen Content Determination	43	
	Effect of Increasing Concentrations of K ₂ S ₂ O ₈ on Dye Uptake and		
	Dye Fixation	45	
	Effect of Liquor Ratio on Dye Uptake and Dye Fixation	46	
	Effect of Temperature on Dye Uptake and Dye Fixation	47	
	Effect of Increasing Concentration of Modifying Agent on		
	Dye Uptake and Dye Fixation	48	
	Effect of Different Dye Types on Dye Uptake and Color Yield	51	
	Effect of Increasing Dye Concentration on Dye Uptake and		
	Color Strength	52	
	Microscopic Analysis of Cross-Section of Dyed Yarn	53	
	Evaluation of Color Fastness to Light	54	
	Dyeing of Unmodified Cotton Fabrics	55	
V	Conclusions	57	
VI	Recommendations	58	

		Page
17	References	. 59
	Appendix	. 61
	Biography	. 63

List of Tables

Tables	F	age)
2-1	Cellulose content in plant material	. 3
2-2	Composition of the typical cotton	. 11
2-3	Properties of cotton fiber	12
3-1	Chemicals used in this project	33
4-1	Whiteness index of modified bleached cotton fabric at various	42
	modifying concentrations	
4-2	Total nitrogen content of MAPTAC cotton before and after H ₂ O ₂ /NaOH	. 43
4-3	Color yield and % dye exhaustion of modified fabrics at various	45
	concentrations of K ₂ S ₂ O ₈ and liquor ratios	
4-4	Color strength and % dye exhaustion of modified fabrics at various	47
	temperature	
4-5	Color yield, % dye exhaustion and % fixation of modified fabrics at	49
	various modifying concentrations	
4-6	Color yield, % dye exhaustion and % fixation of modified dyed fabrics	51
	at different dye types	
4-7	Effect of increasing dye concentration on dyeing properties of	52
	modified fabric	
4-8	Light fastness of dyed cotton fabrics modified with various	55
	concentrations of modifying agent	
4-9	Color yield and % dye exhaustion of unmodified fabrics	56
	dyed with various concentrations of electrolyte	
4-10	Light fastness of unmodified cotton fabrics dyed with various	56
	concentrations of dyes	

List of Figures

Figure:	S	Page
2.1	Scanning electron micrographs of raw cotton fibers	. 4
2.2	Optical micrographs of raw cotton fibers x 184	. 4
2.3	Bilateral structure of mature cotton	. 5
2.4	Hierarchical organization of cellulose	. 6
2.5	Representation of cellulose	. 7
2.6	Cross-section of unit cell in the ab plane	8
2.7	Unit cell in the ac plane	8
2.8	Hydrogen bonding in cellulose	9
2.9	Diagram of fine structure in cellulose	10
2.10	Idealized diagram of cotton morphology	. 10
2.11	Characteristic structural features of a reactive dye	. 19
2.12	Dichlorotriazinyl reactive dye and Monochlorotriazinyl reactive dye	. 20
2.13	Monofluorotriazinyl dyes	21
2.14	The specific base catalyzed addition of the nucleophilic	22
	functional group HY of the textile fiber	
2.15	Figure 2.15	. 27
3.1	The treatment profile of the exhaustion method	. 34
3.2	The dyeing profile of modified fabric using exhaustion method	35
3.3	The dyeing profile of unmodified fabric using exhaustion method	. 37
3.4	Macbeth spectrophotometer	. 38
3.5	UV/Vis spectrophotometer	. 38
3.6	Optical microscope	. 40
3.7	Xenon weather meter	. 41
4.1	Whiteness index chart of modified bleached cotton fabric	. 42
4.2	Dependence of % dye exhaustion on concentrations of K2S2O8 and	45
	Liquor ratios	
4.3	Dependence of color yield on concentrations of $K_2S_2O_8$ and liquor ratios	. 46
4.4	Dependence of dye uptake on the temperature	. 47

Figures (continued)		Page
4.5	Dependence of color strength on the temperature	48
4.6	Dependence of dye uptake on the concentrations of MAPTAC	. 49
4.7	Dependence of % dye fixation on the concentration of MAPTAC	. 50
4.8	Optical micrograph of MAPTAC treated yarn dyed with	. 53
	2% Procion Crimson CX-B	

List of Schemes

Schem	es	Page
	Scheme 2.1	. 7
	Scheme 2.2.	. 17
	Scheme 2.3	. 18
	Scheme 2.4	. 18
	Scheme 2.5	. 18
	Scheme 2.6	. 20
	Scheme 2.7	. 21
	Scheme 2.8	. 21
	Scheme 2.9	. 24
	Scheme 2.10	24
	Scheme 2.11	24
	Scheme 2.12	25
	Scheme 2.13	25
	Scheme 2.14	26
	Scheme 2.15	27
	Scheme 2.16	28
	Scheme 2.17	29
	Scheme 2.18	30
	Scheme 2.19	30
	Scheme 2.20.	31
	Scheme 2.21	32