

REFERENCES

- Alam, M.S., Qasim, M.A., Kamil, M. and Ilyas, M. 1986. Sorbifolin 6-galactoside from *Garcinia andamanica*. *Phytochemistry* 25 (12) : 2900-2901.
- _____, Kamil, M. and Ilyas, M. 1987. 4'-Hydroxywogonin-7-neohesperidoside from *Garcinia andamanica*. *Phytochemistry* 26(6) : 1843-1844.
- Ampofo, S.A. and Waterman, P.G. 1986. Xanthones from three *Garcinia* species. *Phytochemistry* 25(10) : 2351-2355.
- Asai, F., Tosa, H., Tanaka, T. and Iinuma, M. 1995. A xanthone from pericarps of *Garcinia mangostana*. *Phytochemistry* 39(4) : 943-944.
- Asano, J., Chiba, K., Tada, M. and Yoshii, T. 1996. Cytotoxic xanthones from *Garcinia hanburyi*. *Phytochemistry* 41(3) : 815-820.
- Babu, V., Ali, S.M., Sultana, S. and Ilyas, M. 1988. A biflavonoid from *Garcinia nervosa* *Phytcchemistry* 27(10) : 3332-3335.
- Balasubramanian, K. and Rajagopalan, K. 1988. Novel xanthones from *Garcinia mangostana*, structures of BR-xanthone-A and BR-xanthone-B. *Phytochemistry* 27(5) : 1552-1554.
- Bandaranayake, W.M., Selliah, S.S., Sultanbawa, M.U.S. and Ollis, W.D. 1975. Biflavonoids and xanthones of *Garcinia terpnophylla* and *G. echinocarpa*. *Phytochemistry* 14 : 1878-1880.
- Budzikiewicz, H., Djerassi, C. and Williams, D.H. 1964. Structure Elucidation of Natural Products By Mass Spectrometry Vol. II Holden-Day, Inc., San Francisco, pp. 94-108.
- Bennett, G.J. and Lee, H.H. 1989. Xanthones from Gutiferae. *Phytochemistry* 28 (4) : 967-998.
- Burkill, I. H. 1935. A Dictionary of Economic Products of the Malay Peninsula. The Crown Agents for the Colonies, London., pp. 1046-1048.

- Carpenter, I., Locksley, H.D. and Scheinmann, F. 1969. Xanthones in higher plants : Biogenetic proposals and a chemotaxonomic survey. *Phytochemistry* 8 : 2013-2026.
- Chen, F.C., Lin, Y.M. and Hung, J.C. 1975. A new biflavanone glucoside from *Garcinia multiflora*. *Phytochemistry* 14 : 818-820.
- _____, Lin, Y.M. and Hung, J.C. 1975. Phenolic compounds from the heartwood of *Garcinia multiflora*. *Phytochemistry* 14 : 300-303.
- Cotterill, P.J. and Sheinmann, F. 1978. Extractives from Guttiferae Part 34. Kolaflavanone, a new biflavanone from the nuts of *Garcinia kola* Heckel. Applications of ^{13}C nuclear magnetic resonance in elucidation of the structures of flavonoids. *J. Chem. Soc. (Perkin I)* 532-538.
- _____, Scheinmann, F. and Puranik, G.S. 1977. Phenolic compounds from the heartwood of *Garcinia indica*. *Phytochemistry* 16 : 148-149.
- Crichton, E.G. and Waterman, P.G. 1979. Manniflavanone, a new 3,8-linked flavanone dimer from the stem bark of *Garcinia mannii*. *Phytochemistry* 18 : 1553-1557.
- Djerassi, C. 1969. Applications of Mass Spectrometry in the Steroid Field in International Symposium on Chemistry of Natural Products (Mexican Chemical Society) Butterworths, London, pp. 205-215.
- Elderfield, R.C., 1950. Heterocyclic Compounds. John Wiley & sons, Inc.,London. pp.419-444.
- Fukuyama, Y., Kamiyama, A., Mima, Y. and Kodama, M. 1991. Prenylated xanthones from *Garcinia subelliptica*. *Phytochemistry* 30(10) : 3433-3436.
- _____, Kaneshi, A., Tani, N. and Kodama, M. 1993. Subellinone, a polyisoprenylated phloroglucinol derivative from *Garcinia subelliptica*. *Phytochemistry* 33(2) : 483-485.

- Goh, S.H., Jantan, I., Gray, A.I. and Waterman, P.G. 1992. Prenylated xanthones from *Garcinia opaca*. Phytochemistry 31 (4) : 1383-1386.
- Govindachari, T.R., Kalyanaraman, P.S., Muthukumaraswamy, N. and Pai, B.R. 1971. Xanthones of *Garcinia mangostana* Linn. Tetrahedron 27: 3919-3926.
- _____, de Silva, A.M.Y.J., Sotheeswaran, S., Balasubramaniam, S. and Wazeer, M.I.M. 1984. Terpenoid and biflavonoid constituents of *Calophyllum calaba* and *Garcinia spicata* from Sri Lanka. Phytochemistry 23(2) : 323-328.
- _____, Sriyani, H.T.B. and Sotheeswaran, S. 1984. Quaesitol, a phenol from *Garcinia quaesita*. Phytochemistry 23(11) ; 2679-2681.
- _____, Sriyani, H.T.B., Sotheeswaran, S. and Waight, E.S. 1983. 2,5-Dihydroxy-1,6-dimethoxyxanthone and biflavonoids of *Garcinia thwaitesii*. Phytochemistry 22(1) : 233-235.
- Gunatilaka, A.A.L., Sotheeswaran, S., Sriyani, H.T.B. and Waight, E.S. 1982. A revised structure for hermonionic acid. Tetrahedron Lett. 23 (29) : 2987-2990.
- Gustafson, K.R., et al. 1992. The Guttiferones, HIV-inhibitory benzophenones from *Sympodia globulifera*, *Garcinia livingstonei*, *Garcinia ovalifolia* and *Clusia rosea*. Tetrahedron 48(46) : 10093-10102.
- Harrison, L.J., Leong, L.S., Sia, G.L., Sim, K.Y. and Tan, H.T.W. 1993. Xanthones from *Garcinia forbesii*. Phytochemistry 33(3) : 727-728.
- Herbin , G.A. , Jackson, B., Locksley, H.D. , Scheinmann, F . and Wolstenholme, W.A. 1970. Phytochemistry 9 : 221-226.
- Holloway, D.M. and Scheinmann, F. 1975. Phenolic compounds from the heartwood of *Garcinia mangostana*. Phytochemistry 14 : 2517-1518.
- Hooker, J.D, et al. 1872. Flora of British India. Jayyed Press, Delhi., pp. 258-270

- Hussain, R.A., Owegby, A.G., Parimoo, P. and Waterman, P.G. 1982. Kolanone, a novel polyisoprenylated benzophenone with antimicrobial properties from the fruit of *Garcinia kola*. Planta Med. 44 : 78-81.
- _____ and Waterman, P.G. 1982. Lactones, flavonoids and benzophenones from *Garcinia conrauana* and *Garcinia mannii*. Phytochemistry 21(6) : 1393-1396.
- Iinuma, M., Tosa, H., Tanaka, T., Asai, F. and Shimano, R. 1995. Three xanthones from root bark of *Garcinia subelliptica*. Phytochemistry 38 (1) : 247-249.
- _____, Tosa, H., Tanaka, T., Asai, F. and Shimano, R. 1995. Two xanthones with a 1,1-dimethylallyl group in root bark of *Garcinia subelliptica*. Phytochemistry 39(4) : 945-947.
- _____, Tosa, H., Tanaka, T., Asai, F. and Shimano, R. 1995. Two new xanthones from the root bark of *Garcinia subelliptica*. Heterocycles 40(1) : 279-284.
- _____, Tosa, H., Tanaka, T., Shimano, R., Asai, F. and Yonemori S. 1994. Two xanthones from root bark of *Garcinia subelliptica*. Phytochemistry 35(5) :1355-1360.
- Iwu, M. and Igboko, O. 1982. Flavonoids of *Garcinia kola* seeds. J. Nat. Prod. 45 : 650-651.
- Jackson, B., Locksley, H.D. and Scheinmann, F. 1969. Extractives from Guttiferae. Part XIII. Isolation and structure of five xanthones from *Garcinia eugeniiifolia* Wall. J. Chem. Soc. (C) 16 : 2201-2203.
- _____, Locksley,H.D., Scheinmann, F. and Wolstenholme, W.A. 1971. Extractives from Guttiferae. part XXII. The isolation and structure of four novel biflavanones from the heartwoods of *Garcinia buchananii* Baker and *G. eugeniiifolia* Wall. J. Chem. Soc. (C) 3791-3804.

- Joshi, B.S. , Kamat, V.N. and Viswanathan, N 1970. The isolation and structure of two biflavonones from *Garcinia talbofi*. Phytochemistry 9 ; 881-888.
- Kabangu, K., Galeffi, C., Aonzo, E., Nicoletti, M. and Messana, I. 1987. A new biflavanone from the bark of *Garcinia kola*. Planta Med. 53 ; 275-277.
- Karanjagaokar, C.G., Radhakrishnan, P.V. and Venkataraman, K. 1967. Morelloflavone, a 3-(8) flavonylflavanone, from the heartwood of *Garcinia morella*. Tetrahedron Lett. 33 : 3195-3198.
- Karanjgaokar , C.G., Nair, P.M. and Venkataraman, K. 1966 Morellic, Isomorellic and Gambogic acid Tetrahedron Lett. 7 : 687-691.
- _____,Rao, A.V.R. , Venkataraman, K. and Yemul , S.S. 1973. Tetrahedron Lett. 50 : 4977-4980.
- Kartha, G., et al. 1963. The constitution of Morellin. Tetrahedron Lett. 7 : 459-472
- Konoshima, M., et al. 1969. The constitution of flavonoid from *Garcinia spicata* Hook .f. Tetrahedron Lett. 2 : 121-124.
- _____, and Ikeshiro, Y. 1970. Fukugiside. The first biflavonoid glycoside from *Garcinia spicata* HooK.f. Tetrahedron Lett. 20 : 1717- 1720.
- _____, Ikeshiro, Y. and Miyahara, S. 1970. The constitution of biflavonoids from *Garcinia* plants. Tetrahedron Lett. 48 : 4203-4206
- Krishnamurthy, N., Lewis, Y.S. and Ravindranath, B. 1981. On the structures of garcinol, isogarcinol and camboginol. Tetrahedron Lett. 22 : 793-796.
- Lee, H.H. and Chan, H.K. 1977. 1,3,6,-Trihydroxy-7-methoxy-8-(3,7-dimethyl-2,6-octadienyl)xanthone from *Garcinia cowa*. Phytochemistry 16 : 2038-2040.
- Lewis, Y.S. and Neelakantan, S. 1965. (-)-Hydroxycitric acid - the principal acid in the fruits of *Garcinia cambogia* DESR. Phytochemistry 4: 619-625.

- Lin, L.J., Lin, L.Z., Pezzuto, J.M., Cordell, G.A. and Ruangrungsi, N. 1993
 Isogambogic acid and isomorellinol from *Garcinia hanburyi*. Magn. Res. Chem. 31(4) : 340-347.
- Locksley , H.D. , Moore, I. and Scheinmann, F. 1967. Extractives from Guttiferae - VI the significance of maclurin in xanthone biosynthesis. Tetrahedron 23 : 2229-2234.
- Mahabusarakam, W., Wiriyachitra, P. and Taylor, W.C. 1987. Chemical constituents of *Garcinia mangostana*. J. Nat. Prod. 50(3) : 474-478.
- Minami, H., et al. 1994. Antioxidant xanthones from *Garcinia subelliptica*. Phytochemistry 36(2) : 501-506.
 _____, Takahashi, E., Fukuyama, Y., Kodama, M., Yoshizawa, T. and Nakagawa, K. 1995. Novel xanthones with superoxide scavenging activity from *Garcinia subelliptica*. Chem. Pharm. Bull. 43(2) : 347-349.
 _____, Takahashi, E., Kodama, M. and Fukuyama, Y. 1996. Three xanthones from *Garcinia subelliptica*. Phytochemistry 41(2) : 629-633.
- Na Pattalung, P., Thongtheeraparp, W., Wiriyachitra, P. and Taylor, W.C. 1994.
 Xanthones of *Garcinia cowa*. Planta Med. 60 : 365-368.
- _____, Wiriyachitra, P. and Ongsakul , M. 1988. The anti-microbial activities of rubraxanthone isolated from *Garcinia parvifolia* J.Sci. Soc Thailand 14 : 67-71
- Nyemba, A.M., Mpando, T.N., Connolly, J.D. and Rycroft, D.S. 1990. Cycloartane derivatives from *Garcinia lucida* Phytochemistry 29(3) : 994-997.
- Parveen, M. and Khan, N.U. 1987. A new isoprenylated xanthone from *Garcinia mangostana* Linn. Chem. Ind. (London) 12 : 418.
 _____and Khan, N.U. 1988. Two xanthones from *Garcinia mangostana* Phytochemistry. 27 (11) : 3694-3696.

- _____, Khan, N.U.D., Achari, B. and Dutta, P.K. 1991. A triterpene from *Garcinia mangostana*. Phytochemistry 30(1) : 361-362.
- Petter A. 1967. On the question of the structures of GB-1, GB - 1a and GB-2 , A new group of bisflavonoids. Tetrahedron Lett. 19 : 1767 - 1771
- _____, Warrew, R., Chexal, K.K., Handa, B.K. and Rahman, W. 1971. Biflavonyls from Guttiferae - *Garcinia livingstonii*. Tetrahedron 27 : 1625-1634.
- _____, 1968. On the structures of GB-1 , GB-1a and GB-2 and on the biosynthesis of Isoflavones. Tetrahedron Lett. 8 : 897-903.
- Perry, L. M. 1980. Medicinal Plants of East and Southeast Asia. MIT Press, Massachusetts., pp. 173-176.
- Rao, A.V.R., Sarma, M.R., Venkataraman, K. and Yemul, S.S. 1974. Benzo-phenone and xanthone with unusual hydroxylation patterns from the heartwood of *Garcinia pedunculata*. Phytochemistry 13 : 1241-1244.
- _____, Venkatswamy, G. and Pendse, A.D. 1980. Camboginol and cambogin. Tetrahedron Lett. 21 : 1975-1978.
- Sahu, A., Das, B. and Chatterjee, A. 1989. Polyisoprenylated benzophenones from *Garcinia pedunculata*. Phytochemistry 24(4) : 1233-1235
- Sakai, S., Katsura, M., Takayama, H., Aimi, N., Chokethaworn, N. and Sutajit, M. 1993. The structure of Garcinone E. Chem. Pharm. Bull. 41(5) : 958-960.
- Sen, A.K., Sarkar, K.K., Mazumder, P.C., Banerji, N., Uusvuori, R. and Hase, T.A. 1980. A xanthone from *Garcinia mangostana*. Phytochemistry 19 : 2223-2225.
- _____, Sarkar, K.K., Mazumder, P.C., Banerji, N., Uusvuori, R. and Hase, T.A. 1982. The structures of Garcinones A, B and C : three new xanthones from *Garcinia mangostana*. Phytochemistry 21(7) : 1747-1750.
- Silverstein, R.M., Bassler, G.C., and Morril, T.C. 1991. Spectrometric identification of organic compounds. 5th ed. New York : John Wiley & Sons, Inc.

- Smithinand, T. 1980. Thai plant name (Botanical names - vernacular names).
2nd ed. Bangkok : Funny Publishing.
- Sordat - Diserens, I., Hamburger, M., Rogers, C. and Hostettmann, K. 1992.
Dimeric xanthones from *Garcinia livingstonei*. Phytochemistry 31(10) : 3589-3593.
- _____, Marston, A., Hamburger, M., Rogers, C. and Hostettmann, K. 1989.
Novel prenylated xanthones from *Garcinia gerrardii* Harvey. Helv. Chem. Acta 72 : 1001-1007.
- _____, Rogers, C., Sordat, B. and Hostettmann, K. 1992. Prenylated xanthones from *Garcinia livingstonei*. Phytochemistry 31(1) : 313-316.
- Sultanbawa, M.U.S. 1980. Xanthonoids of Tropical plants. Tetrahedron 36 : 1465-1506.
- Waterman P.G. and Crichton, E.G. 1980. Xanthones and biflavonoids from *Garcinia densivenia* stem bark. Phytochemistry 19: 2723-2726.
- _____, and Crichton, E.G. 1980. Xanthones, benzophenones and triterpenes from the stem bark of *Garcinia ovalifolia*. Planta Med. 40 : 351-355.
- _____, and Crichton, E.G. 1980. Pyrones from the bark of *Garcinia conrauana* : conrauanalactone, a novel C₂₀ derived α-pyrone. Phytochemistry 19 : 1187-1189.
- _____, and Hussain, R.A. 1982. Major xanthones from *Garcinia quadrifaria* and *Garcinia staudtii* stem bark. Phytochemistry 21(8) : 2099-2101.
- Wright, J.L.C., et al. 1978. Identification of C-24 alkyl epimers of marine sterols by ¹³C-nuclear magnetic resonance spectroscopy. Can. J. Chem. 56 : 1898-1903.

APPENDIX




Figure 1 The EIMS spectrum of GA-1




Figure 2 The IR spectrum of GA-1 (KBr disc)




Figure 3 The 500 MHz $^1\text{H-NMR}$ spectrum of GA-A1 (in CDCl_3)




Figure 4 The expansion of $^1\text{H-NMR}$ spectrum of GA-1 (in CDCl_3)




Figure 5 The expansion of $^1\text{H-NMR}$ spectrum of GA-1 (in CDCl_3)




Figure 6 The 125 MHz ^{13}C -NMR spectrum of GA-1 (in CDCl_3)




Figure 7 The expansion of ^{13}C -NMR spectrum of GA-1 (in $(\text{CD})_3\text{Cl}$)




Figure 8 The expansion of ^{13}C -NMR spectrum of GA-1 (in CDCl_3)




Figure 9 The EIMS spectrum of GA-2




Figure 10 The IR spectrum of GA-2 (KBr disc)




Figure 11 The UV spectrum of GA-2 (in methanol)




Figure 12 The 600 MHz ^1H -NMR spectrum of GA-2 (in DMSO- d_6)




Figure 13 The expansion of ^1H NMR spectrum of GA-2 (in DMSO-d₆)




Figure 14 The 150 MHz ^{13}C -NMR spectrum of GA-2 (in $\text{DMSO}-d_6$)




Figure 15 The DEPT spectrum of GA-2 (in DMSO- d_6)




Figure 16 The HMQC spectrum of GA-2 (in $\text{DMSO}-\text{d}_6$)




Figure 17 The expansion of HMBC spectrum of GA-2 (in DMSO-d_6)




Figure 18 The HMQC spectrum of GA-2 (in DMSO-d₆)

VITA

Miss Jongdee Kosin was born on January, 9, 1968 in Prachinburi, Thailand. She received her Bachelor of Science (in Pharmacy) in 1993 from the Faculty of Pharmaceutical Sciences, Khon Kaen University, Thailand. At present, she is appointed as a hospital pharmacist at Samutprakan Hospital.