

BIBLIOGRAPHY

ธัญโญ เพ็ญชาติ, ประยูร ตีมา, มุกดา ตฤณานันท์ และ อสัน อิษัวล "อภิปรายเรื่องพิช" วารสารกรมการแพทย์และอนามัย ปีที่ 2 ฉบับที่ 3 (2517) : 303-311, 329-330.

Ahuja, M.L., Brooks, A.G., Veeraraghavan, N. and Menon, I.G.K.
"A Note on the Action of Heparin on Russell's Viper Venom"
Indian J. Med. Res. 32(2), (1946) : 317-322.

Arthur, R.C., Harris, P.E., Hurst and Saker, B.M. "Renal Failure After Snake Bite" Med. J. Aust. 2(11), (1967) : 409-411.

Aung - Khin, M. "Histological and Ultrastructural Change of Kidney in Renal Failure After Viper Envenomation" Toxicon 16 (1978) : 71-75.

Bell, R.D., Mandal, A.K. and Paker, D.E. "The Effect of Splenectomy on Renal Function in Epinephrine Induced Renal Failure"
Proc. Soc. Exp. Biol. Med. 167(1981) : 12-14.

Bhat, R.N. "Viperine Snake Bite Poisoning in Jammu" J. Indian M.A. 63(12), (1974) : 383-392.

Blair - West, J.R., Coghlan , J.P., Denton, D.A., Goding J.R., Munro, J.A., Peterson, R.E. and Wintour M. "Humoral Stimulation of Adrenal cortical secretion" J. Clin. Invest. 41(1962) : 1606-1627.

Bruninga, G.L. "Complement : A Review of the Chemistry and Reduction Mechanism" Am. J. Clin. Path. 55(1971) : 273-282.

Chaiyabutr, N., Faulkner, A. and Peaker, M. "Effects of Starvation on the Cardiovascular System, Water Balance and Milk Secretion in Lactating Goat" Res. Vet. Sci. 28(1980) : 291-295.

Chopra, R.N. and Chowhan, J.S. "Action of the Indian Daboia (Vipera russellii) Venom on the Circulatory System" Indian J. Med. Res. 21(3), (1934) : 493-506.

Chugh, K.S., Aikat, B.K., Sharma, B.K., Dash, S.C., Mathew, M.T. and Das, K.C. "Acute Renal Failure Following Snakebite" Am. J. Trop. Med. 24(24), (1975) : 692-697.

Chugh, K.S., Aikat, B.K., Sharma, B.K., Dash, S.C., Mathew, M.T. and Das, K.C. "Renal Lesion Following Envenomation of Russell's Viper" J. Med. Ass. Thailand 61 (1978) : 78.

Culp, J.R., Erdos, E.G., Hinshaw, L.B. and Holmes, D.D. "Effects of Anti-inflammatory Drugs in Shock Caused by Infection of Living E. coli Cell" Proc. Soc. Exp. Biol. Med. 137(1971) : 219-222.

Dunn, M.J. and Hood, V.L. "Prostaglandins and the Kidney" Am. J. Physiol 233(3), (1977) : F 169-F 184.

Dunn, M.J. and Zambraski, E.J. "Renal Effects of Drug that Inhibit PG Synthesis" Kidney int. 18(1980) : 609-622.

Erdos, E.G., Hinshaw, L.B. and Gill, C.C. "Effect of Indomethacin in Endotoxin Shock in Dog" Proc. Soc. Exp. Biol. Med. 125(1967) : 916-919.

Feigen, L.P., Klainer, E., Chapnick, B.M. and Kadowitz, P.J. "The Effect of Indomethacin on Renal Function in Pentobarbital Anesthetized Dog" J. Pharmacol Exp. Ther. 198(2), (1976) : 457-463.

Ffoulkes - Crabbe, D.J.O., Creighton, R.E., Volgyesi, G.A., Steward, D.J. and Nisbet, H.I.A. "The Effect of Splenectomy on Circulatory Adjustments to Hypoxemia in the Anesthetized Dog" Br. J. Anaesth. 48(1976) : 639-641.

Frolich, J.C., Hollifield, J.W., Dormois, J.C., Frolich, B.L., Seyberth, H., Michelakis, A.M. and Oates, J.A. "Suppression of Plasma Renin Activity by Indomethacin in Man" Circ. Res. 39(1976) : 447-452.

Ganguly, S.N. and Malkana, M.T. "Daboia Venom : Its Chemical Composition, Protein Fraction and Their Physiological Action" Indian J. Med. Res. 23(1), (1935) : 131-140.

Gerber, J.G., Olson, R.D. and Nies, A.S. "Interrelationship between Prostaglandins and Renin Release" Kidney Int. 19(1981) : 816-821.

Guyton, A.C. The Adrenocortical hormones in Textbook of Medical Physiology 6th ed. pp. 944-947, W.B. Saunders company, Tokyo, 1981.

Hadler, W.A. and Brazil, O.V. "Pharmacology of Crystalline Crotoxin. IV Nephrotoxicity" Mem. Inst. Butantan. 33(1966) : 1001-1008.

Hall, J.E., Guyton, A.G., Jackson, T.E., Coleman, T.G., Lohmeier, T.E. and Trippodo, N.C."Control of glomerular filtration rate by Renin-Angiotensin System" Am. J. Physiol. 233(5), (1977) : F 366-372.

Hamilton, W.F., Riley, R.L., Attyah, A.m., Andre Cournand, D.M., Fawell, A., Himmelstein,R.P., Noble, J.W., Remington, D.W., Richard, Jr., Wheeler, N.C. and Witham, A.C. "Comparison of the Fick and Dye Injection Method of Measuring the Cardiac Output in Man" Am. J. Physiol. 153(1948) : 309-321.

Harris, R.C., Hurst, P.E. and Saker, B.M. "Renal Failure After Snake Bite" Med. J. Aust. 2(1976) : 409-411.

Henrich, W.L., Schrier, R.W. and Berl, T. "Mechanism of Renin Secretion During Hemorrhage in the Dog" J. Clin. Invest. 64 (1979) : 1-7.

Henrich, W.L. "Role of Prostaglandin in Renin Secretion" Kidney Int. 19(1981) : 822-830.

Hinshaw, L.B., Gilbert, R.P., Kuida, H. and Visscher, M.B. "Peripheral Resistance Change and Blood Pooling After Endotoxin in Eviscerated Dog" Am. J. Physiol. 195(3), (1958) : 631-634.

Hsu, C.H. and Kurtz, T.W. "Renal Hemodynamic in Experimental ARF" Nephron 27(1981) : 204-208.

Ishwariah, V. and David, J.C. "Pharmacological Action of the Venom of Russell's Viper" Indian J. Med. Res. 19(4). (1932) : 1035-1040.

Iwanaga, S. and Suzuki, T. Enzymes in Snake Venom in Snake Venoms.

(Lee, C.Y. ed) pp. 61-158. Springer-Verlag Berlin Heidelberg, New York, 1979,

Kleinman, A., Page, R.C. and Preisler, P.W. "Effect of prothrombin Clotting Time of the Concentrations of Calcium and other Salts" J. Lab. Clin. Med. 30(1945) : 448.

Knochel, J.P., White, M.G. and Dallas, "The Role of aldosterone in Renal Physiology" Arch. Intern. Med. 131(1973) : 876-884.

Kolmer, J.A., Spaulding, E.H. and Robinson, H.W. in Approved Laboratory Technique. pp. 66-69, Attetton-Century Crofts, Inc. New York, 1951.

Lee, C.Y. "Toxicological Studies on the Venom of Vipera russellii formosensis IV on the Cause of Death in Rabbits" Folia Pharmacol. Jap. 40(1944) : 53-54.

Lee, C.Y. "Toxicological Studies on the Venom of Vipera russellii formosensis maki" J.F.M.A. 47(1948) : 65-98.

Lee, C.Y., Johnson, S.A. and Seegers, W.H. "Clotting of Blood with Russell's Viper Venom" J. Mich. State Med. Soc. 54(1955) : 801-824.

Levens, N.R., Peach, M.J. and Carey, R.M. "Role of the Intrarenal Renin-Angiotensin System in the Control of Renal Function" Circ Res. 48(2), (1981) : 157-167.

Mandal, A.K., Haygood, C.C., Bell, R.D., Sethney, T., James, T.M., Nordquist, J.A., Yunice, A.A. and Lindeman, R.D. "Effects of Acute and Chronic Splenectomy on Experimental Acute Renal Tubular Lesions" J. Lab. Clin. Med. 92(5), (1978) : 698-711.

- Mandal, A.K. "The Spleen and Acute Renal Failure : Mechanisms of Renal Protection by Splenectomy. Involvement of Prostaglandins" Prostaglandins Leukotrienes Med. 9(1982) : 85-107.
- Nashat, F.S. and Portal, R.W., "The Effects of Changes in Hematocrit on Renal Function" J. Physiol 193(1967) : 513-522.
- Oram, S., Ross, G., Pell, L. and Winteler, J. "Renal Cortical Calcification After Snake Bite" Br. Med. J. 1(1963) : 1647-1648.
- Page, R.C., DeBeer, E.J. and Orr, M.L. "Prothrombin Studies Using Russell's Viper Venom. III Effect of Lecithinized Venom on Prothrombin Clotting Time" J. Lab. Clin. Med. 27(1942) : 830.
- Parratt, J.R. and Sturgess, R.M. "The Effect of Indomethacin on the Cardiovascular and Metabolic Responses to E. coli Endotoxin in the Cat" Br.J. Pharmac. 50(1974) : 177-183.
- Plotch, D.W. and Navar, L.G. "Intrarenal effects of the Renin-Angiotensin system" Fed. Proc. 38(1979) : 2280-2285.
- Raab, W. and Kaiser, E. "Nephrotoxic Action of Snake Venom" Mem. Inst. Butantan 33(1966) : 1017-1020.
- Reid, H.A., Chan, K.E. and Thean, P.C. "Plasma Coagulation Defect (Defibrination Syndrome) in Malayan Viper Bite" Lancet 1(1963) : 621-626.
- Ruddy, S., Gigli, I. and Austen, K.F. "The Complement System of Man (Fourth of Four Parts)" N. Engl. J. Med. 287(1972) : 642-646.

Rumpf, K.W., Frenzel, S., Lowitz, H.D. and Scheler, F. "The Effect of Indomethacin on Plasma Renin Activity in Man Under Normal Condition and After Stimulation of the Renin-Angiotensin System" Prostaglandins 10(1975) : 641-648.

Sant, S.M. and Purandare, N.M. "Autopsy of Cases of Snake Bite with Special Reference to Renal Lesion" J. Postgrad. 18(1972) : 181-188.

Sarangi, A., Patnaik, B.C., Das, G.C., Tripathy, N., Misra, G., Snain, A.K. and Das, G.K. "Renal Involvement in Viperine Snake bite" Indian J. Med. Res. 71(1980) : 918-923.

Schreiner, G.E. and Maher, J.F. "Toxic Nephropathy" Am J. Med. 38 (1965) : 409.

Seymour, A.A. and ZeHr, J.E. "Influence of Renal Prostaglandin Synthesis on Renin Control Mechanism in the Dog" Circ Res. 45(1979) : 13-25.

Shastry, J.C., Date, A., Carman, R.H. and Johny, K.V. "Renal Failure Following Snake Bite" Am. J. Trop. Med. 26(3), (1977) : 1032-2038.

Shoukas, A.A., MacAnespie, C.L., Brunner, M.J. and Watermeier, L. "The Importance of the Spleen in Blood Volume Shifts of the Systemic Vascular Bed Caused by the Carotid Sinus Baroreceptor Reflex in the Dog" Circ Res. 49(1981) : 759-767.

Sitprija, V. and Boonpucknavig, V. "Further Observation of Renal Insufficiency in Snake Bite" Nephron 13(1974) : 396-403.

Sitprija, V. and Boonpucknavig, V. "The Kidney in Tropical Snakebite" Clin. Nephrology 8(3), (1977) : 377-383.

Sitprija, V. and Boonpucknavig, V. Snake Venoms and Nephrotoxicity,
in Snake Venom (Lee, C.Y. ed) pp. 997-1018, SpringerVerlag
Berlin Heidelberg, New York, 1979.

Sjodin, J.G., Haggmark, S. and Reiz, S. "Effect of Indomethacin on
Central, Renal and Coronary Hemodynamic" Scand, J. Urol.
Nephrol. 17(1983) : 73-79.

Smith, H.W. in Principle of Renal Physiology, pp. 196-217, Oxford
University press, Inc, 1962.

Starling, E.H. and Lavatt, E.C. in Principles of Human Physiology,
pp. 210, 13th Churchill London, 1962.

Trisnananda, M. "Incidence, Clinical Manifestation and General Manage-
ment of Snake Bite" South Asian J. Trop. Med. Pub. HLH 10(2),
(1979) : 248-254.

Tungthanathanich, P. "Effects of Russell's Viper Venom on Renal func-
tions in Dogs" Master's Thesis, Department of Inter-Depart-
ment of Physiology, Graduate School, Chulalongkorn University,
1983.

Varaguman, T. and Panabokke, R.G. "Bilateral Cortical Necrosis of
the Kidney Following Snakebite" Postgrad Med. J. 46(1970) :
449-451.

Vick, J.A., Ciucuta, H.P. and Menthe, J.H. "Pathophysiological
Studies of Ten Snake Venoms in Animal Toxin. (Russell, F.F.
and Saunders, P.R. eds) pp. 269-282, Oxford Pergamon, 1967.

Whitaker, A.N., McKay, D.G. and Csavossy, I. "Studies on Catecholamine Shock I. Disseminated Intravascular coagulation" Am. J. Path. 56(1969) : 153-176.

Young, D.B., MacCAA, R.E., Pan Yi-JEN and Guyton, A.C. "Effectiveness of the aldosterone, sodium and Potassium Feedback Control System" Am. J. Physiol. 231(3), (1976) : 945-953.

Table 1 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group 1, No 1, weighing 14.6 kg.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 143.3 | 93.3 | 121.1 | 130 |
| HR (beat/min) | 197 | 160 | 164 | 176 |
| CO (ml/min/kg.bw.) | 78.90 | - | 54.01 | 77.26 |
| PV (ml/kg.bw.) | 39.66 | - | 30.41 | 35.96 |
| BV (ml/kg.bw.) | 68.36 | - | 62.12 | 73.42 |
| PCV (%) | 42 | 51 | 51 | 51 |
| TPR (dyne-sec/cm ⁵) | 9,948 | - | 12,326 | 9217 |
| RVR (dyne-sec/cm ⁵) | 34,298 | 181,295 | 207,279 | 277,782 |
| V (μl/min/kg.bw.) | 8.9 | 5.07 | 6.64 | 5.07 |
| RPF (ml/min/kg.bw.) | 13.27 | 1.40 | 1.57 | 1.26 |
| RBF (ml/min/kg.bw.) | 22.89 | 2.82 | 3.21 | 2.56 |
| GFR (ml/min/kg.bw.) | 2.05 | 0.56 | 0.65 | 0.62 |
| F.F (%) | 15.45 | 40.19 | 41.28 | 49.02 |
| R.F. (%) | 29.01 | - | 5.95 | 3.32 |
| P _{Na} (mEq/L) | 149 | 147 | 147 | 147 |
| P _K (mEq/L) | 3.4 | 2.4 | 2.7 | 3.0 |
| P _{Cl} (mEq/L) | 119 | 120 | 117 | 117 |
| U _{Na} V (μEq/min/kg.bw.) | 0.373 | 0.37 | 0.266 | 0.177 |
| U _K V (μEq/min/kg.bw.) | 1.3 | 0.405 | 0.551 | 0.390 |
| U _{Cl} V (μEq/min/kg.bw.) | 0.276 | 0.035 | 0.027 | 0.020 |
| F.E. of Na (%) | 0.122 | 0.449 | 0.278 | 0.196 |
| F.E. of K (%) | 18.65 | 30.12 | 31.42 | 21.13 |
| F.E. of Cl (%) | 0.113 | 0.053 | 0.035 | 0.028 |
| P _{Osm} (mOsm/Kg.) | 294 | 297.2 | 298 | 301.4 |
| U _{Osm} (mOsm/Kg.) | 959.2 | 635.6 | 493 | 514.1 |
| U _{Osm} V (μOsm/min/kg.bw.) | 8.54 | 3.22 | 3.27 | 2.61 |
| C _{Osm} (μOsm/min/kg.bw.) | 29.04 | 10.84 | 10.98 | 8.65 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 20.14 | - 5.77 | - 4.34 | - 3.58 |

Table 2 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group I No 2 weighing 11.5 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|-----------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 136.6 | 88.3 | 58.3 | 78.3 |
| HR (beat/min) | 170 | 180 | 155 | 174 |
| CO (ml/min/kg.bw.) | 88.52 | - | 38.09 | 38.0 |
| PV (ml/kg.bw.) | 42.43 | - | 41.13 | 51.83 |
| BV (ml/kg.bw.) | 59.74 | - | 77.65 | 160 |
| PCV (%) | 29 | 48 | 47 | 45 |
| TPR (dyne-sec/cm ⁵) | 10,732 | - | 16,123 | 14,330 |
| RVR (dyne-sec/cm ⁵) | 49,525 | 429,837 | 506,279 | 2,182,032 |
| V (μL/min/kg.bw.) | 22.61 | 2.43 | 2.87 | 1.13 |
| RPF (ml/min/kg.bw.) | 13.63 | 0.74 | 0.42 | 0.14 |
| RBF (ml/min/kg.bw.) | 19.18 | 1.43 | 0.80 | 0.25 |
| GFR (ml/min/kg.bw.) | 5.27 | 0.35 | 0.24 | 0.08 |
| F.F. (%) | 38.70 | 46.96 | 57.17 | 58.22 |
| R.F. (%) | 21.67 | - | 2.10 | 0.66 |
| P _{Na} (mEq/L.) | 140 | 137 | 135 | 143 |
| P _K (mEq/L.) | 3.5 | 3.0 | 2.7 | 3.2 |
| P _{Cl} (mEq/L.) | 121 | 120 | 116 | 117 |
| U _{Na} V (μEq/min/kg.bw.) | 7.03 | 0.022 | 0.017 | 0.007 |
| U _K V (μEq/min/kg.bw.) | 1.42 | 0.214 | 0.201 | 0.071 |
| U _{Cl} V (μEq/min/kg.bw.) | 5.25 | 0.01 | 0.01 | 0.006 |
| F.E. of Na (%) | 0.952 | 0.046 | 0.053 | 0.059 |
| F.E. of K (%) | 7.72 | 20.45 | 30.67 | 27.82 |
| F.E. of Cl (%) | 0.822 | 0.023 | 0.041 | 0.06 |
| P _{Osm} (mOsm/Kg.) | 289 | 294.6 | 276.4 | 287.2 |
| U _{Osm} (mOsm/Kg.) | 1055.7 | 418.2 | 369 | 337.8 |
| U _{Osm} V (μOsm/min/kg.bw.) | 23.87 | 1.02 | 1.05 | 0.39 |
| C _{Osm} (μOsm/min/kg.bw.) | 82.59 | 3.45 | 3.83 | 1.33 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 59.98 | - 1.02 | - 0.96 | - 0.2 |

Table 3 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group I No 3 weighing 12.1 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 120 | 81.67 | 85 | 110.67 |
| HR (beat/min) | 110 | 86 | 73 | 67 |
| CO (ml/min/kg.bw.) | 151.65 | - | 83.55 | 65.37 |
| PV (ml/kg.bw.) | 38.26 | - | 28.18 | 28.93 |
| BV (ml/kg.bw.) | 58.84 | - | 61.24 | 54.38 |
| PCV (%) | 35 | 54 | 54 | 46 |
| TPR (dyne-sec/cm ⁵) | 5,230 | - | 6,724 | 11,190 |
| RVR (dyne-sec/cm ⁵) | 34,354 | 37,458 | 59,854 | 181,084 |
| V (μ L/min/kg.bw.) | 35.54 | 8.1 | 9.67 | 5.79 |
| RPF (ml/min/kg.bw.) | 15.01 | 6.63 | 4.32 | 2.18 |
| RBF (ml/min/kg.bw.) | 23.09 | 14.41 | 9.39 | 4.04 |
| GFR (ml/min/kg.bw.) | 3.01 | 1.23 | 2.19 | 1.03 |
| FF (%) | 20.06 | 18.56 | 50.62 | 47.31 |
| RF (%) | 15.22 | - | 11.23 | 6.18 |
| P _{Na} (mEq/L.) | 149 | 139 | 149 | 141 |
| P _K (mEq/L.) | 3.5 | 3.0 | 3.2 | 3.3 |
| P _{Cl} (mEq/L.) | 114 | 112 | 114 | 120 |
| U _{Na} V (μ Eq/min/kg.bw.) | 10.55 | 0.93 | 1.09 | .492 |
| U _K V (μ Eq/min/kg.bw.) | 2.84 | 0.94 | 1.61 | 1.18 |
| U _{Cl} V (μ Eq/min/kg.bw.) | 8.07 | 0.58 | 0.198 | 0.13 |
| F.E. of Na (%) | 2.35 | 0.545 | 0.335 | 0.338 |
| F.E. of K (%) | 26.99 | 25.4 | 23.08 | 34.65 |
| F.E. of Cl (%) | 2.35 | 0.423 | 0.078 | 0.107 |
| P _{Osm} (mOsm/Kg.) | 279.2 | 285.4 | 282.4 | 280.6 |
| U _{Osm} (mOsm/Kg.) | 943.1 | 710.4 | 859.3 | 988.6 |
| U _{Osm} V (μ Osm/min/kg.bw.) | 33.52 | 5.75 | 8.31 | 5.72 |
| C _{Osm} (μ Osm/min/kg.bw.) | 120.04 | 20.16 | 29.42 | 20.40 |
| C _{H₂O} (μ Osm/min/kg.bw.) | - 84.5 | - 12.06 | - 19.75 | - 14.61 |

Table 4 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group I No 4, weighing 10.0 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 135.0 | 53.33 | 86.67 | 100.0 |
| HR (beat/min) | 162 | 106 | 147 | 130 |
| CO (ml/min/kg.bw.) | 124.9 | - | 77.3 | 67.0 |
| PV (ml/kg.bw.) | 43.2 | - | 56.0 | 42.8 |
| BV (ml/kg.bw.) | 66.5 | - | 93.3 | 70.2 |
| PCV (%) | 35 | 35 | 40 | 39 |
| TPR (dyne-sec/cm ⁵) | 8,644 | - | 8,967 | 11,937 |
| RVR (syne-sec/cm ⁵) | 65,969 | 56,780 | 80,932 | 70,785 |
| V (μL/min/kg.bw.) | 77 | 42 | 153 | 146 |
| RPF (ml/min/kg.bw.) | 10.63 | 4.88 | 5.14 | 7.14 |
| RBG (ml/min/kg.bw.) | 16.37 | 7.51 | 8.57 | 11.30 |
| GFR (ml/min/kg.bw.) | 4.52 | 1.85 | 2.18 | 3.45 |
| FF (%) | 42.46 | 37.89 | 42.32 | 48.49 |
| RF (%) | 13.10 | - | 11.08 | 16.86 |
| P _{Na} (mEq/L.) | 145 | 151 | 141 | 148 |
| P _K (mEq/L.) | 2.8 | 2.9 | 2.3 | 2.8 |
| P _{Cl} (mEq/L.) | 117 | 117 | 123 | 113 |
| U _{Na} V (μEq/min/kg.bw.) | 11.17 | 1.22 | 4.28 | 3.36 |
| U _K V (μEq/min/kg.bw.) | 1.39 | 1.39 | 0.92 | 0.44 |
| U _{Cl} V (μEq/min/kg.bw.) | 11.47 | 0.76 | 2.75 | 1.31 |
| F.E. of Na (%) | 1.71 | 0.44 | 1.40 | 0.66 |
| F.E. of K (%) | 10.96 | 25.83 | 18.35 | 4.52 |
| F.E. of Cl (%) | 2.17 | 0.35 | 1.03 | 0.34 |
| P _{Osm} (mOsm/Kg.) | 290 | 301.8 | 288.8 | 302 |
| U _{Osm} (mOsm/Kg.) | 455.9 | 275.4 | 157 | 123.1 |
| U _{Osm} V (μOsm/min/kg.bw.) | 35.10 | 11.57 | 24.02 | 17.97 |
| C _{Osm} (μOsm/min/kg.bw.) | 121.05 | 38.33 | 83.11 | 59.51 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 44.05 | + 3.67 | + 69.82 | + 86.49 |

Table 5 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group I, No 5 weighing , 12.5 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 110 | 83.33 | 98.33 | 113.33 |
| HR (beat/min) | 172 | 138 | 144 | 150 |
| CO (ml/min/kg .bw.) | 408.48 | - | 272.0 | 136.56 |
| PV (ml/kg .bw.) | 48.32 | - | 44.32 | 37.84 |
| BV (ml/kg .bw.) | 68.08 | - | 59.12 | 50.48 |
| PCV (%) | 29 | 28 | 25 | 25 |
| TPR (dyne-sec/cm ⁵) | 1,723 | - | 2,313 | 5,309 |
| RVR (dyne-sec/cm ⁵) | 48,156 | 44,393 | 63,137 | 88,456 |
| V (μ L/min.kg .bw.) | 44.8 | 67.2 | 32.0 | 24.0 |
| RPF (ml/min/kg .bw.) | 10.38 | 8.65 | 7.47 | 6.15 |
| RBF (ml/min/kg .bw.) | 14.62 | 12.01 | 9.96 | 8.20 |
| GFR (ml/min/kg .bw.) | 2.92 | 2.67 | 2.81 | 3.47 |
| FF (%) | 28.12 | 30.93 | 37.58 | 56.49 |
| RF (%) | 3.58 | - | 3.66 | 6.00 |
| P _{Na} (mEq/L.) | 146 | 141 | 142 | 146 |
| P _K (mEq/L.) | 2.9 | 2.6 | 2.9 | 3.4 |
| P _{Cl} (mEq/L.) | 109 | 110 | 98 | 112 |
| U _{Na} V (μ Eq/min/kg.bw.) | 10.62 | 11.76 | 7.04 | 5.52 |
| U _K V (μ Eq/min/kg.bw.) | 0.90 | 1.28 | 1.09 | 2.04 |
| U _{Cl} V (μ Eq/min/kg.bw.) | 9.96 | 7.32 | 3.55 | 2.4 |
| F.E. fo Na (%) | 2.49 | 3.12 | 1.77 | 1.09 |
| F.E. of K (%) | 10.60 | 18.36 | 13.36 | 17.28 |
| F.E. of Cl (%) | 3.13 | 2.49 | 1.29 | 0.61 |
| P _{Osm} (mOsm/Kg.) | 284.2 | 285 | 284.8 | 263.8 |
| U _{Osm} (mOsm/Kg.) | 653.5 | 446.6 | 615.2 | 786 |
| U _{Osm} V (μ Osm/minkg.bw.) | 29.28 | 30.01 | 19.69 | 18.86 |
| C _{Osm} (μ Osm/min/kg.bw.) | 103.01 | 105.31 | 69.12 | 71.51 |
| C _{H₂O} (μ Osm/min/kg.bw.) | - 58.21 | - 38.1 | - 37.12 | - 47.51 |

Table 6 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group II No 1, weighing 12.0 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|--------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 95.0 | 85.0 | 98.5 | 68.3 |
| HR (beat/min) | 161 | 157 | 172 | 134 |
| CO (ml/min/kg.bw.) | 123.5 | - | 75.58 | 61.67 |
| PV (ml/kg.bw.) | 50.92 | - | 56.42 | 36.08 |
| BV (ml/kg.bw.) | 70.67 | - | 71.33 | 44.0 |
| PCV (%) | 22 | 21 | 21 | 18 |
| TPR (dyne-sec/cm ⁵) | 5,126 | - | 8,668 | 7,382 |
| RVR (dyne-sec/cm ⁵) | 47,643 | 77,553 | 77,665 | 61,879 |
| V (μL/min/kg.bw.) | 16.67 | 8.5 | 8.75 | 7.92 |
| RPF (ml/min/kg.bw.) | 6.93 | 5.77 | 6.66 | 6.03 |
| RBF (ml/min/kg.bw.) | 9.63 | 7.31 | 8.44 | 7.36 |
| GFR (ml/min/kg.bw.) | 2.07 | 1.25 | 1.87 | 2.00 |
| F.F (%) | 29.86 | 22.09 | 28.09 | 33.20 |
| R.F. (%) | 7.79 | - | 11.16 | 11.93 |
| P _{Na} (mEq/L.) | 137 | 143 | 143 | 129 |
| P _K (mEq/L.) | 4.0 | 3.1 | 3.4 | 3.1 |
| P _{Cl} (mEq/L.) | 125 | 118 | 125 | 119 |
| U _{Na} V (μEq/min/kg.bw.) | 3.60 | 1.22 | 1.04 | 1.14 |
| U _K V (μEq/min/kg.bw.) | 1.92 | 1.28 | 1.6 | 1.73 |
| U _{Cl} V (μEq/min/kg.bw.) | 2.42 | 0.32 | 0.53 | 0.43 |
| F.E. of Na (%) | 1.27 | 0.67 | 0.39 | 0.44 |
| F.E. of K (%) | 23.15 | 32.47 | 25.16 | 27.8 |
| F.E. of Cl (%) | 0.93 | 0.21 | 0.23 | 0.18 |
| P _{Osm} (mOsm/Kg.) | 276 | 274 | 273.2 | 271 |
| U _{Osm} (mOsm/Kg.) | 1191.7 | 1343.4 | 1315.8 | 1350.2 |
| U _{Osm} V (μOsm/min/kg.bw.) | 19.87 | 11.42 | 11.52 | 10.69 |
| C _{Osm} (μOsm/min/kg.bw.) | 71.98 | 41.67 | 42.14 | 39.46 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 55.31 | - 33.17 | - 33.39 | 31.54 |

Table 7 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group II No 2 weighing 13.0 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 126.6 | 73.3 | 98.3 | 108.3 |
| HR (beat/min) | 228 | 190 | 184 | 194 |
| CO (ml/min/kg.bw.) | 133.23 | - | 44.46 | 48.23 |
| PV (ml/kg.bw.) | 35.77 | - | 25.69 | 23.15 |
| BV (ml/kg.bw.) | 55.08 | - | 40.15 | 36.23 |
| PCV (%) | 35 | 36 | 36 | 36 |
| TPR (dyne-sec/cm ⁵) | 5,846 | - | 13,602 | 13,814 |
| RVR (syne-sec/cm ⁵) | 45,956 | 41,899 | 63.609 | 110,187 |
| V (μL/min/kg.bw.) | 95.38 | 24.62 | 83.08 | 61.54 |
| RPF (ml/min/kg.bw.) | 11.02 | 6.89 | 6.07 | 3.87 |
| RBF (ml/min/kg.bw.) | 16.95 | 10.76 | 9.51 | 6.05 |
| GFR (ml/min/kg/bw.) | 2.94 | 2.79 | 3.08 | 2.16 |
| FF (%) | 26.65 | 40.52 | 50.68 | 55.73 |
| RF (%) | 12.72 | - | 21.38 | 12.53 |
| P _{Na} (mEq/L.) | 127 | 138 | 143 | 141 |
| P _K (mEq/L.) | 3.2 | 2.8 | 3.0 | 3.5 |
| P _{Cl} (mEq/L.) | 115 | 115 | 120 | 117 |
| U _{Na} V (μEq/min/kg.bw.) | 2.29 | 1.31 | 2.24 | 2.46 |
| U _K V (μEq/min/kg.bw.) | 2.10 | 1.60 | 1.41 | 1.42 |
| U _{Cl} V (μEq/min/kg.bw.) | 1.53 | 0.54 | 1.08 | 0.49 |
| F.E. of Na (%) | 0.61 | 0.34 | 0.51 | 0.81 |
| F.E. of K (%) | 22.33 | 20.47 | 15.29 | 18.75 |
| F.E. of Cl(%) | 0.45 | 0.17 | 0.29 | 0.20 |
| P _{Osm} (mOsm/Kg.) | 278.2 | 259.2 | 277.2 | 273.4 |
| U _{Osm} (mOsm/Kg.) | 172.5 | 386.8 | 117.8 | 156.6 |
| U _{Osm} V (μOsm/min/kg.bw.) | 16.45 | 9.52 | 9.79 | 9.64 |
| C _{Osm} (μOsm/min/kg.bw.) | 59.14 | 36.74 | 35.31 | 35.25 |
| C _{H₂O} (μOsm/min/kg.bw.) | + 36.24 | - 12.12 | + 47.77 | + 26.29 |

Table 8 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group II No 3 weighing 13.5 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 100.0 | 83.3 | 98.3 | 113.3 |
| HR (beat/min) | 123 | 133 | 125 | 130 |
| CO (ml/min/kg.bw.) | 82.81 | - | 52.96 | 35.85 |
| PV (ml/kg.bw.) | 41.48 | - | 36.74 | 34.96 |
| BV (ml/kg.bw.) | 54.59 | - | 46.44 | 43.63 |
| PCV (%) | 24 | 24 | 21 | 20 |
| TPR(dyne-sec/cm ⁵) | 7,154 | - | 10,996 | 18,723 |
| RVR (dyne-sec/cm ⁵) | 20,450 | 59,137 | 111,740 | 152,017 |
| V (μL/min/kg.bw.) | 48.15 | 16.30 | 11.85 | 14.07 |
| RPF (ml/min/kg.bw.) | 20.28 | 6.34 | 4.12 | 3.53 |
| RBF (ml/min/kg.bw.) | 28.96 | 8.35 | 5.21 | 4.42 |
| GFR (ml/min/kg.bw.) | 4.54 | 1.49 | 1.17 | 1.36 |
| F.F. (%) | 22.40 | 23.53 | 28.41 | 38.62 |
| RF (%) | 34.98 | - | 9.84 | 12.32 |
| P _{Na} (mEq/L.) | 133 | 150 | 118 | 130 |
| P _K (mEq/L.) | 3.4 | 3.1 | 2.7 | 3.1 |
| P _{Cl} (mEq/L.) | 123 | 115 | 121 | 122 |
| U _{Na} V (μEq/min/kg.bw.) | 6.6 | 0.23 | 0.13 | 0.34 |
| U _K V (μEq/min/kg.bw.) | 3.42 | 1.52 | 1.15 | 1.52 |
| U _{Cl} V (μEq/min/kg.bw.) | 6.16 | 0.16 | 0.95 | 0.10 |
| F.E. of Na (%) | 1.09 | 0.10 | 0.09 | 0.19 |
| F.E. of K (%) | 22.14 | 32.75 | 36.40 | 35.94 |
| F.E. of Cl (%) | 1.10 | 0.10 | 0.07 | 0.06 |
| P _{Osm} (mOsm/Kg.) | 262.6 | 276.2 | 276.2 | 282.4 |
| U _{Osm} (mOsm/Kg.) | 579.4 | 407.8 | 432 | 441.4 |
| U _{Osm} V (μOsm/min/kg.bw.) | 27.90 | 6.65 | 5.12 | 6.21 |
| C _{Osm} (μOsm/min/kg.bw.) | 106.24 | 24.07 | 18.53 | 21.99 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 58.09 | - 7.77 | - 6.68 | - 7.97 |

Table 9 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group II No 4 weighing 14.0 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|--------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 123.3 | 95 | 108.3 | 115 |
| HR (beat/min) | 170 | 144 | 138 | 138 |
| CO (ml/min/kg.bw.) | 179.29 | - | 134.29 | 223.57 |
| PV (ml/kg.bw.) | 42.09 | - | 43.21 | 79.36 |
| BV (ml/kg.bw.) | 56.07 | - | 56.86 | 104.43 |
| PCV (%) | 24 | 24 | 24 | 24 |
| TPR (dyne-sec/cm ⁵) | 3,929 | - | 4,607 | 2,939 |
| RVR (dyne-sec/cm ⁵) | 57,298 | 40,898 | 75,478 | 99,327 |
| V (μL/min/kg.bw.) | 25.0 | 134.29 | 78.57 | 40.0 |
| RPF (ml/min/kg.bw.) | 9.22 | 10.09 | 6.23 | 5.03 |
| RBF (ml/min/kg.bw.) | 12.29 | 13.27 | 8.14 | 6.61 |
| GFR (ml/min/kg.bw.) | 3.50 | 3.63 | 2.66 | 2.23 |
| FF (%) | 37.98 | 35.98 | 42.74 | 44.36 |
| RF (%) | 6.86 | - | 6.10 | 2.96 |
| P _{Na} (mEq/L.) | 141 | 145 | 162 | 133 |
| P _K (mEq/L.) | 3.0 | 2.0 | 2.4 | 2.4 |
| P _{Cl} (mEq/L.) | 121 | 116 | 120 | 118 |
| U _{Na} V (μEq/min/kg.bw.) | 1.98 | 2.43 | 1.26 | 1.28 |
| U _K V (μEq/min/kg.bw.) | 2.03 | 2.55 | 0.63 | 1.36 |
| U _{Cl} V (μEq/min/kg.bw.) | 1.88 | 0.67 | 0.86 | 0.44 |
| F.E. of Na (%) | 0.40 | 0.50 | 0.29 | 0.43 |
| F.E. of K (%) | 19.28 | 35.16 | 9.84 | 25.41 |
| F.E. of Cl (%) | 0.44 | 0.16 | 0.27 | 0.17 |
| P _{Osm} (mOsm/Kg.) | 276.8 | 291 | 272 | 290.4 |
| U _{Osm} (mOsm/Kg.) | 723.8 | 131.6 | 107 | 272.6 |
| U _{Osm} V (μOsm/min/kg.bw.) | 18.09 | 17.67 | 8.41 | 10.9 |
| C _{Osm} (μOsm/min/kg.bw.) | 65.37 | 60.73 | 30.91 | 37.55 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 40.37 | + 73.56 | + 47.66 | + 2.45 |

Table 10 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group II No 5 weighing 9.5 kgs.

| parameter | control | Post venom injection | | |
|--|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 123.3 | 116.67 | 11.3.3 | 96.67 |
| HR (beat/min) | 210 | 234 | 234 | 222 |
| CO (ml/min/kg.bw.) | 177.89 | - | 186.11 | - |
| PV (ml/kg.bw.) | 62.21 | - | 50.53 | 50.74 |
| BV (ml/kg.bw.) | 81.89 | - | 64.84 | 65.05 |
| PCV (%) | 24 | 24 | 22 | 22 |
| TPR (dyne-sec/cm ⁵) | 5,835 | - | 5,125 | - |
| RVR (dyne-sec/cm ⁵) | 45,904 | 48,166 | 61,640 | 58.273 |
| V (μ L/min/kg.bw.) | 12.63 | 11.58 | 9.47 | 8.74 |
| RPF (ml/min/kg.bw.) | 17.19 | 15.30 | 12.07 | 10.89 |
| RBF (ml/min/kg.bw.) | 22.61 | 20.39 | 15.47 | 13.97 |
| GFR (ml/min/kg.bw.) | 3.60 | 3.19 | 2.79 | 2.38 |
| F.F (%) | 20.97 | 20.88 | 23.09 | 21.89 |
| R.F. (%) | 12.71 | - | 8.32 | - |
| P _{Na} (mEq/L.) | 140 | 124 | 116 | 129 |
| P _K (mEq/L.) | 4.0 | 2.7 | 2.5 | 2.7 |
| P _{Cl} (mEq/L.) | 118 | 124 | 123 | 130 |
| U _{Na} V (μ Eq/min/kg.bw.) | 0.62 | 0.82 | 0.82 | 0.72 |
| U _K V (μ Eq/min/kg.bw.) | 2.88 | 2.39 | 1.88 | 1.99 |
| U _{Cl} V (μ Eq/min/kg.bw.) | 1.10 | 0.38 | 0.33 | 2.62 |
| F.E. of Na (%) | 0.12 | 0.21 | 0.25 | 0.23 |
| F.E. of K (%) | 19.98 | 27.65 | 26.92 | 30.94 |
| F.E. of Cl (%) | 0.26 | 0.10 | 0.10 | 0.09 |
| P _{Osm} (mOsm/Kg.) | 246.6 | 288.2 | 288.2 | 282 |
| U _{Osm} (mOsm/Kg.) | 1808 | 1439.6 | 1429 | 1298 |
| U _{Osm} V (μ Osm/min/kg.bw.) | 22.84 | 16.67 | 13.53 | 11.34 |
| C _{Osm} (μ Osm/min/kg.bw.) | 92.6 | 57.83 | 46.96 | 40.21 |
| C _{H2O} (μ Osm/min/kg.bw.) | - 79.97 | - 46.25 | - 37.49 | - 31.47 |

Table 11 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group III No 1 weighing 11.0 kgs.

| parameter | control | Post venom injection | | |
|--------------------------------------|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 135 | 108.33 | 113.33 | 121.67 |
| HR (beat/min) | 196 | 180 | 216 | 210 |
| CO (ml/min/kg.bw.) | 291.82 | - | 183.64 | 149.09 |
| PV (ml/kg.bw.) | 62.91 | - | 38.18 | 40.35 |
| BV (ml/kg.bw.) | 85.0 | - | 51.64 | 54.82 |
| PCV (%) | 26 | 26 | 26 | 26 |
| TPR (dyne-sec/cm ⁵) | 3,664 | - | 4,487 | 5,934 |
| RVR (dyne-sec/cm ⁵) | 112,824 | 100,770 | 110,781 | 91,631 |
| V (μL/min/kg.bw.) | 8.18 | 11.82 | 9.09 | 12.73 |
| RPF (ml/min/kg.bw.) | 6.44 | 5.78 | 5.50 | 7.14 |
| RBF (ml/min/kg.bw.) | 8.70 | 7.82 | 7.44 | 9.65 |
| GFR (ml/min/kg.bw.) | 2.80 | 2.90 | 2.72 | 2.85 |
| F.E. (%) | 43.48 | 50.10 | 49.45 | 53.91 |
| R.F. (%) | 2.98 | - | 4.05 | 6.48 |
| P _{Na} (mEq/L.) | 137 | 124 | 143 | 126 |
| P _K (mEq/L.) | 3.4 | 2.9 | 3.1 | 3.1 |
| P _{Cl} (mEq/L.) | 125 | 112 | 122 | 110 |
| U _{Na} V (μEq/min/kg.bw.) | 0.87 | 1.83 | 2.01 | 3.27 |
| U _K V (μEq/min/kg.bw.) | 2.17 | 2.33 | 2.87 | 3.30 |
| C _{Cl} V (μEq/min/kg.bw.) | 0.16 | 0.36 | 0.25 | 0.27 |
| F.E. of Na (%) | 0.23 | 0.51 | 0.52 | 0.67 |
| F.E. of K (%) | 22.78 | 27.70 | 34.05 | 27.93 |
| F.E. of Cl(%) | 0.04 | 0.11 | 0.07 | 0.06 |
| P _{Osm} (mOsm/ Kg.) | 285 | 275.2 | 270 | 272.6 |
| U _{Osm} (mOsm/ Kg.) | 1913 | 1686.7 | 1959.4 | 1929 |
| U _{Osm} V (μOsm/min/kg.bw.) | 15.65 | 19.94 | 17.81 | 24.56 |
| C _{Osm} (μOsm/min/kg.bw.) | 54.91 | 72.44 | 65.97 | 90.08 |
| C _{H2O} (μOsm/min/kg.bw.) | - 46.73 | - 60.62 | - 56.88 | - 77.35 |

Table 12 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group III No 2 weighing 15.0 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|--------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 103.33 | 128.33 | 128.33 | 121.67 |
| HR (beat/min) | 126 | 140 | 152 | 156 |
| CO (ml/min/kg.bw.) | 80.93 | - | 51.0 | 44.2 |
| PV (ml/kg.bw.) | 50.2 | - | 28.67 | 27.67 |
| BV (ml/kg.bw.) | 71.73 | - | 49.4 | 47.67 |
| PCV (%) | 30 | 42 | 42 | 42 |
| TPR (dyne-sec/cm ⁵) | 6,808 | - | 13,417 | 14,677 |
| RVR (dyne-sec/cm ⁵) | 64,339 | 91,445 | 78,662 | 108,449 |
| V (μ L/min/kg.bw.) | 9.33 | 11.33 | 14.67 | 10.0 |
| RPF (ml/min/kg.bw.) | 5.99 | 4.34 | 5.05 | 3.47 |
| RBF (ml/min/kg.bw.) | 8.56 | 7.48 | 8.70 | 5.98 |
| GFR (ml/min/kg.bw.) | 2.95 | 2.69 | 2.67 | 1.88 |
| FF (%) | 49.24 | 61.87 | 52.99 | 54.23 |
| R.F. (%) | 10.58 | - | 14.67 | 13.53 |
| P _{Na} (mEq/L.) | 128 | 148 | 129 | 152 |
| P _K (mEq/L.) | 3.2 | 3.6 | 3.3 | 3.7 |
| P _{Cl} (mEq/L.) | 111 | 107 | 107 | 121 |
| U _{Na} V (μ Eq/min/kg.bw.) | 2.01 | 1.12 | 0.98 | 0.59 |
| U _K V (μ Eq/min/kg.bw.) | 2.05 | 1.68 | 1.80 | 1.54 |
| U _{Cl} V (μ Eq/min/kg.bw.) | 0.46 | 0.19 | 0.22 | 0.14 |
| F.E. of Na (%) | 0.53 | 0.28 | 0.28 | 0.21 |
| F.E. of K (%) | 21.74 | 17.35 | 20.45 | 22.12 |
| F.E. of Cl (%) | 0.14 | 0.07 | 0.08 | 0.06 |
| P _{Osm} (mOsm/Kg.) | 248.2 | 278.8 | 295 | 287.6 |
| U _{Osm} (mOsm/Kg.) | 2064 | 1729 | 1334.7 | 1196.6 |
| U _{Osm} V (μ Osm/min/kg.bw.) | 19.26 | 19.59 | 19.58 | 11.97 |
| C _{Osm} (μ Osm/min/kg.bw.) | 77.59 | 70.26 | 66.37 | 41.61 |
| C _{H₂O} (μ Osm/min/kg.bw.) | - 68.26 | - 58.93 | - 51.7 | - 31.61 |

Table 13 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group III No 3 weighing 13.5 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 140 | 136.67 | 135.0 | 123.33 |
| HR (beat/min) | 186 | 162 | 156 | 186 |
| CO (ml/min/kg.bw.) | 154.07 | - | 89.41 | 105.19 |
| PV (ml/kg.bw.) | 65.63 | - | 55.63 | 55.33 |
| BV (ml/kg.bw.) | 91.11 | - | 75.11 | 73.78 |
| PCV (%) | 28 | 30 | 26 | 25 |
| TPR (dyne-sec/cm ⁵) | 5,383 | - | 8,946 | 6,946 |
| RVR (dyne-sec/cm ⁵) | 75,194 | 62,327 | 79,556 | 75,005 |
| V (μL/min/kg.bw.) | 8.15 | 7.78 | 9.26 | 10.0 |
| RPF (ml/min/kg.bw.) | 7.93 | 0.09 | 7.44 | 7.31 |
| RBF (ml/min/kg.bw.) | 11.03 | 12.99 | 10.05 | 9.74 |
| GFR (ml/min/kg.bw.) | 3.49 | 3.59 | 3.41 | 2.70 |
| F.E. (%) | 43.92 | 39.43 | 45.80 | 37.01 |
| R.F. (%) | 7.16 | - | 11.24 | 9.26 |
| P _{Na} (mEq/L.) | 133 | 127 | 125 | 144 |
| P _K (mEq/L.) | 2.7 | 2.7 | 2.7 | 3.4 |
| P _{Cl} (mEq/L.) | 115 | 124 | 108 | 120 |
| U _{Na} V (μEq/min/kg.bw.) | 1.58 | 0.48 | 1.09 | 1.18 |
| U _K V (μEq/min/kg.bw.) | 1.76 | 2.29 | 2.42 | 2.44 |
| U _{Cl} V (μEq/min/kg.bw.) | 0.42 | 0.19 | 0.18 | 0.15 |
| F.E. of Na (%) | 0.34 | 0.11 | 0.26 | 0.30 |
| F.E. of K (%) | 18.69 | 23.70 | 26.27 | 26.54 |
| F.E. of Cl (%) | 0.11 | 0.04 | 0.05 | 0.05 |
| P _{Osm} (mOsm/Kg.) | 299.2 | 275.4 | 282 | 300.8 |
| U _{Osm} (mOsm/Kg.) | 2087 | 1357.8 | 1458.2 | 1287.7 |
| U _{Osm} V (μOsm/min/kg.bw.) | 17.01 | 10.56 | 13.50 | 12.88 |
| C _{Osm} (μOsm/min/kg.bw.) | 56.85 | 38.35 | 47.88 | 42.81 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 48.7 | - 30.57 | - 38.62 | - 32.81 |

Table 14 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group III No 4 weighing 14.0 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 136.67 | 128.33 | 138.33 | 136.67 |
| HR (beat/min) | 172 | 160 | 150 | 174 |
| CO (ml/min/kg.bw.) | 348.57 | - | 46.0 | 114.07 |
| PV (ml/kg.bw.) | 58.14 | - | 41.86 | 43.79 |
| BV (ml/kg.bw.) | 83.07 | - | 64.5 | 66.36 |
| PCV (%) | 30 | 39 | 35 | 34 |
| TPR (dyne-sec/cm ⁵) | 2,240 | - | 9,189 | 6,844 |
| RVR (dyne-sec/cm ⁵) | 83,620 | 88,344 | 79,772 | 86,355 |
| V (μ L/min/kg : bw.) | 10.36 | 9.64 | 11.43 | 12.5 |
| RPF (ml/min/kg .bw.) | 6.54 | 5.06 | 6.44 | 5.79 |
| RBF (ml/min/kg .bw.) | 9.38 | 8.30 | 9.91 | 9.04 |
| GFR (ml/ming/kg .bw.) | 3.08 | 2.65 | 3.00 | 3.06 |
| FF (%) | 47.15 | 52.25 | 46.63 | 51.32 |
| RF (%) | 2.68 | - | 11.52 | 7.93 |
| P _{Na} (mEq/L.) | 135 | 143 | 141 | 140 |
| P _K (mEq/L.) | 2.7 | 2.7 | 3.4 | 3.8 |
| P _{Cl} (mEq/L.) | 103 | 120 | 114 | 107 |
| U _{Na} V (μ Eq/min/kg .bw.) | 3.15 | 2.45 | 1.49 | 2.68 |
| U _K V (μ Eq/min/kg .bw.) | 0.94 | 1.52 | 1.38 | 1.85 |
| U _{Cl} V (μ Eq/min/kg .bw.) | 1.33 | 0.52 | 0.47 | 0.50 |
| F.E. of Na (%) | 0.76 | 0.65 | 0.35 | 0.62 |
| F.E. of K (%) | 11.33 | 21.33 | 13.54 | 15.90 |
| F.E. of Cl (%) | 0.42 | 0.16 | 0.14 | 0.15 |
| P _{Osm} (mOsm/Kg.) | 294.8 | 292.8 | 297.2 | 291.2 |
| U _{Osm} (mOsm/Kg.) | 1964 | 1438.5 | 1460.9 | 1479 |
| U _{Osm} V (μ Osm/min.kg.bw.) | 20.35 | 13.87 | 16.70 | 18.49 |
| C _{Osm} (μ Osm/min/kg.bw.) | 69.02 | 47.36 | 56.18 | 63.49 |
| C _{H₂O} (μ Osm/min/kg.bw.) | - 58.66 | - 37.72 | - 44.75 | - 50.99 |

Table 15 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group III No 5 weighing 13.6 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 138.33 | 121.67 | 133.33 | 133.33 |
| HR (beat/min) | 144 | 150 | 156 | 162 |
| CO (ml/min/kg.bw.) | 128.68 | - | 65.29 | 53.75 |
| PV ((ml/kg.bw.) | 45.15 | - | 35.59 | 48.46 |
| BV (ml/kg.bw.) | 72.79 | - | 82.79 | 105.39 |
| PCV (%) | 38 | 63 | 57 | 54 |
| TPR (dyne-sec/cm ⁵) | 6,322 | - | 12,009 | 14,588 |
| RVR (dyne-sec/cm ⁵) | 54.402 | 83,774 | 117,184 | 165,715 |
| V (μL/min/kg.bw.) | 9.93 | 4.04 | 6.1 | 5.15 |
| RPF (ml/min/kg.bw.) | 8.64 | 3.16 | 2.88 | 2.18 |
| RBG (ml/min/kg.bw.) | 13.92 | 8.54 | 6.69 | 4.73 |
| GFR (ml/min/kg.bw.) | 3.28 | 1.65 | 1.40 | 0.97 |
| FF (%) | 37.99 | 52.19 | 48.73 | 44.76 |
| RF (%) | 10.83 | - | 10.25 | 8.80 |
| P _{Na} (mEq/L.) | 140 | 138 | 142 | 123 |
| P _K (mEq/L.) | 3.3 | 3.1 | 3.2 | 3.0 |
| P _{Cl} (mEq/L.) | 120 | 115 | 116 | 124 |
| U _{Na} V (μEq/min/kg.bw.) | 0.60 | 0.06 | 0.05 | 0.04 |
| U _K V (μEq/min/kg.bw.) | 1.61 | 0.95 | 0.89 | 0.67 |
| U _{Cl} V (μEq/min/kg.bw.) | 0.14 | 0.05 | 0.04 | 0.04 |
| F.E. of Na (%) | 0.01 | 0.03 | 0.03 | 0.03 |
| F.E. of K (%) | 14.85 | 18.51 | 19.72 | 22.89 |
| F.E. of Cl (%) | 0.04 | 0.03 | 0.03 | 0.03 |
| P _{Osm} (mOsm/Kg.) | 289.8 | 284.8 | 281.2 | 276.2 |
| U _{Osm} (mOsm/Kg.) | 1983.5 | 1608.6 | 1060.4 | 828.1 |
| U _{Osm} V (μOsm/min/kg.bw.) | 19.70 | 6.49 | 6.47 | 4.26 |
| C _{Osm} (μOsm/min/kg.bw.) | 67.69 | 22.82 | 23.00 | 15.44 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 55.03 | - 18.78 | - 16.9 | - 10.29 |

Table 16 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group IV, No 1 weighing 13.0 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 145 | 161.67 | 165 | 170 |
| HR (beat/min) | 198 | 186 | 174 | 180 |
| CO (ml/min/kg.bw.) | 241.69 | - | 72.85 | 64.92 |
| PV (ml/kg.bw.) | 58.0 | - | 88.85 | 46.77 |
| BV (ml/kg.bw.) | 80.62 | - | 123.46 | 71.0 |
| PCV (%) | 28 | 28 | 28 | 28 |
| TPR (dyne-sec/cm ⁵) | 3,691 | - | 13,935 | 16,110 |
| RVR (dyne-sec/cm ⁵) | 72,718 | 86,145 | 111.224 | 192,341 |
| V(μL/min/kg.bw.) | 11.92 | 14.62 | 10.38 | 5.15 |
| RPF (ml/min/kg.bw.) | 8.83 | 7.78 | 6.57 | 3.92 |
| RBF (ml/min/kg.bw.) | 12.26 | 11.15 | 9.13 | 5.44 |
| GFR (ml/min/kg.bw.) | 4.46 | 2.80 | 3.46 | 2.19 |
| FF (%) | 50.52 | 35.94 | 52.70 | 55.89 |
| RF (%) | 5.08 | - | 12.53 | 8.38 |
| P _{Na} (mEq/L.) | 136 | 134 | 142 | 142 |
| P _K (mEq/L.) | 2.6 | 3.1 | 3.5 | 3.5 |
| P _{Cl} (mEq/L.) | 108 | 119 | 113 | 107 |
| U _{Na} V (μEq/min/kg.bw.) | 4.76 | 3.52 | 2.09 | 0.90 |
| U _K V (μEq/min/kg.bw.) | 0.68 | 0.45 | 0.57 | 0.87 |
| U _{Cl} V (μEq/min/kg.bw.) | 4.29 | 1.59 | 0.56 | 0.16 |
| F.E. of Na (%) | 0.78 | 0.94 | 0.43 | 0.29 |
| F.E. of K (%) | 5.86 | 5.22 | 4.72 | 11.37 |
| F.E. of Cl (%) | 0.89 | 0.48 | 0.14 | 0.07 |
| P _{Osm} (mOsm/Kg.) | 293.8 | 291 | 294.2 | 291 |
| U _{Osm} (mOsm/Kg.) | 1798.4 | 948.9 | 1234.6 | 1416.1 |
| U _{Osm} V (μOsm/min/kg.bw.) | 21.43 | 13.82 | 12.82 | 7.29 |
| C _{Osm} (μOsm/min/kg.bw.) | 72.97 | 47.67 | 43.56 | 25.06 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 61.04 | - 33.05 | - 33.18 | - 19.91 |

Table 17 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group IV, No 2 weighing 11.5 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 125 | 116.67 | 90.00 | 88.33 |
| HR (beat/min) | 174 | 198 | 168 | 180 |
| CO (ml/min/kg.bw.) | 197.48 | - | 115.22 | 138.52 |
| PV (ml/kg.bw.) | 63.74 | - | 46.61 | 62.87 |
| BV (ml/kg.bw.) | 87.39 | - | 60.61 | 78.61 |
| PCV (%) | 27 | 25 | 23 | 20 |
| TPR (dyne-sec/cm ⁵) | 4,402 | - | 5,433 | 4,435 |
| RVR (dyne-sec/cm ⁵) | 69,981 | 90,975 | 69,287 | 74,177 |
| V (μL/min/kg.bw.) | 25.22 | 16.09 | 17.83 | 13.91 |
| RPF (ml/min/kg.bw.) | 9.07 | 6.69 | 6.97 | 6.63 |
| RBF (ml/min/kg.bw.) | 12.42 | 8.92 | 9.03 | 8.28 |
| GFR (ml/min/kg.bw.) | 3.33 | 2.47 | 2.95 | 2.44 |
| FF (%) | 36.67 | 36.96 | 42.43 | 36.83 |
| RF (%) | 6.29 | - | 7.84 | 5.97 |
| P _{Na} (mEq/L.) | 137 | 126 | 122 | 121 |
| P _K (mEq/L.) | 3.4 | 3.0 | 2.8 | 3.0 |
| P _{Cl} (mEq/L.) | 109 | 120 | 111 | 119 |
| U _{Na} V (μEq/min/kg.bw.) | 0.83 | 0.71 | 1.05 | 0.61 |
| U _K V (μEq/min/kg.bw.) | 2.22 | 2.08 | 2.09 | 2.35 |
| U _{Cl} V (μEq/min/kg.bw.) | 1.10 | 0.13 | 0.14 | 0.14 |
| F.E. of Na (%) | 0.18 | 0.23 | 0.29 | 0.21 |
| F.E. of K (%) | 19.63 | 27.99 | 25.24 | 32.12 |
| F.E. of Cl (%) | 0.03 | 0.04 | 0.04 | 0.05 |
| P _{Osm} (mOsm/Kg.) | 266.2 | 282.2 | 295.4 | 296 |
| U _{Osm} (mOsm/Kg.) | 1218.8 | 1166.4 | 1265.3 | 1234 |
| U _{Osm} V (μOsm/min/kg.bw.) | 30.74 | 18.77 | 22.69 | 17.16 |
| C _{Osm} (μOsm/min/kg.bw.) | 115.47 | 66.50 | 76.37 | 57.99 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 90.25 | - 50.41 | - 58.54 | - 44.08 |

Table 18 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group IV, No 3 weighing 12.8 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|---------|---------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 100.0 | 101.67 | 78.33 | 70.0 |
| HR (beat/min) | 168 | 200 | 186 | 198 |
| CO (ml/min/kg.bw.) | 370.39 | - | - | 139.92 |
| PV (ml/kg.bw.) | 59.45 | - | 54.77 | 44.14 |
| BV (ml/kg.bw.) | 79.22 | - | 70.23 | 55.16 |
| PCV (%) | 25 | 23 | 22 | 20 |
| TPR (dyne-sec/cm ⁵) | 1,687 | - | - | 3,126 |
| RVR (dyne-sec/cm ⁵) | 58,865 | 81,348 | 51,673 | 46,807 |
| V (μL/min/kg.bw.) | 17.19 | 12.11 | 15.23 | 12.5 |
| RPF (ml/min/kg.bw.) | 7.96 | 6.01 | 7.39 | 7.48 |
| RBF (ml/min/kg.bw.) | 10.61 | 7.81 | 9.47 | 9.34 |
| GFR (ml/min/kg.bw.) | 2.29 | 1.96 | 2.02 | 2.15 |
| FF (%) | 28.81 | 32.52 | 27.33 | 28.75 |
| EF (%) | 2.87 | - | - | 6.68 |
| P _{Na} (mEq/L.) | 144 | 144 | 144 | 143 |
| P _K (mEq/L.) | 2.9 | 2.5 | 2.8 | 3.4 |
| P _{Cl} (mEq/L.) | 98 | 93 | 99 | 96 |
| U _{Na} V (μEq/min/kg.bw.) | 0.96 | 1.05 | 1.31 | 0.89 |
| U _K V (μEq/min/kg.bw.) | 1.55 | 1.44 | 1.66 | 1.76 |
| U _{Cl} V (μEq/min/kg.bw.) | 0.17 | 0.25 | 0.29 | 0.26 |
| F.E. of Na (%) | 0.29 | 0.37 | 0.45 | 0.29 |
| F.E. of K (%) | 23.25 | 29.48 | 29.37 | 24.12 |
| F.E.of Cl (%) | 0.08 | 0.14 | 0.15 | 0.13 |
| P _{Osm} (mOsm/Kg.) | 302.2 | 309.6 | 307.4 | 305.4 |
| U _{Osm} (mOsm/Kg.) | 1185.4 | 1022.7 | 1158.1 | 1075.4 |
| U _{Osm} V (μOsm/min/kg.bw.) | 20.38 | 12.38 | 17.64 | 13.44 |
| C _{Osm} (μOsm/min/kg.bw.) | 67.43 | 40.0 | 57.38 | 44.02 |
| C _{H₂O} (μOsm/min/kg.bw.) | - 50.24 | - 27.89 | - 42.15 | - 31.52 |

Table 19 The effects of intravenous injection of 0.1 mg/kg.bw. of Russell's viper venom on cardiovascular and renal parameter of dogs group IV, No 4 weighing 12.0 kgs.

| parameter | control | Post venom injection | | |
|---|---------|----------------------|--------|--------|
| | | 1 hr | 2 hrs | 3 hrs |
| MAP (mm.Hg.) | 100.0 | 88.33 | 115.0 | 115.0 |
| HR (beat/min) | 192 | 174 | 204 | 210 |
| CO (ml/min/kg.bw.) | 357.5 | - | 156.17 | 169.58 |
| PV (ml/kg.bw.) | 76.83 | - | 56.5 | 67.83 |
| BV (ml/kg.bw.) | 102.5 | - | 72.42 | 84.83 |
| PCV (%) | 25 | 22 | 22 | 20 |
| TPR (dyne-sec/cm ⁵) | 1,864 | - | 4,908 | 4,520 |
| RVR (dyne-sec/cm ⁵) | 45,679 | 41,952 | 56,759 | 50,412 |
| V (μL/min/kg.bw.) | 12.08 | 15.0 | 16.0 | 15.83 |
| RPF (ml/min/kg.bw.) | 10.94 | 10.95 | 10.53 | 12.16 |
| RBF (ml/min/kg.bw.) | 14.59 | 14.03 | 13.50 | 15.20 |
| GFR (ml/min/kg.bw.) | 4.68 | 4.17 | 4.55 | 4.48 |
| FF (%) | 42.79 | 38.09 | 43.24 | 36.86 |
| RF (%) | 4.08 | - | 8.65 | 8.97 |
| P _{Na} (mEq/L.) | 139 | 141 | 150 | 149 |
| P _K (mEq/L.) | 3.0 | 3.1 | 3.3 | 3.4 |
| P _{Cl} (mEq/L.) | 118 | 109 | 120 | 105 |
| U _{Na} V (μEq/min/kg.bw.) | 1.91 | 3.12 | 2.67 | 1.96 |
| U _K V (μEq/min/kg.bw.) | 1.96 | 2.43 | 2.17 | 1.95 |
| U _{Cl} V (μEq/min/kg.bw.) | 1.60 | 2.04 | 1.87 | 1.55 |
| F.E. of Na (%) | 0.29 | 0.53 | 0.39 | 0.29 |
| F.E. of K (%) | 13.93 | 18.76 | 14.42 | 12.78 |
| F.E. of Cl (%) | 0.29 | 0.45 | 0.34 | 0.33 |
| P _{Osm} (mOsm/Kg.) | 287.8 | 292.6 | 292.8 | 294 |
| U _{Osm} (mOsm/Kg.) | 1386 | 1360.7 | 1230.6 | 1031.8 |
| U _{Osm} V (μOsm/min/kg.bw.) | 16.74 | 20.41 | 19.69 | 16.33 |
| C _{Osm} (μOsm/min/kg.bw.) | 58.18 | 69.76 | 67.25 | 55.56 |
| C _{H₂O} (μOsm/min/kg.bw.) | 46.10 | 54.76 | 51.25 | 39.73 |

BIOGRAPHY

Miss Somchit Tongyongchai was born on May, 1, 1953 in Ubol - Ratchathanee and graduated with a B.Sc. in Nursing from Khon Kaen University in 1976. At present she is a instructor of Nursing in Sanpasitthiprasong Nursing collage, Ubol Ratchathanee.

