#### REFERENCES

- Arena, F., Horrell, B., Cocke, D., Parmaliana, A., and Giordano, N. (1991)

  Magnesia-supported nickel catalysts: I. Factors affecting the structure and morphological properties. <u>Journal of Catalysis</u>, 132, 58-67.
- Ashcroft, A.T., Cheethan, A.K., Green, M.L.H., and Vernon, P.D.F. (1991) Partial oxidation of methane to synthesis gas using carbon dioxide. <u>Nature</u>, 352, 225-226.
- Bhat, R.N. and Sachtler, W.M.H. (1997) Potential of zeolite supported rhodium catalysts for the CO<sub>2</sub> reforming of CH<sub>4</sub>. Applied Catalysis A: General, 150, 279-296.
- Chang, J.S., Park, S.E., and Chon, H. (1996) Catalytic activity and coke resistance in the carbon dioxide reforming of methane to synthesis gas over zeolite-supported Ni catalysts. Applied Catalysis A: General, 145, 111-124.
- Chang, J.S., Park, S.E., Yoo, J.W., and Park, J.N. (2000) Catalytic behavior of supported KNiCa catalyst and mechanistic consideration for carbon dioxide reforming of methane. <u>Journal of Catalysis</u>, 195, 1–11.
- Cheng, Z., Wu, Q., Li, J., and Zhu, Q. (1996) Effects of promoters and preparation procedures on reforming of methane with carbon dioxide over Ni/Al<sub>2</sub>O<sub>3</sub> catalyst. <u>Catalysis Today</u>, 30, 147-155.
- Dong, W.S., Roh, H.S., Jun, K.W., Park, S.E., and Oh, Y.S. (2002) Methane reforming over Ni/Ce-ZrO<sub>2</sub> catalyst: effect of nickel content. <u>Applied Catalysis A: General</u>, 226, 63-72.
- Edwards, J.H. and Maitra, A.M. (1995) The chemistry of methane reforming with carbon dioxide and its current and potential applications. <u>Fuel Processing</u>
  <u>Technology</u>, 42, 269-289.
- Erdohelyi, A., Cserenyi, J., Papp, E., and Solymosi, F. (1994) Catalytic reaction of methane with carbon dioxide over supported palladium. <u>Applied Catalysis</u>
  <u>A: General</u>, 108, 205-219.
- Gadalla, A.M. and Bower, B. (1988) The role of catalyst support on the activity of nickel for reforming methane with CO<sub>2</sub>. Chemical Engineering Science, 42, 3049-3062.

- Gesser, H.D., Hunter, N.R., Shigapov, A.N., and Januati, V. (1994) Carbon dioxide reforming with methane to CO and H<sub>2</sub> in a hot wire thermal diffusion column (TDC) reactor. Energy & Fuels, 8, 1123-1125.
- Iwamoto, M., Hasuwa, T., Furukawa, H., and Kagawa, S. (1983) Water gas shift reaction catalyzed by metal ion-exchanged zeolites. <u>Journal of Catalysis</u>, 79, 291-297.
- Kuijpers, E., Jansen, J., van Dillen, A.J., and Geus, J.W. (1981) The reversible decomposition of methane on a Ni/SiO<sub>2</sub> catalyst. <u>Journal of Catalysis</u>, 72, 75-82.
- McDaniel, C.V. and Maher, P.K. (1984) Zeolite stability and ultra stable zeolites. Zeolite Chemical Catalysis, 4, 225-232.
- Nakamura, J., Aikawa, K., Sato, K., and Uchijima, T. (1994) Role of support in reforming of CH<sub>4</sub> with CO<sub>2</sub> over Rh catalysts. <u>Catalysis Letters</u>, 25, 265-270.
- Nimwattanakul, W., Luengnaruemitchai, A., and Jitkarnka, S. (2005) Potential of Ni supported on clinoptiolite catalysts for carbon dioxide reforming of methane. <u>International of Hydrogen Energy</u>, *In Press*.
- Parmaliana, A., Arena, F., Frusteri, F., Coluccia, S., Marchese, L., Martra, G., and Chuvilin, A. (1993) Magnesia-supported nickel catalysts: II. Surface properties and reactivity in methane steam Reforming. <u>Journal of Catalysis</u>, 141, 34-47.
- Portugal, U.L., Santos, A.C.S.F., Damyanova, S., Marques, C.M.P., and Buenob, J.M.C. (2002) CO<sub>2</sub> reforming of CH<sub>4</sub> over Rh-containing catalysts.

  <u>Journal of Molecular Catalysis A: Chemical</u>, 184, 311-322.
- Qin, D. and Lapszewicz, J. (1994) Study of mixed steam and CO<sub>2</sub> reforming of CH<sub>4</sub> to syngas on MgO-supported metals. <u>Catalysis Today</u>, 21, 551-560.
- Richardson, J.T. and Paripatyadar, S.A. (1990) Carbon dioxide reforming of methane with supported rhodium. <u>Applied Catalysis</u>, 61(1), 293-309.
- Roh, H.S., Jun, K.W., and Park, S.E. (2003) Methane-reforming reactions over Ni/Ce-ZrO<sub>2</sub>/θ-Al<sub>2</sub>O<sub>3</sub> catalysts. <u>Applied Catalysis A: General</u>, 251, 275-283.

- Roh, H.S., Jun, K.W., Dong, W.S., Chang, J.S., Park, S.E., and Joe, Y.I. (2002) Highly activity and stable Ni/Ce-ZrO<sub>2</sub> catalyst for H<sub>2</sub> production from methane. Journal of Molecular Catalysis A: Chemical, 181, 137-142.
- Rostrup-Nielsen, J.R., in: J. Anderson, M. Boudart (Eds.). (1984) <u>Catalysis Science</u> and <u>Technology</u>, Vol. 5, Springer, New York, 1984.
- Stagg, S.M., Romeo, E., Padro, C., and Resasco, D.E. (1998) Effect of promotion with Sn on supported Pt catalysts for CO<sub>2</sub> reforming of CH<sub>4</sub>. <u>Journal of Catalysis</u>, 178, 137-145.
- Suthupanya, M., Wongkasemjit, S., and Gulari, E. (2003) Zeolite synthesis directly from alumatrane and silarane via sol-gel process and microwave technique. Ph.D. Thesis in Polymer Science Program, The Petroleum and Petrochemical College, Chulalongkorn University.
- Tomishige, K., Kanazawa, S., Ito, S., and Kunimori, K. (2003) Catalyst development for direct heat supply from combustion to reforming in methane reforming with CO<sub>2</sub> and O<sub>2</sub>. <u>Applied Catalysis A: General.</u> 244, 71–82.
- Tsipouriari, V.A., Efstahiou, A.M., and Verykios, X.E. (1994) Reforming of methane with carbon dioxide to synthesis gas over supported Rh catalysts.

  <u>Catalysis Today</u>, 21, 579-587.
- Tsuchida, T. (1993) Preparation of high surface area  $\alpha$ -Al<sub>2</sub>O<sub>3</sub> and its surface properties. Applied Catalysis A: General, 105, 141-149.
- Vernon, P.D.F., Green, M.L.H., Cheethan, A.K., and Ashcroft, A.T. (1992) Partial oxidation of methane to synthesis gas, and carbon dioxide as an oxidizing agent for methane conversion. <u>Catalysis Today</u>, 13, 417-426.
- Wang, S. and Lu, G.Q. (1996) Carbon dioxide reforming of methane to produce synthesis gas over metal-supported catalysts: State of art. Energy & Fuels, 10, 896-904.
- Wang, S. and Lu, G.Q. (1998) Role of CeO<sub>2</sub> in Ni/CeO<sub>2</sub>-Al<sub>2</sub>O<sub>3</sub> catalysts for carbon dioxide reforming of methane. <u>Applied Catalysis B: Environmental</u>, 19, 267-277.

Wang, S. and Lu, G.Q. (1999) A comprehensive study on carbon dioxide reforming of methane over Ni/γ-Al<sub>2</sub>O<sub>3</sub> catalysts. <u>Industrial Engineering Chemistry Research</u>, 38, 2615-2625.

#### **APPENDIX**

### Assumptions, definitions, and calculations.

To facilitate the calculations in this work, some assumptions were made as follows:

- 1. All the gaseous behaviors obey the ideal gas law.
- 2. Pressure drop across the system is very small and can be negligible.
- 3. The pressure in the system equals to the atmospheric pressure.

# 1. Catalyst Preparation

1.1 Amount of Ni loading

Prepared 0.2 g of 8wt%Ni/KH Zeolite

- To prepare 0.2 g of catalyst with 8wt%Ni (MW=58.69 g/mole) need to have 0.2\*8/100 = 0.016 g of Ni
- Amount of Ni(NO<sub>3</sub>)<sub>2</sub>•6H<sub>2</sub>O (MW=290.81 g/mole) 0.016\*290.81/58.69 = 0.07928 g
- Amount of KH Zeolite

$$0.2 - 0.016 = 0.184 g$$

## 2. Conversion and Selectivity

2.1 Methane Conversion

CH<sub>4</sub> conversion = 
$$\frac{CH_{4,in} - CH_{4,out}}{CH_{4,in}} \times 100$$
Where CH<sub>4,in</sub> = CH<sub>4</sub> fed to the reactor
$$CH_{4,out} = CH_4 \text{ left the reactor}$$

## 2.2 Carbon Dioxide Conversion

$$CO_2 \text{ conversion} = \frac{CO_{2,in} - CO_{2,out}}{CO_{2,in}} \times 100$$

Where  $CO_{2,in} = CO_2$  fed to the reactor

$$CO_{2,out} = CO_2$$
 left the reactor

### 2.3 Hydrogen Selectivity

Calculation H<sub>2</sub> selectivity vs CO selectivity:

$$CH_4 + CO_2 \longrightarrow 2H_2 + 2 CO$$

$$H_2 \text{ selectivity} = \frac{\frac{1}{2} * F_{,out} * y_{(H_2,out)}}{F_{,in} * y_{(CH_4,in)} - F_{,out} * y_{(CH_4,out)}} * 100$$

CO selectivity = 
$$\frac{\frac{1}{2} * F_{,out} * y_{(CO,out)}}{F_{,in} * y_{(CH_4,in)} - F_{,out} * y_{(CH_4,out)}} * 100$$

Therefore, selectivity of H<sub>2</sub> vs CO = 
$$\frac{H_2 selectivity}{CO selectivity + H_2 selectivity}*100$$

$$= \frac{\frac{1}{2} * y_{(H_2,out)}}{(\frac{1}{2} * y_{(H_2,out)}) + (\frac{1}{2} * y_{(CO,out)})} *100$$

$$= \frac{y_{(H_2,out)}}{y_{(H_2,out)} + y_{(CO,out)}} *100$$

Where  $F_{,in}$  = Total flow rate of feed stream that fed to the reactor

F<sub>out</sub> = Total flow rate that left the reactor

 $y_{(H_2,out)}$  = Mole fraction of H<sub>2</sub> in the effluent stream

 $y_{(CO,out)}$  = Mole fraction of CO in the effluent stream

 $y_{(CH_4,in)}$  = Mole fraction of CH<sub>4</sub> in the feed stream

 $y_{(CH_4,out)}$  = Mole fraction of CH<sub>4</sub> in the effluent stream

### **CURRICULUM VITAE**

Name: Ms. Athiya Kaengsilalai

Date of Birth: December 28, 1980

Nationality: Thai

## University Education:

1999-2003 Bachelor Degree of Industrial Chemistry, Faculty of Applied Science, King's Mongkut Institute of Technology North Bangkok, Bangkok, Thailand

### Proceedings:

- Kaengsilalai, A., Luengnaruemitchai, A., Jitkarnka, S., and Wongkasemjit, S. (2004, December 1-3) Carbon Dioxide Reforming of Methane over Ni/K-H Zeolite. <u>Proceedings of the RSCE 2004</u>, Bangkok, Thailand.
- Kaengsilalai, A., Luengnaruemitchai, A., Jitkarnka, S., and Wongkasemjit, S. (2004, December 1-3) Hydrogen Production from Carbon Dioxide Reforming of Methane over Ni/Zeolite Catalysts: Effect of an Upscale Synthesis. <u>Proceedings</u> of the Sustainable Energy and Environment (SEE), Hua Hin, Thailand.
- Kaengsilalai, A., Luengnaruemitchai, A., Jitkarnka, S., and Wongkasemjit, S. (2005, July 13-15) Influence of Batch Scale Synthesis and Morphology of KH Zeolite on Hydrogen Production from Carbon Dioxide Reforming of Methane. <u>Proceedings of the International Hydrogen Energy Conference 2005</u>. Istanbul, Turkey.